

NILPOTENT BICONE AND CHARACTERISTIC SUBMODULE OF A REDUCTIVE LIE ALGEBRA

JEAN-YVES CHARBONNEL

ANNE MOREAU

Université Paris 7 - CNRS
Institut de Mathématiques de Jussieu
Théorie des groupes
Case 7012, 2 Place Jussieu
75251 Paris Cedex 05, France
jyc@math.jussieu.fr

ETH Zürich
Departement Mathematik
HG G66.4
Rämistrasse 101
8092 Zürich, Switzerland
anne.moreau@math.ethz.ch

Abstract. Let \mathfrak{g} be a finite-dimensional complex reductive Lie algebra and $S(\mathfrak{g})$ its symmetric algebra. The *nilpotent bicone* of \mathfrak{g} is the subset of elements (x, y) of $\mathfrak{g} \times \mathfrak{g}$ whose subspace generated by x and y is contained in the nilpotent cone. The nilpotent bicone is naturally endowed with a scheme structure, as nullvariety of the augmentation ideal of the subalgebra of $S(\mathfrak{g}) \otimes_{\mathbb{C}} S(\mathfrak{g})$ generated by the 2-order polarizations of invariants of $S(\mathfrak{g})$. The main result of this paper is that the nilpotent bicone is a complete intersection of dimension $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g})$, where $\mathfrak{b}_{\mathfrak{g}}$ and $\text{rk } \mathfrak{g}$ are the dimensions of Borel subalgebras and the rank of \mathfrak{g} , respectively. This affirmatively answers a conjecture of Kraft and Wallach concerning the nullcone [KrW2]. In addition, we introduce and study in this paper the *characteristic submodule* of \mathfrak{g} . The properties of the nilpotent bicone and the characteristic submodule are known to be very important for the understanding of the commuting variety and its ideal of definition. The main difficulty encountered for this work is that the nilpotent bicone is not reduced. To deal with this problem, we introduce an auxiliary reduced variety, *the principal bicone*. The nilpotent bicone, as well as the principal bicone, are linked to jet schemes. We study their dimensions using arguments from motivic integration. Namely, we follow methods developed by Mustață in [Mu]. Finally, we give applications of our results to invariant theory.

0. Introduction

0.1. Main results

Let \mathfrak{g} be a finite-dimensional complex reductive Lie algebra, let G be its adjoint group and let $\mathfrak{N}_{\mathfrak{g}}$ be its nilpotent cone. For (x, y) in $\mathfrak{g} \times \mathfrak{g}$, we denote by $P_{x,y}$ the subspace generated by x and y . A subset of $\mathfrak{g} \times \mathfrak{g}$ is called *bicone* if it is stable under the maps $(x, y) \mapsto (sx, ty)$ where s and t are in \mathbb{C}^* . The *nilpotent bicone* of \mathfrak{g} is the subset

$$\mathcal{N}_{\mathfrak{g}} := \{(x, y) \in \mathfrak{g} \times \mathfrak{g} \mid P_{x,y} \subset \mathfrak{N}_{\mathfrak{g}}\}.$$

Since $\mathfrak{N}_{\mathfrak{g}}$ is a G -invariant closed cone of \mathfrak{g} , the subset $\mathcal{N}_{\mathfrak{g}}$ is a closed bicone of $\mathfrak{g} \times \mathfrak{g}$, invariant under the diagonal action of G on $\mathfrak{g} \times \mathfrak{g}$.

DOI: 10.1007/s00031-008-

Received February 23, 2008. Accepted August 9, 2008.

Let $S(\mathfrak{g})$ be the symmetric algebra of \mathfrak{g} and let $S(\mathfrak{g})^{\mathfrak{g}}$ be the subalgebra of G -invariant elements of $S(\mathfrak{g})$. According to a Chevalley result, the algebra $S(\mathfrak{g})^{\mathfrak{g}}$ is polynomial in $\mathrm{rk} \mathfrak{g}$ variables, where $\mathrm{rk} \mathfrak{g}$ is the rank of \mathfrak{g} . Let $p_1, \dots, p_{\mathrm{rk} \mathfrak{g}}$ be homogeneous generators of $S(\mathfrak{g})^{\mathfrak{g}}$ of degrees $d_1, \dots, d_{\mathrm{rk} \mathfrak{g}}$ respectively. We can suppose that the sequence $d_1, \dots, d_{\mathrm{rk} \mathfrak{g}}$ is weakly increasing. For $i = 1, \dots, \mathrm{rk} \mathfrak{g}$, the 2-order polarizations $p_{i,m,n}$ of p_i are the unique elements of $(S(\mathfrak{g}) \otimes_{\mathbb{C}} S(\mathfrak{g}))^{\mathfrak{g}}$ satisfying the following relation:

$$p_i(ax + by) = \sum_{m+n=d_i} a^m b^n p_{i,m,n}(x, y), \quad (1)$$

for all (a, b) in \mathbb{C}^2 and (x, y) in $\mathfrak{g} \times \mathfrak{g}$. The nilpotent cone $\mathfrak{N}_{\mathfrak{g}}$ is the nullvariety in \mathfrak{g} of the ideal generated by the polynomials $p_1, \dots, p_{\mathrm{rk} \mathfrak{g}}$. Therefore, according to relation (1), $\mathcal{N}_{\mathfrak{g}}$ is the nullvariety in $\mathfrak{g} \times \mathfrak{g}$ of the ideal of $S(\mathfrak{g}) \otimes_{\mathbb{C}} S(\mathfrak{g})$ generated by the polynomials $p_{i,m,n}$ for $i = 1, \dots, \mathrm{rk} \mathfrak{g}$ and $m + n = d_i$. Thus, $\mathcal{N}_{\mathfrak{g}}$ is naturally endowed with a scheme structure. From here, we study $\mathcal{N}_{\mathfrak{g}}$ as the subscheme of $\mathfrak{g} \times \mathfrak{g}$ corresponding to the ideal generated by the polynomials $p_{i,m,n}$ for $i = 1, \dots, \mathrm{rk} \mathfrak{g}$ and $m + n = d_i$.

Let $b_{\mathfrak{g}}$ be the dimension of a Borel subalgebra of \mathfrak{g} . By a classical result [Bou], we have $d_1 + \dots + d_{\mathrm{rk} \mathfrak{g}} = b_{\mathfrak{g}}$. So $\mathcal{N}_{\mathfrak{g}}$ is the nullvariety in $\mathfrak{g} \times \mathfrak{g}$ of $b_{\mathfrak{g}} + \mathrm{rk} \mathfrak{g}$ polynomial functions. As a result, the dimension of any irreducible component of $\mathcal{N}_{\mathfrak{g}}$ is at least $3(b_{\mathfrak{g}} - \mathrm{rk} \mathfrak{g})$, since $\mathfrak{g} \times \mathfrak{g}$ has dimension $2(b_{\mathfrak{g}} - \mathrm{rk} \mathfrak{g})$. The main result of this paper is the following theorem (see Theorem 33(iv), Proposition 36(ii), and Theorem 45).

Theorem 1. *The nilpotent bicone is a nonreduced complete intersection in $\mathfrak{g} \times \mathfrak{g}$ of dimension $3(b_{\mathfrak{g}} - \mathrm{rk} \mathfrak{g})$. Moreover, the images of any irreducible component of $\mathcal{N}_{\mathfrak{g}}$ by the first and second projections from $\mathfrak{g} \times \mathfrak{g}$ to \mathfrak{g} are equal to $\mathfrak{N}_{\mathfrak{g}}$.*

This result affirmatively answers a conjecture [KrW2, Sect. 1] of Kraft and Wallach concerning the *nullcone* of the G -module $\mathfrak{g} \times \mathfrak{g}$ (see Theorem 41). Clearly, the images of $\mathcal{N}_{\mathfrak{g}}$ by the first and second projections from $\mathfrak{g} \times \mathfrak{g}$ to \mathfrak{g} are equal to $\mathfrak{N}_{\mathfrak{g}}$. Theorem 1 specifies that this is true for any irreducible component of $\mathcal{N}_{\mathfrak{g}}$. Notice that this statement is a much stronger result.

We introduce in addition in this paper the *characteristic submodule* of \mathfrak{g} . It is an $S(\mathfrak{g}) \otimes_{\mathbb{C}} S(\mathfrak{g})$ -submodule of $S(\mathfrak{g}) \otimes_{\mathbb{C}} S(\mathfrak{g}) \otimes_{\mathbb{C}} \mathfrak{g}$. The properties of the nilpotent bicone and the characteristic submodule are known to be very important for the understanding of the commuting variety. Recall that the *commuting variety* $\mathcal{C}_{\mathfrak{g}}$ of \mathfrak{g} is the set of elements (x, y) of $\mathfrak{g} \times \mathfrak{g}$ such that $[x, y] = 0$. The commuting variety has been studied for many years. According to a result of Richardson [Ri1], $\mathcal{C}_{\mathfrak{g}}$ is irreducible. Moreover, $\mathcal{C}_{\mathfrak{g}}$ is the nullvariety of the ideal generated by the elements $(x, y) \mapsto \langle v, [x, y] \rangle$, where v runs through \mathfrak{g} . An old unsolved question is to know whether this ideal is prime [LS]. In other words, we want to know if this ideal is the ideal of definition of $\mathcal{C}_{\mathfrak{g}}$ since $\mathcal{C}_{\mathfrak{g}}$ is irreducible. The study of the commuting variety and of its ideal of definition is a main motivation for our work.

0.2. Description of the paper

We denote by $\mathfrak{g}_{\mathrm{reg}}$ the subset of regular elements of \mathfrak{g} and we set

$$\Omega_{\mathfrak{g}} := \{(x, y) \in \mathfrak{g} \times \mathfrak{g} \mid P_{x,y} \setminus \{0\} \subset \mathfrak{g}_{\mathrm{reg}}, \dim P_{x,y} = 2\}.$$

NILPOTENT BICONE

The properties of the subset $\Omega_{\mathfrak{g}}$ were studied by Bolsinov in [Bol] and recently by Panyushev and Yakimova in [PYa]. The set of smooth points of $\mathcal{N}_{\mathfrak{g}}$ with respect to its scheme structure is the set of elements $(x, y) \in \mathfrak{g} \times \mathfrak{g}$ at which the differentials of the $p_{i,m,n}$, for $i = 1, \dots, \text{rk } \mathfrak{g}$ and $m + n = d_i$, are linearly independent. The set $\Omega_{\mathfrak{g}}$ is an open subset of $\mathfrak{g} \times \mathfrak{g}$ and its intersection with the nilpotent bicone turns out to be the set of smooth points of $\mathcal{N}_{\mathfrak{g}}$ (see Proposition 13 and Remark 3). The description of this subset is therefore very important for us. We give various properties of the subset $\Omega_{\mathfrak{g}}$ in Section 1. Next, we introduce in this section the *characteristic submodule* $B_{\mathfrak{g}}$ of \mathfrak{g} (see Definition 1) whose study is heavily related to $\Omega_{\mathfrak{g}}$. The characteristic submodule was introduced by the first author a few years ago in a paper on the commuting variety. We prove here that $B_{\mathfrak{g}}$ is a free $S(\mathfrak{g}) \otimes_{\mathbb{C}} S(\mathfrak{g})$ -module of rank $b_{\mathfrak{g}}$, and we provide an explicit basis for $B_{\mathfrak{g}}$ (see Theorem 11). Even if this result is not directly useful for the nilpotent bicone, it could be interesting in itself.

The results obtained in Section 1 concerning $\Omega_{\mathfrak{g}}$ do not enable us to provide “enough” smooth points of $\mathcal{N}_{\mathfrak{g}}$ to apply the criterion of Kostant [Kos2] (see Remark 3). Actually, the intersection of $\mathcal{N}_{\mathfrak{g}}$ and $\Omega_{\mathfrak{g}}$ is an empty set in many cases. This observation makes the study of the scheme $\mathcal{N}_{\mathfrak{g}}$ very difficult. In order to deal with this problem, we introduce an auxiliary reduced scheme, the *principal bicone* of \mathfrak{g} ,

$$\mathcal{X}_{\mathfrak{g}} := \{(x, y) \in \mathfrak{g} \times \mathfrak{g} \mid P_{x,y} \subset \mathfrak{X}_{\mathfrak{g}}\},$$

where $\mathfrak{X}_{\mathfrak{g}}$ is the *principal cone* of \mathfrak{g} , that is the Zariski closure of the set of principal semisimple elements (see Definitions 2 and 3). The study of the varieties $\mathfrak{X}_{\mathfrak{g}}$ and $\mathcal{X}_{\mathfrak{g}}$ is the main topic of Section 2.

We observe that $\mathcal{N}_{\mathfrak{g}}$ and $\mathcal{X}_{\mathfrak{g}}$ can be identified with constructible subsets of jet schemes of $\mathfrak{N}_{\mathfrak{g}}$ and $\mathfrak{X}_{\mathfrak{g}}$ respectively. In [Mu], Mustața uses the theory of motivic integration, as developed by Kontsevich [Kon], Denef and Loeser [DL2], and Batyrev [Ba], to prove a result concerning the jet schemes of locally complete intersections. In particular, his result can be applied to $\mathfrak{N}_{\mathfrak{g}}$ and $\mathfrak{X}_{\mathfrak{g}}$. Thus, in the Appendix of [Mu], Eisenbud and Frenkel extend results of Kostant concerning the nilpotent cone of a reductive Lie algebra in the setting of jet schemes. In Section 3, after reviewing some facts about motivic integration, we state technical results concerning motivic integrals, following methods developed by Mustața in [Mu], which will be useful for Section 4. Using all of this, we prove in Section 4 the following theorem (see Theorem 33(i) and Proposition 36(i)).

Theorem 2. *The principal bicone of \mathfrak{g} is a reduced complete intersection of dimension $3(b_{\mathfrak{g}} - \text{rk } \mathfrak{g} + 1)$. Moreover, the images of any irreducible component of $\mathcal{X}_{\mathfrak{g}}$ by the first and second projection from $\mathfrak{g} \times \mathfrak{g}$ to \mathfrak{g} are equal to the principal cone $\mathfrak{X}_{\mathfrak{g}}$.*

Then, we deduce Theorem 1 from Theorem 2. We give in Section 5 applications of our results to invariant theory, mainly in relation with the nullcone of the G -module $\mathfrak{g} \times \mathfrak{g}$. In Section 6 we obtain additional properties about the irreducible components of the nilpotent bicone. We prove that $\mathcal{N}_{\mathfrak{g}}$ is a nonreduced scheme (see Theorem 45) and we give a lower bound for the number of irreducible components of $\mathcal{N}_{\mathfrak{g}}$ (see Proposition 52).

We note that Section 3 is entirely independent of Sections 1 and 2 and does not deal with Lie algebras.

0.3. Additional notations and conventions

In this paper, the ground field is \mathbb{C} . All topological terms refer to the Zariski topology. If X is an algebraic variety, an open subset of X whose complementary in X has codimension at least 2 is called a *big open subset* of X .

The set of integers, the set of nonnegative integers and the set of positive integers are denoted by \mathbb{Z} , \mathbb{N} and \mathbb{N}^* respectively. As usual, the subset of nonzero elements of \mathbb{C} is denoted by \mathbb{C}^* .

If E is a finite set, its cardinality is denoted by $|E|$.

For x in \mathfrak{g} , we denote by $\mathfrak{g}(x)$ the centralizer of x in \mathfrak{g} . Thus x belongs to $\mathfrak{g}_{\text{reg}}$ if and only if $\mathfrak{g}(x)$ has dimension $\text{rk } \mathfrak{g}$.

The action of G in $\mathfrak{g} \times \mathfrak{g}$ will always be the diagonal action. For x in \mathfrak{g} or in $\mathfrak{g} \times \mathfrak{g}$, we denote by $G.x$ its G -orbit under the corresponding G -action.

If \mathfrak{g} is commutative, then $\mathcal{N}_{\mathfrak{g}}$ is reduced to $\{(0, 0)\}$. Henceforth, we suppose that \mathfrak{g} is not commutative.

Let $\mathfrak{g}_1, \dots, \mathfrak{g}_m$ be the simple factors of \mathfrak{g} . Since the nilpotent cone $\mathfrak{N}_{\mathfrak{g}}$ is the product of the nilpotent cones $\mathfrak{N}_{\mathfrak{g}_1}, \dots, \mathfrak{N}_{\mathfrak{g}_m}$, $\mathcal{N}_{\mathfrak{g}}$ is the product of the nilpotent bicones $\mathcal{N}_{\mathfrak{g}_1}, \dots, \mathcal{N}_{\mathfrak{g}_m}$. Furthermore, as the equality

$$3(\text{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g}) = 3(\text{b}_{\mathfrak{g}_1} - \text{rk } \mathfrak{g}_1) + \dots + 3(\text{b}_{\mathfrak{g}_m} - \text{rk } \mathfrak{g}_m)$$

holds, it suffices to prove Theorem 1 in the case where \mathfrak{g} is simple. It will be sometimes useful to make the assumption that \mathfrak{g} is simple (see, e.g., Sections 2, 5 and 6).

The first and second projections from $\mathfrak{g} \times \mathfrak{g}$ to \mathfrak{g} are denoted by ϖ_1 and ϖ_2 respectively. The following lemma will be very handy throughout the paper.

Lemma 3. *Let Y be a G -invariant closed bicone in $\mathfrak{g} \times \mathfrak{g}$. Then the subsets $\varpi_1(Y)$ and $\varpi_2(Y)$ are G -invariant closed cones of \mathfrak{g} .*

Proof. As Y is a G -invariant bicone, $\varpi_1(Y)$ and $\varpi_2(Y)$ are G -invariant cones of \mathfrak{g} . Moreover, $\varpi_1(Y) \times \{0\}$ and $\{0\} \times \varpi_2(Y)$ are the intersections of Y with $\mathfrak{g} \times \{0\}$ and $\{0\} \times \mathfrak{g}$ respectively, since Y is a closed bicone. So $\varpi_1(Y)$ and $\varpi_2(Y)$ are closed subsets of \mathfrak{g} . \square

We fix a principal \mathfrak{sl}_2 -triple (e, h, f) of \mathfrak{g} . Thus, the following relations:

$$[h, e] = 2e, \quad [e, f] = h, \quad [h, f] = -2f,$$

are satisfied, e and f are regular nilpotent elements and h is a regular semisimple element of \mathfrak{g} . If ad is the adjoint representation of \mathfrak{g} , then $\text{ad } h$ induces a \mathbb{Z} -grading on \mathfrak{g} and we have $\mathfrak{g} = \sum_{i \in \mathbb{Z}} \mathfrak{g}_i$, where \mathfrak{g}_i is the i -eigenspace of $\text{ad } h$. Moreover, all the eigenvalues of $\text{ad } h$ are even integers. The centralizer $\mathfrak{g}(h) = \mathfrak{g}_0$ of h in \mathfrak{g} is a Cartan subalgebra, that we will also denote by \mathfrak{h} . The unique Borel subalgebra containing e is

$$\mathfrak{b} := \mathfrak{h} \oplus \sum_{i > 0} \mathfrak{g}_i,$$

NILPOTENT BICONE

and the nilpotent radical of \mathfrak{b} is $\mathfrak{u} := \sum_{i>0} \mathfrak{g}_i$. We additionally set

$$\mathfrak{b}_- := \sum_{i \leq 0} \mathfrak{g}_i, \quad \mathfrak{u}_- := \sum_{i < 0} \mathfrak{g}_i.$$

Let \mathbf{B} and \mathbf{B}_- be the normalizers of \mathfrak{b} and \mathfrak{b}_- in G and let \mathbf{H} and $N_G(\mathfrak{h})$ be the centralizer and the normalizer of \mathfrak{h} in G . Then the quotient $W_G(\mathfrak{h})$ of $N_G(\mathfrak{h})$ by \mathbf{H} is the Weyl group of \mathfrak{g} with respect to \mathfrak{h} .

Let \mathcal{R} be the root system of $(\mathfrak{g}, \mathfrak{h})$, let \mathcal{R}_+ be the positive root system of \mathcal{R} defined by \mathfrak{b} and let Π be the basis of \mathcal{R}_+ . For any α in \mathcal{R} , we denote by \mathfrak{g}^α the corresponding root subspace. Denote by w_0 the longest element of $W_G(\mathfrak{h})$ with respect to Π . Then $w_0(\Pi)$ is equal to $-\Pi$ and for any representative g_0 of w_0 in $N_G(\mathfrak{h})$, $g_0(\mathfrak{b})$ is equal to \mathfrak{b}_- .

Let $\langle \cdot, \cdot \rangle$ be a nondegenerate G -invariant bilinear form on $\mathfrak{g} \times \mathfrak{g}$ which extends the Killing form of the semisimple part of \mathfrak{g} . In the remainder of this paper, the orthogonality will refer to $\langle \cdot, \cdot \rangle$.

Acknowledgments. We are grateful to Michel Duflo for bringing [Mu] to our attention. We would also like to thank François Loeser for his e-mails concerning motivic integration questions, and Karin Baur for her comments. In addition, we thank Vladimir Popov for his support and the referees for their numerous and judicious advice and their careful attention to our paper.

1. Characteristic submodule

In this section we introduce the characteristic submodule of \mathfrak{g} and we describe some of its properties.

1.1. Preliminaries

Let us recall that $p_1, \dots, p_{\text{rk } \mathfrak{g}}$ are homogeneous generators of $S(\mathfrak{g})^{\mathfrak{g}}$ of degrees $d_1, \dots, d_{\text{rk } \mathfrak{g}}$ respectively such that the sequence $d_1, \dots, d_{\text{rk } \mathfrak{g}}$ is weakly increasing. For $i = 1, \dots, \text{rk } \mathfrak{g}$, let ε_i be the element of $S(\mathfrak{g}) \otimes_{\mathbb{C}} \mathfrak{g}$ defined by the following relation:

$$\langle \varepsilon_i(x), v \rangle = p'_i(x)(v),$$

for x, v in \mathfrak{g} , where $p'_i(x)$ is the differential of p_i at x . The first statement of the following lemma comes from [Ri3, Lemma 2.1] while the second statement comes from [Kos2, Theorem 9].

Lemma 4. *Let x be in \mathfrak{g} .*

- (i) *For $i = 1, \dots, \text{rk } \mathfrak{g}$, $\varepsilon_i(x)$ belongs to the centre of $\mathfrak{g}(x)$.*
- (ii) *The elements $\varepsilon_1(x), \dots, \varepsilon_{\text{rk } \mathfrak{g}}(x)$ are linearly independent in \mathfrak{g} if and only if x is regular. Moreover, if so, $\varepsilon_1(x), \dots, \varepsilon_{\text{rk } \mathfrak{g}}(x)$ is a basis of $\mathfrak{g}(x)$.*

Let i be in $\{1, \dots, \text{rk } \mathfrak{g}\}$. The 2-order polarizations $p_{i,m,n}$ of p_i are defined by the following relation:

$$p_i(ax + by) = \sum_{m+n=d_i} a^m b^n p_{i,m,n}(x, y),$$

for any (a, b) in \mathbb{C}^2 and any (x, y) in $\mathfrak{g} \times \mathfrak{g}$. For $(m, n) \in \mathbb{N}^2$ such that $m + n = d_i$, let $\varepsilon_{i,m,n}$ be the element of $S(\mathfrak{g}) \otimes_{\mathbb{C}} S(\mathfrak{g}) \otimes_{\mathbb{C}} \mathfrak{g}$ such that the linear functional $v \mapsto \langle \varepsilon_{i,m,n}(x, y), v \rangle$ on \mathfrak{g} is the differential at x of the function $x \mapsto p_{i,m,n}(x, y)$ for (x, y) in $\mathfrak{g} \times \mathfrak{g}$. In particular, $\varepsilon_{i,0,d_i} = 0$ for $i = 1, \dots, \text{rk } \mathfrak{g}$. In addition, since $d_1 + \dots + d_{\text{rk } \mathfrak{g}}$ is equal to $\mathfrak{b}_{\mathfrak{g}}$, the cardinality of the family $\{\varepsilon_{i,1,d_i-1}, \dots, \varepsilon_{i,d_i,0} \mid i = 1, \dots, \text{rk } \mathfrak{g}\}$ is equal to $\mathfrak{b}_{\mathfrak{g}}$.

Lemma 5. *For $i = 1, \dots, \text{rk } \mathfrak{g}$ and $(x, y) \in \mathfrak{g} \times \mathfrak{g}$, we have*

$$\varepsilon_i(ax + by) = \sum_{m=1}^{d_i} a^{m-1} b^{d_i-m} \varepsilon_{i,m,d_i-m}(x, y),$$

for any (a, b) in \mathbb{C}^2 . In particular, $\varepsilon_{i,1,d_i-1}(x, y) = \varepsilon_i(y)$ and $\varepsilon_{i,d_i,0}(x, y) = \varepsilon_i(x)$.

Proof. Let v be in \mathfrak{g} . For any (a, b) in $\mathbb{C}^* \times \mathbb{C}$, we have

$$\begin{aligned} \langle \varepsilon_i(ax + by), v \rangle &= \left. \frac{d}{dt} p_i(ax + by + tv) \right|_{t=0} \\ &= \sum_{m+n=d_i} a^m b^n \left. \frac{d}{dt} p_{i,m,n}(x + ta^{-1}v, y) \right|_{t=0} \\ &= \sum_{m+n=d_i} a^m b^n \langle \varepsilon_{i,m,n}(x, y), a^{-1}v \rangle, \end{aligned}$$

whence the lemma, since $\langle \cdot, \cdot \rangle$ is a nondegenerate G -invariant bilinear form on $\mathfrak{g} \times \mathfrak{g}$. \square

Recall that we have introduced the subset

$$\Omega_{\mathfrak{g}} := \{(x, y) \in \mathfrak{g} \times \mathfrak{g} \mid P_{x,y} \setminus \{0\} \subset \mathfrak{g}_{\text{reg}}, \dim P_{x,y} = 2\},$$

which is clearly G -invariant. We denote by \mathfrak{h}' , \mathfrak{g}'_2 and \mathfrak{g}'_{-2} the intersections of $\mathfrak{g}_{\text{reg}}$ with \mathfrak{h} , \mathfrak{g}_2 and \mathfrak{g}_{-2} respectively. Set

$$\mathfrak{u}_+ := \sum_{i \geq 4} \mathfrak{g}_i.$$

In the following lemma, we explicitly provide elements of $\Omega_{\mathfrak{g}}$.

Lemma 6.

- (i) *If $x \in \mathfrak{h}' + \mathfrak{u}$ and $y \in \mathfrak{g}'_2 + \mathfrak{u}_+$, then $(x, y) \in \Omega_{\mathfrak{g}}$.*
- (ii) *If $x \in \mathfrak{g}'_{-2}$ and $y \in \mathfrak{g}'_2$, then $(x, y) \in \Omega_{\mathfrak{g}}$.*

Proof. (i) The elements x and y are clearly linearly independent. In addition, for any t in \mathbb{C} , $x + ty$ is conjugate to x under G . So $x + ty$ is regular for any t in \mathbb{C} . Since y is regular, (x, y) belongs to $\Omega_{\mathfrak{g}}$.

(ii) The elements x and y are clearly linearly independent. As x is regular, $x + sy$ is regular for any s in an open subset of \mathbb{C} containing 0. Let $t \mapsto g(t)$ be the one-parameter subgroup of \mathbf{H} generated by $\text{ad } h$. As $\mathfrak{g}_{\text{reg}}$ is G -invariant, for any t in \mathbb{C}^* , $g(t)(x + sy)$ is regular for any s in an open subset of \mathbb{C} containing 0. So, from the relations $[h, x] = -2x$ and $[h, y] = 2y$, we deduce that for any t in \mathbb{C}^* , $t^{-2}x + st^2y$ is regular for any s in an open subset of \mathbb{C} containing 0. As $\mathfrak{g}_{\text{reg}}$ is an open cone, $x + st^4y$ is consequently regular for any s in an open subset of \mathbb{C} containing $\{0\}$. So $x + ty$ is regular for any t in \mathbb{C} . As y is regular, we deduce that (x, y) belongs to $\Omega_{\mathfrak{g}}$. \square

NILPOTENT BICONE

Remark 1. The subset $\Omega_{\mathfrak{g}}$ is stable under the involution $(x, y) \mapsto (y, x)$. So, by Lemma 6, the images of the subsets $(\mathfrak{h}' + \mathfrak{u}) \times (\mathfrak{g}'_2 + \mathfrak{u}_+)$ and $\mathfrak{g}'_{-2} \times \mathfrak{g}'_2$ by this involution are contained in $\Omega_{\mathfrak{g}}$. For example, the elements (e, h) , (f, h) and (e, f) are in $\Omega_{\mathfrak{g}}$.

The following lemma is well known.

Lemma 7. *Let x be in $\mathfrak{g}_{\text{reg}}$.*

- (i) *If x belongs to \mathfrak{b} , then $\mathfrak{g}(x)$ is contained in \mathfrak{b} .*
- (ii) *If x belongs to \mathfrak{u} , then $\mathfrak{g}(x)$ is contained in \mathfrak{u} .*

For (x, y) in $\mathfrak{g} \times \mathfrak{g}$, we denote by $\mathfrak{V}(x, y)$ the subspace generated by the elements

$$\varepsilon_{i,1,d_i-1}(x, y), \dots, \varepsilon_{i,d_i,0}(x, y), \quad i = 1, \dots, \text{rk } \mathfrak{g},$$

and we set

$$\mathfrak{V}'(x, y) := \sum_{(a,b) \in \mathbb{C}^2 \setminus (0,0)} \mathfrak{g}(ax + by).$$

We collect in the following lemma some results concerning the subset $\Omega_{\mathfrak{g}}$ in part obtained by Bolsinov in [Bol] and recently by Panyushev and Yakimova in [PYa].

Lemma 8. *Let (x, y) be in $\mathfrak{g} \times \mathfrak{g}$.*

- (i) *The subspace $\mathfrak{V}(x, y)$ has dimension at most $b_{\mathfrak{g}}$. Moreover, it has dimension $b_{\mathfrak{g}}$ if and only if (x, y) belongs to $\Omega_{\mathfrak{g}}$.*
- (ii) *The subspace $\mathfrak{V}(x, y)$ is contained in $\mathfrak{V}'(x, y)$. Moreover, the equality occurs when (x, y) belongs to $\Omega_{\mathfrak{g}}$.*
- (iii) *If (x, y) belongs to $\mathfrak{b} \times \mathfrak{b}$, then $\mathfrak{V}(x, y)$ is contained in \mathfrak{b} .*

Proof. (i) As the sum of $d_1, \dots, d_{\text{rk } \mathfrak{g}}$ is equal to $b_{\mathfrak{g}}$, $\mathfrak{V}(x, y)$ has dimension at most $b_{\mathfrak{g}}$. Let us suppose that (x, y) belongs to $\Omega_{\mathfrak{g}}$. Then by [PYa, Theorem 2.4], the subspace $\mathfrak{V}(x, y)$ has dimension $b_{\mathfrak{g}}$. Conversely, let us suppose that $\mathfrak{V}(x, y)$ has dimension $b_{\mathfrak{g}}$. In particular, $\varepsilon_1(x), \dots, \varepsilon_{\text{rk } \mathfrak{g}}(x)$ are linearly independent. Hence, by Lemma 4(ii), x is regular. By Lemma 5, for any (a, b) in $\mathbb{C}^2 \setminus \{(0, 0)\}$, $\mathfrak{V}(x, y)$ is equal to $\mathfrak{V}(ax + by, y)$. Hence (x, y) belongs to $\Omega_{\mathfrak{g}}$.

(ii) By Lemma 4, $\mathfrak{V}'(x, y)$ contains $\varepsilon_i(ax + by)$ for $i = 1, \dots, \text{rk } \mathfrak{g}$ and any (a, b) in \mathbb{C}^2 . Hence by Lemma 5, $\mathfrak{V}'(x, y)$ contains $\mathfrak{V}(x, y)$. Moreover, by (i) and Lemma 4, $\mathfrak{V}(x, y)$ is equal to $\mathfrak{V}'(x, y)$ when (x, y) belongs to $\Omega_{\mathfrak{g}}$.

(iii) Let us suppose that (x, y) belongs to $\mathfrak{b} \times \mathfrak{b}$. By Lemma 6 and Remark 1, the intersection of $\Omega_{\mathfrak{g}}$ and $\mathfrak{b} \times \mathfrak{b}$ is a nonempty open subset. Moreover, when (x, y) belongs to this intersection, for any (a, b) in $\mathbb{C}^2 \setminus \{(0, 0)\}$, $\mathfrak{g}(ax + by)$ is contained in \mathfrak{b} , by Lemmas 4(ii) and 7(i). Hence for $i = 1, \dots, \text{rk } \mathfrak{g}$ and $m = 1, \dots, d_i$, $\varepsilon_{i,m,d_i-m}(x, y)$ belongs to \mathfrak{b} , by (ii). So $\mathfrak{V}(x, y)$ is contained in \mathfrak{b} . \square

1.2. Closed irreducible subsets of $\mathfrak{s} \times \mathfrak{s}$, invariant under the actions of \mathbf{S} and $\mathbf{GL}_2(\mathbb{C})$

The following automorphisms give an action of $\mathbf{GL}_2(\mathbb{C})$ in $\mathfrak{g} \times \mathfrak{g}$:

$$(x, y) \mapsto (ax + by, cx + dy), \quad \text{where } g = \begin{bmatrix} a & b \\ c & d \end{bmatrix}.$$

Let \mathfrak{s} be the subspace of \mathfrak{g} generated by e, h, f so that \mathfrak{s} is isomorphic to $\mathfrak{sl}_2(\mathbb{C})$. Let \mathbf{S} be the closed connected subgroup of G whose Lie algebra is $\mathfrak{ad}\mathfrak{s}$. We start this subsection by describing the closed irreducible subsets of $\mathfrak{s} \times \mathfrak{s}$ invariant under the actions of \mathbf{S} and $\mathrm{GL}_2(\mathbb{C})$. This will be used in Corollary 10 and Lemma 18.

Let T_3 be the closed bicone of $\mathfrak{s} \times \mathfrak{s}$ generated by the diagonal of $\mathfrak{N}_{\mathfrak{s}} \times \mathfrak{N}_{\mathfrak{s}}$, let T_4 be the closed bicone of $\mathfrak{s} \times \mathfrak{s}$ generated by the diagonal of $\mathfrak{s} \times \mathfrak{s}$ and let T_5 be the subset of elements (x, y) of $\mathfrak{s} \times \mathfrak{s}$ such that x and y belong to the same Borel subalgebra of \mathfrak{s} . As defined, the sets T_3, T_4 and T_5 are irreducible closed subsets of $\mathfrak{s} \times \mathfrak{s}$, invariant under the actions of \mathbf{S} and $\mathrm{GL}_2(\mathbb{C})$. Moreover, they have dimensions 3, 4 and 5 respectively. The verification of these claims is left to the reader.

Lemma 9. *The subsets $\{0\}, T_3, T_4, T_5$, and $\mathfrak{s} \times \mathfrak{s}$ are the only nonempty irreducible closed subsets of $\mathfrak{s} \times \mathfrak{s}$, invariant under the actions of \mathbf{S} and $\mathrm{GL}_2(\mathbb{C})$.*

Proof. Let us first remark that $\mathfrak{N}_{\mathfrak{s}}$ is the only proper \mathbf{S} -invariant closed cone of \mathfrak{s} . Let T be a nonempty irreducible closed subset of $\mathfrak{s} \times \mathfrak{s}$, invariant under the actions of \mathbf{S} and $\mathrm{GL}_2(\mathbb{C})$. By Lemma 3 and the preceding remark, if $0 < \dim T < 4$, then $\varpi_1(T)$ is equal to $\mathfrak{N}_{\mathfrak{s}}$, whence $T = T_3$ by $\mathrm{GL}_2(\mathbb{C})$ -invariance. We assume now $\dim T \geq 4$. As $\varpi_1(T) \supset \mathfrak{N}_{\mathfrak{s}}$ and $\dim T \geq 4$, T contains an element $(e, ae + bf + ch)$, with b or c different from 0. Hence T contains $(e, bf + ch)$. If $c \neq 0$, then $\varpi_1(T)$ contains $bf + ch$ which is semisimple. Otherwise, T contains (e, f) and $\varpi_1(T)$ contains $e + f$ which is semisimple, too. So, in any case, $\varpi_1(T) = \mathfrak{s}$. In particular, $T \supset T_4$ and the equality holds as soon as $\dim T = 4$. Finally, let us suppose $\dim T \geq 5$. As $\varpi_1(T) = \mathfrak{s}$, there exist a, b, c in \mathbb{C} such that $(h, ae + bf + ch)$ belongs to T with a or b different from 0. If $ab = 0$, using the invariance of T under \mathbf{S} and $\mathrm{GL}_2(\mathbb{C})$, we deduce that $\{h\} \times \mathfrak{b}_{\mathfrak{s}}$ is contained in T . In this case, T contains T_5 . If $ab \neq 0$, using the invariance of T under $\mathrm{GL}_2(\mathbb{C})$ and the one-parameter subgroup of \mathbf{H} generated by $\mathrm{ad}h$, we deduce that $(h, at^2e + bt^{-2}f)$ belongs to T for any t in \mathbb{C}^* . So $\{h\} \times \mathfrak{s}$ is contained in T and T has dimension 6, whence the lemma. \square

Corollary 10. *The subset $\Omega_{\mathfrak{g}}$ is a big open subset of $\mathfrak{g} \times \mathfrak{g}$.*

Proof. Suppose that $\Omega_{\mathfrak{g}}$ is not a big open subset of $\mathfrak{g} \times \mathfrak{g}$. Then, the complementary of $\Omega_{\mathfrak{g}}$ in $\mathfrak{g} \times \mathfrak{g}$ has an irreducible component Σ of codimension 1 in $\mathfrak{g} \times \mathfrak{g}$. As $\Omega_{\mathfrak{g}}$ is invariant under the action of G and $\mathrm{GL}_2(\mathbb{C})$, Σ is invariant under these actions too. The intersection T of Σ and $\mathfrak{s} \times \mathfrak{s}$ contains $(0, 0)$. So T is a nonempty closed cone of $\mathfrak{s} \times \mathfrak{s}$, invariant under the actions of \mathbf{S} and $\mathrm{GL}_2(\mathbb{C})$. As Σ is a hypersurface of $\mathfrak{g} \times \mathfrak{g}$, T has codimension at most 1 in $\mathfrak{s} \times \mathfrak{s}$. Hence Lemma 9(ii), implies that (h, e) belongs to T . But, by Lemma 6(i), (h, e) belongs to $\Omega_{\mathfrak{g}}$, whence the expected contradiction. \square

Remark 2. We can also deduce Corollary 10 from [PYa]. Indeed, according to [PYa, Lemma 3.1], for any $x \in \mathfrak{g}_{\mathrm{reg}}$, the set $\{y \in \mathfrak{g} \mid (x, y) \in \Omega_{\mathfrak{g}}\}$ is a big open subset of \mathfrak{g} , whence we can readily deduce Corollary 10.

1.3. Characteristic submodule

By a result of Dixmier [Di, §2], the $S(\mathfrak{g})$ -submodule of elements φ in $S(\mathfrak{g}) \otimes_{\mathbb{C}} \mathfrak{g}$ such that $\varphi(x)$ belongs to $\mathfrak{g}(x)$ for all x in \mathfrak{g} , is a free module of basis $\varepsilon_1, \dots, \varepsilon_{\mathrm{rk}\mathfrak{g}}$.

NILPOTENT BICONE

Definition 1. The characteristic submodule $B_{\mathfrak{g}}$ of \mathfrak{g} is the $S(\mathfrak{g}) \otimes_{\mathbb{C}} S(\mathfrak{g})$ -submodule of elements φ in $S(\mathfrak{g}) \otimes_{\mathbb{C}} S(\mathfrak{g}) \otimes_{\mathbb{C}} \mathfrak{g}$ such that $\varphi(x, y)$ belongs to $\mathfrak{A}'(x, y)$, for any (x, y) in a nonempty open subset of $\mathfrak{g} \times \mathfrak{g}$.

The following result can be viewed as a generalization of the previous result of Dixmier.

Theorem 11. *The submodule $B_{\mathfrak{g}}$ of $S(\mathfrak{g}) \otimes_{\mathbb{C}} S(\mathfrak{g}) \otimes_{\mathbb{C}} \mathfrak{g}$ is a free $S(\mathfrak{g}) \otimes_{\mathbb{C}} S(\mathfrak{g})$ -module of rank $b_{\mathfrak{g}}$. Moreover, the family $\{\varepsilon_{i,1,d_i-1}, \dots, \varepsilon_{i,d_i,0} \mid i = 1, \dots, \text{rk } \mathfrak{g}\}$ is a basis of $B_{\mathfrak{g}}$.*

Proof. By Lemma 8(ii), for $i = 1, \dots, \text{rk } \mathfrak{g}$ and $m = 1, \dots, d_i$, ε_{i,m,d_i-m} belongs to $B_{\mathfrak{g}}$. Moreover, by Lemma 8(i), these elements are linearly independent over $S(\mathfrak{g}) \otimes_{\mathbb{C}} S(\mathfrak{g})$. It remains to prove that they generate $B_{\mathfrak{g}}$ as an $S(\mathfrak{g}) \otimes_{\mathbb{C}} S(\mathfrak{g})$ -module. Let φ be in $B_{\mathfrak{g}}$. By Lemma 8(i) and (ii), $\varphi(x, y)$ belongs to $\mathfrak{A}'(x, y)$ for any (x, y) in $\Omega_{\mathfrak{g}}$. So there exist regular functions ψ_{i,m,d_i-m} on $\Omega_{\mathfrak{g}}$ for $i = 1, \dots, \text{rk } \mathfrak{g}$ and $m = 1, \dots, d_i$, such that

$$\varphi(x, y) = \sum_{\substack{i=1, \dots, \text{rk } \mathfrak{g} \\ m=1, \dots, d_i}} \psi_{i,m,d_i-m}(x, y) \varepsilon_{i,m,d_i-m}(x, y),$$

for any (x, y) in $\Omega_{\mathfrak{g}}$. By Corollary 10, $\Omega_{\mathfrak{g}}$ is a big open subset of $\mathfrak{g} \times \mathfrak{g}$. Hence the regular functions ψ_{i,m,d_i-m} have regular extensions to $\mathfrak{g} \times \mathfrak{g}$ since $\mathfrak{g} \times \mathfrak{g}$ is normal. As a result, the family $\{\varepsilon_{i,1,d_i-1}, \dots, \varepsilon_{i,d_i,0} \mid i = 1, \dots, \text{rk } \mathfrak{g}\}$ is a basis of $B_{\mathfrak{g}}$. \square

1.4. Smooth points of $\mathcal{N}_{\mathfrak{g}}$

In this subsection we establish a link between the open subset $\Omega_{\mathfrak{g}}$ and the nilpotent bicone. For $i = 1, \dots, \text{rk } \mathfrak{g}$, we denote by σ_i the map

$$\mathfrak{g} \times \mathfrak{g} \longrightarrow \mathbb{C}^{d_i+1}, \quad (x, y) \mapsto (p_{i,0,d_i}(x, y), \dots, p_{i,d_i,0}(x, y)).$$

Lemma 12. *For (x, y) in $\mathfrak{g} \times \mathfrak{g}$ and $i = 1, \dots, \text{rk } \mathfrak{g}$, the differential of σ_i at (x, y) is the linear map*

$$(v, w) \mapsto (\langle \varepsilon_{i,1,d_i-1}(x, y), w \rangle, \langle \varepsilon_{i,1,d_i-1}(x, y), v \rangle + \langle \varepsilon_{i,2,d_i-2}(x, y), w \rangle, \dots, \langle \varepsilon_{i,d_i-1,1}(x, y), v \rangle + \langle \varepsilon_{i,d_i,0}(x, y), w \rangle, \langle \varepsilon_i(x), v \rangle).$$

Proof. For (x, y) in $\mathfrak{g} \times \mathfrak{g}$, we denote by $p'_{i,m,d_i-m}(x, y)$ the differential of p_{i,m,d_i-m} at (x, y) . From Lemma 5 and the equality

$$p_i(tx + y) = \sum_{m=0}^{d_i} t^m p_{i,m,d_i-m}(x, y)$$

for (x, y) in $\mathfrak{g} \times \mathfrak{g}$, we deduce the equality

$$\sum_{m=1}^{d_i} t^m \langle \varepsilon_{i,m,d_i-m}(x, y), v + t^{-1}w \rangle = \sum_{m=0}^{d_i} t^m p'_{i,m,d_i-m}(x, y)(v, w),$$

for x, y, v, w in \mathfrak{g} and t in \mathbb{C}^* , since $p_{i,0,d_i}$ is the map $(x, y) \mapsto p_i(y)$. Hence we get, for $m = 0, \dots, d_i - 1$,

$$p'_{i,m,d_i-m}(x, y)(v, w) = \langle \varepsilon_{i,m,d_i-m}(x, y), v \rangle + \langle \varepsilon_{i,m+1,d_i-m-1}(x, y), w \rangle.$$

In addition, $p'_{i,d_i,0}(x, y)$ is the linear functional $(v, w) \mapsto \langle \varepsilon_i(x), v \rangle$. \square

Let σ be the map

$$\mathfrak{g} \times \mathfrak{g} \longrightarrow \mathbb{C}^{b_{\mathfrak{g}} + \text{rk } \mathfrak{g}}, \quad (x, y) \mapsto (\sigma_1(x, y), \dots, \sigma_{\text{rk } \mathfrak{g}}(x, y)).$$

Proposition 13. *Let (x, y) be in $\mathfrak{g} \times \mathfrak{g}$. Then σ is smooth at (x, y) if and only if (x, y) belongs to $\Omega_{\mathfrak{g}}$.*

Proof. We denote by $\sigma'(x, y)$ the differential of σ at (x, y) and we denote by $\ker \sigma'(x, y)$ its kernel in $\mathfrak{g} \times \mathfrak{g}$. Let us suppose that (x, y) belongs to $\Omega_{\mathfrak{g}}$. For v in \mathfrak{g} , we denote by X_v the subset of elements w in \mathfrak{g} such that (v, w) is in $\ker \sigma'(x, y)$. By Lemmas 12 and 4(ii), v belongs to the orthogonal complement of $\mathfrak{g}(x)$ in \mathfrak{g} . In addition, X_v is an affine subspace whose tangent space is equal to the orthogonal complement of $\mathfrak{V}(x, y)$ in \mathfrak{g} . By Lemma 8(i), $\mathfrak{V}(x, y)$ has dimension $b_{\mathfrak{g}}$ since (x, y) belongs to $\Omega_{\mathfrak{g}}$. Consequently, $\ker \sigma'(x, y)$ has dimension at most $b_{\mathfrak{g}} - \text{rk } \mathfrak{g} + 2(b_{\mathfrak{g}} - \text{rk } \mathfrak{g}) = 3(b_{\mathfrak{g}} - \text{rk } \mathfrak{g})$, since x is regular. Hence the image of $\sigma'(x, y)$ has dimension at least $b_{\mathfrak{g}} + \text{rk } \mathfrak{g}$. So $\sigma'(x, y)$ is surjective and σ is smooth at (x, y) .

Conversely, suppose that σ is smooth at (x, y) . For (a, b) in \mathbb{C}^2 , we denote by $\pi_{a,b}$ the linear map

$$(z_{i,0,d_i}, \dots, z_{i,d_i,0}, i = 1, \dots, \text{rk } \mathfrak{g}) \mapsto \left(\sum_{m+n=d_i} a^m b^n z_{i,m,n}, i = 1, \dots, \text{rk } \mathfrak{g} \right),$$

from $\mathbb{C}^{b_{\mathfrak{g}} + \text{rk } \mathfrak{g}}$ to $\mathbb{C}^{\text{rk } \mathfrak{g}}$. The linear map $\pi_{a,b}$ is surjective as soon as $(a, b) \neq (0, 0)$. Since σ is smooth at (x, y) , we deduce that the compound map $\sigma_{a,b} := \pi_{a,b} \circ \sigma$ is smooth at (x, y) for any (a, b) in $\mathbb{C}^2 \setminus \{(0, 0)\}$. As

$$\sum_{m+n=d_i} a^m b^n p_{i,m,n}(x', y') = p_i(ax' + by'),$$

for $i = 1, \dots, \text{rk } \mathfrak{g}$ and (x', y') in $\mathfrak{g} \times \mathfrak{g}$, $\sigma_{a,b}$ maps (x', y') to $(p_1(ax' + by'), \dots, p_{\text{rk } \mathfrak{g}}(ax' + by'))$. Moreover, it is the compound map of the two following maps:

$$\begin{aligned} \mathfrak{g} \times \mathfrak{g} &\longrightarrow \mathfrak{g}, & (x', y') &\mapsto (ax' + by'), \\ \mathfrak{g} &\longrightarrow \mathbb{C}^{\text{rk } \mathfrak{g}}, & z &\mapsto (p_1(z), \dots, p_{\text{rk } \mathfrak{g}}(z)). \end{aligned}$$

Therefore, the second map is smooth at $(ax + by)$ for any (a, b) in $\mathbb{C}^2 \setminus \{(0, 0)\}$. So, by Lemma 4(ii), (x, y) belongs to $\Omega_{\mathfrak{g}}$. \square

Remark 3. Recall that $\mathcal{N}_{\mathfrak{g}}$ is the subscheme of $\mathfrak{g} \times \mathfrak{g}$ defined by the ideal generated by the polynomials $p_{i,m,n}$. Therefore, by Proposition 13, the intersection of $\mathcal{N}_{\mathfrak{g}}$ and $\Omega_{\mathfrak{g}}$ is nothing but the set of smooth points of $\mathcal{N}_{\mathfrak{g}}$. The elements of $\Omega_{\mathfrak{g}}$ provided by Lemma 6(i), do not belong to $\mathcal{N}_{\mathfrak{g}}$. Consider the elements described in Lemma 6(ii). If \mathfrak{g} is equal to \mathfrak{sl}_3 , the subset $\mathfrak{g}'_{-2} \times \mathfrak{g}'_2$ has a nonempty intersection with $\mathcal{N}_{\mathfrak{g}}$. Indeed, if we set

$$e := \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}, \quad h := \begin{bmatrix} 2 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -2 \end{bmatrix}, \quad x := \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & -1 & 0 \end{bmatrix},$$

NILPOTENT BICONE

then $(x, e) \in \mathcal{N}_{\mathfrak{g}} \cap (\mathfrak{g}'_{-2} \times \mathfrak{g}'_2)$. But in general this intersection can be empty. For instance, it is an empty set if $\mathfrak{g} = \mathfrak{sl}_n$ for many $n > 3$. Whatever the case, we will see a posteriori that there exists at least one irreducible component of $\mathcal{N}_{\mathfrak{g}}$ which has an empty intersection with $\Omega_{\mathfrak{g}}$ (see Theorem 45). So we cannot hope to apply the criterion of Kostant [Kos2]. This observation was originally the reason why we introduced another subscheme (see the following section).

2. Principal cone and principal bicone

In this section we suppose that \mathfrak{g} is simple. Then we can suppose that p_1 is the Casimir element of $S(\mathfrak{g})^{\mathfrak{g}}$, that is, $p_1(x) = \langle x, x \rangle$ for any x in \mathfrak{g} . Recall that (e, h, f) is a principal \mathfrak{sl}_2 -triple of \mathfrak{g} and use the notations of the Introduction, Subsection 0.3.

2.1. Principal cone

Since e is a regular nilpotent element of \mathfrak{g} , the nilpotent cone $\mathfrak{N}_{\mathfrak{g}}$ is the G -invariant closed cone generated by e . According to Kostant's results [Kos2], the nilpotent cone is a complete intersection of codimension $\text{rk } \mathfrak{g}$. Moreover, it is proved in [He] that it has rational singularities. In this subsection we intend to prove analogous properties for *the principal cone* introduced in

Definition 2. The *principal cone* $\mathfrak{X}_{\mathfrak{g}}$ of \mathfrak{g} is the G -invariant closed cone generated by h .

Recall that w_0 is the longest element of the Weyl group $W_G(\mathfrak{h})$. The simple following well known lemma turns out to be useful.

Lemma 14. *The element $w_0(h)$ is equal to $-h$. Moreover, there exists a representative g_0 of w_0 in $N_G(\mathfrak{h})$ such that $g_0(e)$ is equal to f .*

For $i = 2, \dots, \text{rk } \mathfrak{g}$, we define the element q_i of $S(\mathfrak{g})^{\mathfrak{g}}$ as follows:

$$q_i = \begin{cases} p_i, & \text{if } d_i \text{ is odd;} \\ p_1(h)^{d_i/2} p_i - p_i(h) p_1^{d_i/2}, & \text{otherwise.} \end{cases}$$

The polynomial q_i is homogeneous of degree d_i . As the eigenvalues of $\text{ad } h$ are integers, not all equal to zero, $p_1(h) \neq 0$. In addition, as

$$p_i(h) = p_i(w_0(h)) = p_i(-h) = (-1)^{d_i} p_i(h),$$

$p_i(h) = 0$ as soon as d_i is odd. This forces $q_i(h) = 0$ for any $i = 2, \dots, \text{rk } \mathfrak{g}$. So $\mathfrak{X}_{\mathfrak{g}}$ is contained in the nullvariety of the functions $q_2, \dots, q_{\text{rk } \mathfrak{g}}$.

Lemma 15.

- (i) *The nullvariety of p_1 in $\mathfrak{X}_{\mathfrak{g}}$ is equal to $\mathfrak{N}_{\mathfrak{g}}$.*
- (ii) *The principal cone $\mathfrak{X}_{\mathfrak{g}}$ is the nullvariety of the functions $q_2, \dots, q_{\text{rk } \mathfrak{g}}$.*
- (iii) *The subset $(\mathfrak{X}_{\mathfrak{g}} \setminus \mathfrak{g}_{\text{reg}})$ is equal to $(\mathfrak{N}_{\mathfrak{g}} \setminus G.e)$ and the codimension of $(\mathfrak{X}_{\mathfrak{g}} \setminus \mathfrak{g}_{\text{reg}})$ in $\mathfrak{X}_{\mathfrak{g}}$ is equal to 3.*

Proof. (i) Prove first that $\mathfrak{N}_{\mathfrak{g}}$ is contained in $\mathfrak{X}_{\mathfrak{g}}$. Since $\mathfrak{X}_{\mathfrak{g}}$ is a G -invariant closed cone, the relation $\exp(-t \operatorname{ad} e)(h) = h + 2te$, for any $t \in \mathbb{C}$, implies $e \in \mathfrak{X}_{\mathfrak{g}}$. So $\mathfrak{N}_{\mathfrak{g}}$ is contained in $\mathfrak{X}_{\mathfrak{g}}$ as closure of $G.e$. Let X be the nullvariety of the functions $q_2, \dots, q_{\operatorname{rk} \mathfrak{g}}$. As $p_1(h) \neq 0$, the nullvariety of p_1 in X is the nullvariety of $p_1, \dots, p_{\operatorname{rk} \mathfrak{g}}$. So $\mathfrak{N}_{\mathfrak{g}}$ is the nullvariety of p_1 in X . Moreover, $\mathfrak{N}_{\mathfrak{g}}$ is also the nullvariety of p_1 in $\mathfrak{X}_{\mathfrak{g}}$ since $\mathfrak{X}_{\mathfrak{g}}$ is a subset of X which contains $\mathfrak{N}_{\mathfrak{g}}$.

(ii) Let X be as in (i). We need to prove that X is contained in $\mathfrak{X}_{\mathfrak{g}}$. Let x be an element of X which is not nilpotent. By (i), $p_1(x) \neq 0$. So there exists $t \in \mathbb{C}^*$ such that $p_1(tx) = p_1(h)$. Then $p_i(tx) = p_i(h)$, for $i = 1, \dots, \operatorname{rk} \mathfrak{g}$, since x is in X . As h is regular and semisimple, tx and h are G -conjugate. Hence x belongs to $\mathfrak{X}_{\mathfrak{g}}$.

(iii) By (i) and (ii), the subset of elements of $\mathfrak{X}_{\mathfrak{g}}$ which are not regular in \mathfrak{g} is equal to $(\mathfrak{N}_{\mathfrak{g}} \setminus G.e)$. As this subset has codimension 2 in $\mathfrak{N}_{\mathfrak{g}}$, it has codimension 3 in $\mathfrak{X}_{\mathfrak{g}}$ by (i). \square

Recall that \mathfrak{b}_- is the Borel subalgebra containing \mathfrak{h} and “opposite” to \mathfrak{b} , with nilpotent radical \mathfrak{u}_- . Let \mathfrak{b}_0 be the subspace of \mathfrak{b} generated by h and \mathfrak{u} and let $\mathfrak{b}_{0,-}$ be the subspace of \mathfrak{b}_- generated by h and \mathfrak{u}_- . The following results are partially proved in [Ri2, Prop. 10.3] as pointed out in Remark 4 below.

Corollary 16. *The principal cone of \mathfrak{g} is normal and it is a complete intersection of codimension $\operatorname{rk} \mathfrak{g} - 1$ in \mathfrak{g} . The regular elements of \mathfrak{g} which belong to $\mathfrak{X}_{\mathfrak{g}}$ are smooth points of $\mathfrak{X}_{\mathfrak{g}}$. Finally, $\mathfrak{X}_{\mathfrak{g}}$ has rational singularities.*

Proof. By Lemma 15(i) and (ii), $\mathfrak{X}_{\mathfrak{g}}$ is a complete intersection of codimension $\operatorname{rk} \mathfrak{g} - 1$. By Lemma 4(ii), the differentials at x of $p_1, \dots, p_{\operatorname{rk} \mathfrak{g}}$ are linearly independent as soon as x is a regular element of \mathfrak{g} . Hence the same goes for the differentials at x of $q_2, \dots, q_{\operatorname{rk} \mathfrak{g}}$. Then, any x in the union of $G.e$ and $(\mathbb{C}^*)G.h$ is a smooth point of $\mathfrak{X}_{\mathfrak{g}}$. So by Lemma 15, (ii), $\mathfrak{X}_{\mathfrak{g}}$ is regular in codimension 1. By Serre’s normality criterion [Ma, Chap. 8, Theorem 23.8], $\mathfrak{X}_{\mathfrak{g}}$ is a normal subvariety.

The subalgebra \mathfrak{b}_0 is an ideal of \mathfrak{b} . The contracted product $G \times_{\mathbf{B}} \mathfrak{b}_0$ is defined as the quotient of $G \times \mathfrak{b}_0$ under the right action of \mathbf{B} given by $(g, x).b = (gb, b^{-1}(x))$. The map $(g, x) \mapsto g(x)$ from $G \times \mathfrak{b}_0$ to \mathfrak{g} factorizes through the canonical map from $G \times \mathfrak{b}_0$ to $G \times_{\mathbf{B}} \mathfrak{b}_0$. Since $G.\mathfrak{b}_0$ is equal to $\mathfrak{X}_{\mathfrak{g}}$, we get a surjective morphism π from $G \times_{\mathbf{B}} \mathfrak{b}_0$ to $\mathfrak{X}_{\mathfrak{g}}$. Moreover, the morphism π is proper since G/\mathbf{B} is a projective variety. Let K be the field of rational functions on $G \times_{\mathbf{B}} \mathfrak{b}_0$ and let K_0 be the field of rational functions on $\mathfrak{X}_{\mathfrak{g}}$. Let g_0 be as in Lemma 14. By Lemma 14, $g_0(h)$ is equal to $-h$. In particular, the fibre of π at h has at least two elements. Let (g, x) be an element of $G \times \mathfrak{b}_0$ such that $g(x)$ is equal to h . In particular, x is a regular semisimple element of \mathfrak{g} which belongs to \mathfrak{b}_0 . Hence, there exists b in \mathbf{B} such that $b(x)$ belongs to the line generated by h . As $\operatorname{ad} h$ and $\operatorname{ad} b(x)$ have the same eigenvalues, $b(x)$ is equal to h or $-h$, since the eigenvalues of $\operatorname{ad} h$ are integers, not all equal to zero. If $b(x) = h$, then (g, x) and $(\mathbf{1}_{\mathfrak{g}}, h)$ are equal in $G \times_{\mathbf{B}} \mathfrak{b}_0$. Otherwise, (g, x) and $(g_0, -h)$ are equal in $G \times_{\mathbf{B}} \mathfrak{b}_0$. Hence, the fibre of π at h has two elements. So the same applies for the fibre of π at any element of the G -invariant cone generated by h since π is G -equivariant. Hence, K is an algebraic extension of degree 2 of K_0 . In particular, it is a Galois extension. Let A be the integral closure of $\mathbb{C}[\mathfrak{X}_{\mathfrak{g}}]$ in K and let $\mathfrak{X}'_{\mathfrak{g}}$ be an affine algebraic variety such that $\mathbb{C}[\mathfrak{X}'_{\mathfrak{g}}]$ is equal to A . Then A is stable under the Galois group of the

NILPOTENT BICONE

extension K of K_0 and $\mathbb{C}[\mathfrak{X}_{\mathfrak{g}}]$ is the subalgebra of invariant elements of this action since $\mathfrak{X}_{\mathfrak{g}}$ is normal. Hence, by [El2, Lemma 1], it is enough to prove that $\mathfrak{X}'_{\mathfrak{g}}$ has rational singularities.

The variety $G \times_{\mathbf{B}} \mathfrak{b}_0$ is smooth as a vector bundle over G/\mathbf{B} . Hence the morphism π factors through the canonical morphism from $\mathfrak{X}'_{\mathfrak{g}}$ to $\mathfrak{X}_{\mathfrak{g}}$. Let π' be the morphism from $G \times_{\mathbf{B}} \mathfrak{b}_0$ to $\mathfrak{X}'_{\mathfrak{g}}$ such that π is the compound map of π' with the canonical morphism from $\mathfrak{X}'_{\mathfrak{g}}$ to $\mathfrak{X}_{\mathfrak{g}}$. Then $(G \times_{\mathbf{B}} \mathfrak{b}_0, \pi')$ is a desingularization of $\mathfrak{X}'_{\mathfrak{g}}$ since π' is proper and birational. By the corollary of [He, Theorem B], for $i \in \mathbb{N}^*$, the i -th cohomology group $H^i(G \times_{\mathbf{B}} \mathfrak{b}_0, \mathcal{O}_{G \times_{\mathbf{B}} \mathfrak{b}_0})$ is equal to zero. Hence by [Ha, Chap. III, Prop. 8.5], for $i \in \mathbb{N}^*$, $R^i \pi'_*(\mathcal{O}_{G \times_{\mathbf{B}} \mathfrak{b}_0}) = 0$ and $\mathfrak{X}'_{\mathfrak{g}}$ has rational singularities since $\mathfrak{X}'_{\mathfrak{g}}$ is normal. \square

Remark 4. The G -invariant closed cone generated by a semisimple element of \mathfrak{g} is not a normal variety in general [Ri2, Prop. 10.1]. Nevertheless, whenever x is the semisimple element of an \mathfrak{sl}_2 -triple of \mathfrak{g} , the closed cone generated by the regular orbit whose closure contains x is normal and Cohen–Macaulay [Ri2, Prop. 10.3].

2.2. Principal bicone

In this subsection we study various properties of the *principal bicone*.

Definition 3. The *principal bicone* of \mathfrak{g} is the subset

$$\begin{aligned} \mathcal{X}_{\mathfrak{g}} &:= \{(x, y) \in \mathfrak{g} \times \mathfrak{g} \mid P_{x,y} \subset \mathfrak{X}_{\mathfrak{g}}\} \\ &= \{(x, y) \in \mathfrak{g} \times \mathfrak{g} \mid ax + by \in \mathfrak{X}_{\mathfrak{g}}, \forall (a, b) \in \mathbb{C}^2\}. \end{aligned}$$

As $\mathcal{N}_{\mathfrak{g}}$, the subset $\mathcal{X}_{\mathfrak{g}}$ is a closed bicone of $\mathfrak{g} \times \mathfrak{g}$, and it is invariant under the actions of G and $\mathrm{GL}_2(\mathbb{C})$. For $i = 2, \dots, \mathrm{rk} \mathfrak{g}$, we define the elements $q_{i,m,n}$ of $(\mathbb{S}(\mathfrak{g}) \otimes_{\mathbb{C}} \mathbb{S}(\mathfrak{g}))^{\mathfrak{g}}$ by the following relation:

$$q_i(ax + by) = \sum_{m+n=d_i} a^m b^n q_{i,m,n}(x, y), \quad (2)$$

for (a, b) in \mathbb{C}^2 and (x, y) in $\mathfrak{g} \times \mathfrak{g}$. For w in $W_G(\mathfrak{h})$, we denote by $\mathfrak{b}_{0,w}$ the subspace generated by $w(\mathfrak{h})$ and \mathfrak{u} . It is worth recalling the following well known result for the understanding of the principal bicone.

Lemma 17. *If x is in \mathfrak{h} , then the intersection of $G.x$ and \mathfrak{h} is the orbit of x under $W_G(\mathfrak{h})$.*

Our purpose is to prove that $\mathcal{X}_{\mathfrak{g}}$ is a complete intersection of dimension $3(\mathfrak{b}_{\mathfrak{g}} - \mathrm{rk} \mathfrak{g} + 1)$, which we will do in Section 4.

Lemma 18.

- (i) *The subset $\mathcal{X}_{\mathfrak{g}}$ is the nullvariety of the polynomial functions $q_{i,m,n}$ for $i = 2, \dots, \mathrm{rk} \mathfrak{g}$ and $m + n = d_i$. In particular, any irreducible component of $\mathcal{X}_{\mathfrak{g}}$ has dimension at least $3(\mathfrak{b}_{\mathfrak{g}} - \mathrm{rk} \mathfrak{g} + 1)$.*
- (ii) *The subset $\mathfrak{s} \times \mathfrak{s}$ is contained in $\mathcal{X}_{\mathfrak{g}}$.*
- (iii) *For x in \mathfrak{b} , $(w(\mathfrak{h}), x)$ belongs to $\mathcal{X}_{\mathfrak{g}}$ if and only if x is in $\mathfrak{b}_{0,w}$.*

Proof. (i) As $\mathfrak{X}_{\mathfrak{g}}$ is the nullvariety of the q_i by Lemma 15(ii), $\mathcal{X}_{\mathfrak{g}}$ is the nullvariety of the polynomial functions $q_{i,m,n}$ for $i = 2, \dots, \text{rk } \mathfrak{g}$ and $m + n = d_i$. As $d_1 = 2$, $\mathcal{X}_{\mathfrak{g}}$ is the nullvariety in $\mathfrak{g} \times \mathfrak{g}$ of $\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g} - 3$ regular functions, whence the second statement.

(ii) Let T be the intersection of $\mathcal{X}_{\mathfrak{g}}$ and $\mathfrak{s} \times \mathfrak{s}$. Then T is a closed, subset of $\mathfrak{s} \times \mathfrak{s}$, invariant under $\text{GL}_2(\mathbb{C})$ and \mathbf{S} . For any t in \mathbb{C} , $h + te$ belongs to $G.h$. So T contains (h, e) . Hence, by Lemma 9, $\dim T \geq 5$ and $T = T_5$ if and only if $\dim T = 5$. As th and $e + t^2f$ are in the same G -orbit for any $t \in \mathbb{C}^*$, (e, f) is in $\mathcal{X}_{\mathfrak{g}}$. So T is equal to $\mathfrak{s} \times \mathfrak{s}$ since (e, f) is not in T_5 .

(iii) For any t in \mathbb{C}^* and any x in \mathfrak{u} , $tw(h) + x$ belongs to the G -orbit of $tw(h)$. Hence $(w(h), tw(h) + x)$ belongs to $\mathcal{X}_{\mathfrak{g}}$. So $\{w(h)\} \times \mathfrak{b}_{0,w}$ is contained in $\mathcal{X}_{\mathfrak{g}}$. Let (x_1, x_2) be in $\mathfrak{h} \times \mathfrak{u}$. We suppose that $(w(h), x_1 + x_2)$ is in $\mathcal{X}_{\mathfrak{g}}$. Then for any t in \mathbb{C} , $w(h) + tx_1 + tx_2$ is in $\mathfrak{X}_{\mathfrak{g}}$. In particular, $x_1 + x_2$ is tangent at $w(h)$ to the cone $\mathfrak{X}_{\mathfrak{g}}$. The tangent space at $w(h)$ of the cone $\mathfrak{X}_{\mathfrak{g}}$ is generated by $w(h)$ and $[w(h), \mathfrak{g}]$. So x_1 and $w(h)$ are collinear. \square

2.3. Smooth points of $\mathcal{X}_{\mathcal{P}}$

As the nilpotent cone $\mathfrak{N}_{\mathfrak{g}}$ is contained in the principal cone $\mathfrak{X}_{\mathfrak{g}}$, the nilpotent bicone $\mathcal{N}_{\mathfrak{g}}$ is contained in the principal bicone $\mathcal{X}_{\mathfrak{g}}$. We now introduce two other varieties “squeezed in between” $\mathcal{N}_{\mathfrak{g}}$ and $\mathcal{X}_{\mathfrak{g}}$. Set

$$\begin{aligned} \mathcal{Y}_{\mathfrak{g}} &:= \{ (x, y) \in \mathcal{X}_{\mathfrak{g}} \mid \langle x, y \rangle = 0 \}, \\ \mathcal{Z}_{\mathfrak{g}} &:= \{ (x, y) \in \mathcal{Y}_{\mathfrak{g}} \mid y \in \mathfrak{N}_{\mathfrak{g}} \}. \end{aligned}$$

Lemma 19.

- (i) *The nilpotent bicone is the nullvariety in $\mathcal{X}_{\mathfrak{g}}$ of $p_{1,2,0}, p_{1,1,1}, p_{1,0,2}$.*
- (ii) *The subset $\mathcal{Y}_{\mathfrak{g}}$ is the nullvariety in $\mathcal{X}_{\mathfrak{g}}$ of $p_{1,1,1}$. Moreover, $\mathcal{Y}_{\mathfrak{g}}$ is a G -invariant closed bicone and any irreducible component of $\mathcal{Y}_{\mathfrak{g}}$ has dimension at least $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g}) + 2$.*
- (iii) *The subset $\mathcal{Z}_{\mathfrak{g}}$ is the nullvariety in $\mathcal{Y}_{\mathfrak{g}}$ of $p_{1,0,2}$. Moreover, $\mathcal{Z}_{\mathfrak{g}}$ is a G -invariant closed bicone and any irreducible component of $\mathcal{Z}_{\mathfrak{g}}$ has dimension at least $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g}) + 1$.*

Proof. (i) As $\mathfrak{N}_{\mathfrak{g}}$ is the nullvariety of p_1 in $\mathfrak{X}_{\mathfrak{g}}$, by Lemma 15(i), $\mathcal{N}_{\mathfrak{g}}$ is the nullvariety in $\mathcal{X}_{\mathfrak{g}}$ of the 2-order polarizations of p_1 , that is $p_{1,2,0}, p_{1,1,1}, p_{1,0,2}$.

(ii) As $p_{1,1,1}$ is the Killing form, $\mathcal{Y}_{\mathfrak{g}}$ is the nullvariety of $p_{1,1,1}$ in $\mathcal{X}_{\mathfrak{g}}$. Moreover, $\mathcal{Y}_{\mathfrak{g}}$ is a G -invariant closed bicone since $p_{1,1,1}$ is G -invariant and bihomogeneous. By Lemma 18(i), any irreducible component of $\mathcal{Y}_{\mathfrak{g}}$ has dimension at least $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g}) + 2$.

(iii) By Lemma 15(i), $\mathfrak{N}_{\mathfrak{g}}$ is the nullvariety of p_1 in $\mathfrak{X}_{\mathfrak{g}}$. Hence $\mathcal{Z}_{\mathfrak{g}}$ is the nullvariety of $p_{1,0,2}$ in $\mathcal{Y}_{\mathfrak{g}}$. Moreover, $\mathcal{Z}_{\mathfrak{g}}$ is a G -invariant closed bicone since $p_{1,0,2}$ is G -invariant and bihomogeneous. By (ii), any irreducible component of $\mathcal{Z}_{\mathfrak{g}}$ has dimension at least $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g}) + 1$. \square

Remark 5. By Lemma 19, the subsets $\mathcal{Y}_{\mathfrak{g}}$ and $\mathcal{Z}_{\mathfrak{g}}$ inherit a natural structure of the scheme.

NILPOTENT BICONE

Let P be a subset of the set of polynomials

$$\{p_{1,2,0}, p_{1,1,1}, p_{1,0,2}, q_{i,m,d_i-m} \mid i = 2, \dots, \text{rk } \mathfrak{g}, m = 0, \dots, d_i\}.$$

We denote by $\mathbb{C}[P]$ the subalgebra of $S(\mathfrak{g}) \otimes_{\mathbb{C}} S(\mathfrak{g})$ generated by P , we denote by σ_P the morphism of affine varieties whose comorphism is the canonical injection from $\mathbb{C}[P]$ to $S(\mathfrak{g}) \otimes_{\mathbb{C}} S(\mathfrak{g})$ and we denote by \mathcal{X}_P the nullvariety of P in $\mathfrak{g} \times \mathfrak{g}$.

Lemma 20. *Let (x, y) be in $\Omega_{\mathfrak{g}}$.*

- (i) *The morphism σ_P is smooth at (x, y) .*
- (ii) *If (x, y) is in \mathcal{X}_P , then (x, y) is a smooth point of \mathcal{X}_P . Moreover, the unique irreducible component of \mathcal{X}_P which contains (x, y) has dimension $2 \dim \mathfrak{g} - |P|$.*

Proof. (i) Let ρ be the morphism whose comorphism is the canonical injection from $\mathbb{C}[P]$ to the subalgebra generated by the polynomials p_{i,m,d_i-m} where $i = 1, \dots, \text{rk } \mathfrak{g}$ and $m = 0, \dots, d_i$. Then σ_P is equal to $\rho \circ \sigma$, where σ is the morphism introduced in Subsection 1.4. As q_{i,m,d_i-m} is a linear combination of p_{i,m,d_i-m} and homogeneous elements in the subalgebra generated by $p_{1,2,0}, p_{1,1,1}, p_{1,0,2}$, for $i = 2, \dots, \text{rk } \mathfrak{g}$ and $m = 0, \dots, d_i$, the morphism ρ is smooth. Then, by Proposition 13, σ_P is a smooth morphism at (x, y) .

(ii) We suppose that (x, y) is in \mathcal{X}_P . By definition, \mathcal{X}_P is the fibre at 0 of the morphism σ_P . So by (i), (x, y) is a smooth point of \mathcal{X}_P and the codimension in $\mathfrak{g} \times \mathfrak{g}$ of the irreducible component of \mathcal{X}_P which contains (x, y) is equal to the dimension of $\mathbb{C}[P]$. By Proposition 13, the polynomials p_{i,m,d_i-m} are algebraically independent for $i = 1, \dots, \text{rk } \mathfrak{g}$ and $m = 0, \dots, d_i$. So the elements of P are algebraically independent since q_{i,m,d_i-m} is a linear combination of p_{i,m,d_i-m} and homogeneous elements in the subalgebra generated by $p_{1,2,0}, p_{1,1,1}, p_{1,0,2}$ for $i = 2, \dots, \text{rk } \mathfrak{g}$ and $m = 0, \dots, d_i$. Hence the irreducible component of \mathcal{X}_P containing (x, y) has dimension $2 \dim \mathfrak{g} - |P|$. \square

The principal bicone $\mathcal{X}_{\mathfrak{g}}$ has irreducible components of the expected dimension as the following proposition shows. Actually this proposition will be not used in the other sections.

Proposition 21. *Let w be in $W_G(\mathfrak{h})$. There is a unique irreducible component $\chi(w)$ of $\mathcal{X}_{\mathfrak{g}}$ containing $(w(h), e)$ and $\chi(w)$ satisfies the following properties:*

- (1) $\chi(w)$ has dimension $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g} + 1)$;
- (2) $\chi(w w_0)$ is equal to $\chi(w)$; and
- (3) $\chi(w)$ contains $(w_0 w(h), f), (e, w(h)), (f, w_0 w(h))$.

Moreover, $\chi(w)$ contains $\mathfrak{s} \times \mathfrak{s}$ if $w = 1$ or $w = w_0$.

Proof. Let P be the subset $\{q_{i,m,d_i-m} \mid i = 2, \dots, \text{rk } \mathfrak{g}, m = 0, \dots, d_i\}$. Then \mathcal{X}_P is equal to $\mathcal{X}_{\mathfrak{g}}$ by Lemma 18(i). By Lemma 18(iii), $\mathcal{X}_{\mathfrak{g}}$ contains $(w(h), e)$ and by Lemma 6(i), $(w(h), e)$ belongs to $\Omega_{\mathfrak{g}}$. Hence by Lemma 20(ii), $(w(h), e)$ is a smooth point of $\mathcal{X}_{\mathfrak{g}}$. So there is a unique irreducible component of $\mathcal{X}_{\mathfrak{g}}$ which contains $(w(h), e)$. Moreover, this component has dimension $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g} + 1)$. Let us denote it by $\chi(w)$. Then $\chi(w)$ satisfies condition (1). As $w w_0(h)$ is equal to

$-w(h)$, by Lemma 14, $\chi(w)$ contains $(w_0w(h), e)$ since $\chi(w)$ is a bicone. Moreover, by Lemma 14, $\chi(w)$ contains $(w_0w(h), f)$ since $\chi(w)$ is G -invariant and $g_0 \cdot (w(h), e)$ is equal to $(w_0w(h), f)$. As $\chi(w)$ is also $\mathrm{GL}_2(\mathbb{C})$ -invariant, it is invariant under the involution $(x, y) \mapsto (y, x)$. So $\chi(w)$ contains $(e, w(h))$ and $(f, w_0w(h))$.

By Lemma 18(ii), $\mathfrak{s} \times \mathfrak{s}$ is contained in $\mathcal{X}_{\mathfrak{g}}$. Hence, there exists an irreducible component of $\mathcal{X}_{\mathfrak{g}}$ which contains $\mathfrak{s} \times \mathfrak{s}$. But $\mathfrak{s} \times \mathfrak{s}$ contains (h, e) , which is a smooth point of $\mathcal{X}_{\mathfrak{g}}$. So $\chi(w)$ contains $\mathfrak{s} \times \mathfrak{s}$ if $w(h)$ is collinear to h , that is to say, $w = 1$ or $w = w_0$. \square

Remark 6. When \mathfrak{g} is simple of type $B_\ell, C_\ell, D_{2\ell}, E_7, E_8, F_4$ or G_2 , w_0 is equal to -1 . So, in these cases, for any w in $W_G(\mathfrak{h})$, $\chi(w)$ contains $(w(h), f)$ and $(f, w(h))$ since $\chi(w)$ is a bicone. A quick computation shows that $\mathcal{X}_{\mathfrak{g}}$ has precisely $|W_G(\mathfrak{h})|/2$ irreducible components when \mathfrak{g} is equal to $\mathfrak{sl}_n(\mathbb{C})$ for $n = 2, 3$.

2.4. Proof of Corollary 23

The goal of this subsection is to prove Corollary 23. This corollary will be crucial for the study of the dimension of $\mathcal{X}_{\mathfrak{g}}$ (see Section 4).

Lemma 22. *Let \mathcal{Z} be an irreducible closed subset of $\mathcal{Z}_{\mathfrak{g}}$ satisfying the following two conditions:*

- (1) $\varpi_1(\mathcal{Z})$ contains a nonzero semisimple element of \mathfrak{g} ; and
- (2) $\varpi_2(\mathcal{Z})$ contains a regular nilpotent element of \mathfrak{g} .

Then \mathcal{Z} has a nonempty intersection with $\Omega_{\mathfrak{g}}$.

Proof. By condition (1), the subset \mathcal{Z}_1 of elements (x, y) in \mathcal{Z} , such that x is a nonzero semisimple element of \mathfrak{g} , is a nonempty open subset of \mathcal{Z} since the subset of semisimple elements of \mathfrak{g} which belong to $\mathfrak{X}_{\mathfrak{g}}$ is an open dense subset of $\mathfrak{X}_{\mathfrak{g}}$. On the other hand, by condition (2), the subset \mathcal{Z}_2 of elements (x, y) in \mathcal{Z} , such that y is a regular nilpotent element of \mathfrak{g} , is a nonempty open subset of \mathcal{Z} since the subset of regular nilpotent elements is an open dense subset of $\mathfrak{N}_{\mathfrak{g}}$. Then the intersection of \mathcal{Z}_1 and \mathcal{Z}_2 is a nonempty subset since \mathcal{Z} is irreducible. Let (x, y) be in the intersection \mathcal{Z}_1 and \mathcal{Z}_2 . Suppose that (x, y) doesn't belong to $\Omega_{\mathfrak{g}}$. We expect a contradiction. Then there exists $t \in \mathbb{C}$ such that $tx + y$ is not a regular element of \mathfrak{g} . Indeed, x is regular as a nonzero semisimple element of $\mathfrak{X}_{\mathfrak{g}}$. As (x, y) belongs to the principal bicone, $tx + y$ is a nonregular element of $\mathfrak{X}_{\mathfrak{g}}$, whence

$$\langle tx + y, tx + y \rangle = 0,$$

by Lemma 15(iii). But the left-hand side of this equality is equal to $t^2 \langle x, x \rangle$ since (x, y) belongs to $\mathcal{Z}_{\mathfrak{g}}$. Hence $t = 0$ since for any nonzero semisimple element z in $\mathfrak{X}_{\mathfrak{g}}$, $\langle z, z \rangle \neq 0$. This contradicts the fact that y is regular. So \mathcal{Z} has a nonempty intersection with $\Omega_{\mathfrak{g}}$. \square

Corollary 23. *Let \mathcal{X} be an irreducible component of $\mathcal{X}_{\mathfrak{g}}$. We denote by \mathcal{X}' the intersection of \mathcal{X} and $\mathcal{Z}_{\mathfrak{g}}$. Let us suppose that \mathcal{X}' has an irreducible component \mathcal{Z} satisfying the following two conditions:*

- (1) $\varpi_1(\mathcal{Z})$ is equal to $\mathfrak{X}_{\mathfrak{g}}$; and
- (2) $\varpi_2(\mathcal{Z})$ is equal to $\mathfrak{N}_{\mathfrak{g}}$.

Then \mathcal{X}' and \mathcal{X} have a nonempty intersection with $\Omega_{\mathfrak{g}}$. In particular, \mathcal{X} has dimension $3(\mathrm{b}_{\mathfrak{g}} - \mathrm{rk} \mathfrak{g} + 1)$.

NILPOTENT BICONE

Proof. By Lemma 22, \mathcal{Z} , \mathcal{X}' and \mathcal{X} have a nonempty intersection with $\Omega_{\mathfrak{g}}$. Then, by Lemma 20(ii), \mathcal{X} has dimension $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g} + 1)$. \square

3. Jet schemes and motivic integration

We plan to study in Section 4 the dimensions of the nilpotent bicone and the principal bicone via motivic integration arguments. In view of this work, we start this section by some basics on motivic integration.

3.1. Jet schemes

Let us first review the definition and properties of jet schemes. Let X be a complex algebraic variety. For $m \in \mathbb{N}$, the m -order jet scheme $J_m(X)$ of X is the scheme whose closed points over $x \in X$ are morphisms

$$\mathcal{O}_{X,x} \longrightarrow \mathbb{C}[t]/t^{m+1}.$$

Thus, the \mathbb{C} -valued points of $J_m(X)$ are in natural bijection with the $\mathbb{C}[t]/t^{m+1}$ -valued points of X . In particular, there are canonical isomorphisms $J_0(X) \simeq X$ and $J_1(X) \simeq \text{TX}$, where TX is the total tangent space of X . The canonical morphisms

$$\pi_{l,m} : J_l(X) \longrightarrow J_m(X),$$

with $l \geq m$, obtained by truncation, induce a projective system $(J_m(X), \pi_{l,m})$. The space of the arcs

$$J_{\infty}(X) := \text{proj} \lim_m J_m(X)$$

is the projective limit of the system $(J_m(X), \pi_{l,m})$. Denote by $\pi_{\infty,m}$ the canonical morphism

$$\pi_{\infty,m} : J_{\infty}(X) \longrightarrow J_m(X),$$

for $m \in \mathbb{N}$, obtained by truncation, too. For ν in $J_{\infty}(X)$ and φ a regular function on X , we denote by $\text{ord}(\nu \circ \varphi)$ the order of the series $\nu \circ \varphi$. If \mathcal{I} is an ideal of \mathcal{O}_X , we denote by $\text{ord}(\mathcal{I}, \nu)$ the smallest integer $\text{ord}(\nu \circ \varphi)$, where φ runs through the ideal $\mathcal{I}_{\pi_{\infty,0}(\nu)}$ generated by \mathcal{I} in the local ring $\mathcal{O}_{X, \pi_{\infty,0}(\nu)}$. The function $F_{\mathcal{I}} : \nu \mapsto \text{ord}(\mathcal{I}, \nu)$ is semialgebraic by [DL2, Theorem 2.1]. As a matter of fact, for $m \in \mathbb{N}$, we can define an analogous function from $J_m(X)$ to $\{1, \dots, m\}$ that we denote by the same symbol. When \mathcal{I} is the ideal of definition of a closed subset Z of X , we rather denote by F_Z the function $F_{\mathcal{I}}$. In particular, if \mathcal{I} is the ideal of definition of a divisor D of X , we denote by F_D the function $F_{\mathcal{I}}$.

In [Mu], Mustařa proves the following result, first conjectured by David Eisenbud and Edward Frenkel.

Theorem 24 (Mustařa). *If X is a locally complete intersection variety, then $J_m(X)$ is irreducible of dimension $\dim X(m+1)$ for all $m \geq 1$ if and only if X has rational singularities.*

Remark 7. The nilpotent cone $\mathfrak{N}_{\mathfrak{g}}$ of the reductive Lie algebra \mathfrak{g} is a complete intersection and, by [He, Theorem A], it has rational singularities. Likewise, by Corollary 16, the principal cone $\mathfrak{X}_{\mathfrak{g}}$ of \mathfrak{g} is a complete intersection and has rational singularities. As a consequence, Theorem 24 can be applied to $\mathfrak{N}_{\mathfrak{g}}$ or $\mathfrak{X}_{\mathfrak{g}}$. In the Appendix of [Mu], Eisenbud and Frenkel apply Theorem 24 to $\mathfrak{N}_{\mathfrak{g}}$ to extend Kostant's results in the setting of jet schemes.

3.2. Motivic integration

Recall now some facts about the theory of motivic integration. The construction of motivic integrals for smooth spaces is due to Kontsevich [Kon] and was generalized by Denef and Loeser to singular spaces in [DL2] and [DL1]. Further in the paper, we will only need the Hodge realization of motivic integrals on the arcs of a smooth variety. We refer to [Ba], [DL2] and [Lo] for more explanations, definitions and proofs (see also [Cr] for an introduction).

Let \mathcal{V} be the category of complex algebraic varieties. Denote by $K_0(\mathcal{V})$ the Grothendieck ring of \mathcal{V} and denote by $[X]$ the class in $K_0(\mathcal{V})$ of an element X in \mathcal{V} . The map $X \mapsto [X]$ naturally extends to the category of constructible subsets of algebraic varieties. Let \mathbb{L} be the class of \mathbb{A}^1 in $K_0(\mathcal{V})$ and let \mathbf{M} be the localization $K_0(\mathcal{V})[\mathbb{L}^{-1}]$. For m in \mathbb{Z} , we denote by $F^m\mathbf{M}$ the subgroup of \mathbf{M} generated by the elements $[X]\mathbb{L}^{-r}$, where $r - \dim X \geq m$. Then we get a decreasing filtration of the ring \mathbf{M} and we denote by $\widehat{\mathbf{M}}$ its separated completion.

Let X be an algebraic variety of pure dimension d . A subset A in $J_\infty(X)$ is called a *cylinder* if A is a finite union of fibres of $\pi_{\infty,m} : J_\infty(X) \rightarrow J_m(X)$ for $m \in \mathbb{N}$. If $A \in J_\infty(X)$ is a cylinder, we say that A is *stable at level* $m \in \mathbb{N}$ if, for any $n \geq m$, the map $\pi_{\infty,n+1}(J_\infty(X)) \rightarrow \pi_{\infty,n}(J_\infty(X))$ is a piecewise trivial fibration over $\pi_{\infty,n}(A)$ with fibre \mathbb{A}^d . When X is smooth, any cylinder is stable and the later additional condition is superfluous.

Proposition 25 ([DL2, Definition-Proposition 3.2]). *There is a well defined sub-algebra \mathbf{B}_X of the Boolean algebra of subsets of $J_\infty(X)$ which contains the cylinders and a map μ_X from \mathbf{B}_X to $\widehat{\mathbf{M}}$ satisfying the following properties:*

- (1) *if $A \in \mathbf{B}_X$ is stable at level m , then $\mu_X(A) = [\pi_{\infty,m}(A)]\mathbb{L}^{-(m+1)d}$;*
- (2) *if Y is a closed subvariety in X of dimension strictly smaller than d , then for any A in \mathbf{B}_X , contained in $J_\infty(Y)$, $\mu_X(A) = 0$;*
- (3) *let A_1, A_2, \dots be a sequence of elements in \mathbf{B}_X whose union A is in \mathbf{B}_X , then the sequence $\mu_X(A_1), \mu_X(A_2), \dots$ converges to $\mu_X(A)$ in $\widehat{\mathbf{M}}$; and*
- (4) *if A and B are in \mathbf{B}_X , A is contained in B and $\mu_X(B)$ belongs to the closure of $F^m\mathbf{M}$ in $\widehat{\mathbf{M}}$, then $\mu_X(A)$ belongs to the closure of $F^m\mathbf{M}$ in $\widehat{\mathbf{M}}$.*

For A in \mathbf{B}_X and $\psi : A \rightarrow \mathbb{Z} \cup \{\infty\}$ a function such that $\psi^{-1}(s) \in \mathbf{B}_X$, for any $s \in \mathbb{Z} \cup \{\infty\}$ and $\mu_X(\psi^{-1}(\infty)) = 0$, we can set

$$\int_A \mathbb{L}^{-\psi} d\mu_X := \sum_{s \in \mathbb{Z}} \mu_X(\psi^{-1}(s))\mathbb{L}^{-s} \quad (3)$$

in $\widehat{\mathbf{M}}$, whenever the right-hand side converges in $\widehat{\mathbf{M}}$. In this case, we say that $\mathbb{L}^{-\psi}$ is *integrable on A* .

Recall that the dimension of a nonirreducible variety is the maximal dimension of its irreducible components. Let u and v be two indeterminate variables. For any smooth projective variety X , the *Hodge-Deligne polynomial* of X is the element of $\mathbb{Z}[u, v]$

$$h(X) := \sum_{k \in \mathbb{N}} \sum_{\substack{(p,q) \in \mathbb{N}^2 \\ p+q=k}} (-1)^k h_{p,q}(H_c^k(X; \mathbb{C})) u^p v^q,$$

NILPOTENT BICONE

where $h_{p,q}(H_c^k(X; \mathbb{C}))$ are the Hodge–Deligne numbers of X . The map h factors through the ring $K_0(\mathcal{V})$. So we have a morphism h from $K_0(\mathcal{V})$ to $\mathbb{Z}[u, v]$ such that $h(X)$ is equal to $h([X])$ for any smooth projective variety. In particular, $h(L)$ is equal to uv . What is important for us is that $h([X])$ is a polynomial whose highest degree term is $c(uv)^{\dim X}$, where c is the number of irreducible components of X of dimension $\dim X$. By continuity, the morphism h uniquely extends to a morphism from $\widehat{\mathbf{M}}$ to the ring $\mathbb{Z}[u, v][[u^{-1}, v^{-1}]]$. The compound map $h \circ \mu_X$, which is now well defined, is *the Hodge realization of the motivic measure*.

3.3. Some technical results

Let V be a finite-dimensional vector space. In this subsection we study various properties of specific subsets in $J_\infty(V)$.

For (x, y) in $V \times V$, we denote by $\nu_{x,y}$ the arc $t \mapsto x + ty$ of $J_\infty(V)$. Let m be in \mathbb{N}^* . We denote by $\nu_{x,y,m}$ the image of $\nu_{x,y}$ by the canonical projection $\pi_{\infty,m}^V$ from $J_\infty(V)$ to $J_m(V)$. Let \mathbf{K} be a connected closed subgroup of $\mathrm{GL}(V)$. The \mathbf{K} -action on V extends to a \mathbf{K} -action on $J_m(V)$ which is compatible with the canonical projections $\pi_{\infty,m}^V$. Let X be a \mathbf{K} -invariant irreducible closed cone in V . We suppose that X is a complete intersection in V with rational singularities and we suppose that X is a finite union of \mathbf{K} -orbits. Let N be the dimension of X and let r be the codimension of X in V .

We now use techniques developed in [Mu, Theorem 3.2] in order to prove Proposition 28. We denote by Z the union of \mathbf{K} -orbits in X which are not of maximal dimension. Let B_X be the blowing up of V whose centre is X and let

$$\tau : B_X \longrightarrow V$$

be the morphism of the blowing-up. As X is a complete intersection in V , $\tau^{-1}(X)$ is an integral divisor on B_X and is locally a complete intersection. Moreover, there exists a regular action of \mathbf{K} on B_X for which τ is a \mathbf{K} -equivariant morphism. By the theorem of embedded desingularization of Hironaka [Hi], there exists a desingularization $(Y, \tilde{\tau})$ of B_X such that $(\tau \circ \tilde{\tau})^{-1}(X)$ is a divisor with normal crossings. Moreover, we can find $(Y, \tilde{\tau})$ such that there exists a regular action of \mathbf{K} on Y for which $\tilde{\tau}$ is an equivariant morphism, and $(\tau \circ \tilde{\tau})^{-1}(X)$ is a \mathbf{K} -invariant divisor. Set $\gamma := \tau \circ \tilde{\tau}$. Denoting by E_1, \dots, E_t the irreducible components of $\gamma^{-1}(X)$, we can assume that the following conditions are fulfilled:

- (a) E_1 is the only prime divisor dominating X ;
- (b) the divisor $\gamma^{-1}(X)$ is equal to $\sum_{i=1}^t a_i E_i$;
- (c) the discrepancy W of γ is equal to $\sum_{i=1}^t b_i E_i$,
- (d) a_1 is equal to 1 and $b_1 + 1$ is equal to r ; and
- (e) $\gamma^{-1}(Z)$ is contained in the union of E_2, \dots, E_t ;

since $\tau^{-1}(X)$ is an integral divisor on B_X and \mathbf{K} has finitely many orbits in X . By condition (b), E_1, \dots, E_t are \mathbf{K} -invariant. Moreover, by condition (e), Z is the image by γ of the union of E_2, \dots, E_t . So there exist $c_2, \dots, c_t \in \mathbb{N}$, such that $\gamma^{-1}(Z) = \sum_{i=2}^t c_i E_i$. For $m \in \mathbb{N} \cup \{\infty\}$, let

$$\gamma_m : J_m(Y) \longrightarrow J_m(V)$$

be the morphism induced by γ . In the remainder of this subsection, for $m \in \mathbb{N} \cup \{\infty\}$ and $n \in \mathbb{N}$ with $m \geq n$, $\pi_{m,n}$ refers to the canonical morphism $J_m(Y) \rightarrow J_n(Y)$.

As X has rational singularities, the canonical injection from its canonical module to its dualizing module is an isomorphism by [Fl, Satz 1.1]. But its canonical module is locally free of rank 1 since X is Gorenstein as a complete intersection in V [Br, Theorem 3.3.7 and Prop. 3.1.20]. So X has canonical singularities. Moreover, by [Mu, Theorem 2.1], $b_i \geq ra_i$ for $i = 1, \dots, t$. Then by the above condition (d), we have $b_i \geq (b_1 + 1)a_i$ for $i = 2, \dots, t$. For $k \in \mathbb{N}$, for J a nonempty subset of $\{1, \dots, t\}$ and for J' a nonempty subset of J , we denote by $\psi_{k,J,J'}$ the affine functional on $\mathbb{Q}^{|J|}$,

$$(\alpha_i, i \in J) \mapsto (N - |J'|)(m + 1) + \sum_{i \in J} (b_i + 1)\alpha_i - \sum_{i \in J \setminus J'} \alpha_i + k \sum_{i \in J} c_i \alpha_i.$$

For $p \in \mathbb{N}^*$, we set

$$A_{J,J',p} := \left\{ (\alpha_i, i \in J) \in \mathbb{N}^{|J|} \mid \begin{array}{l} \sum_{i \in J} a_i \alpha_i \geq p \text{ and } \alpha_i \geq m + 1, i \in J' \\ \text{and } 1 \leq \alpha_i \leq m, i \in J \setminus J' \end{array} \right\}$$

and we denote by $\nu_{k,J,J',p}$ the minimal value of $\psi_{k,J,J'}$ on $A_{J,J',p}$. The two assertions of Lemma 26 are straightforward from the definitions and the preceding inequalities: $b_i \geq (b_1 + 1)a_i$ for $i = 2, \dots, t$. Their verifications are left to the reader.

Lemma 26. *For $k \in \mathbb{N}^*$, $p \geq m + 1$, $J \subseteq \{1, \dots, t\}$ and J' a nonempty subset of J , we have:*

- (i) $\nu_{0,J,J',p} = p(b_1 + 1) + (N - 1)(m + 1) = \nu_{k,J,J',p}$.
- (ii) $\nu_{k,\{1\},\{1\},p} \leq \nu_{k,J,J',p}$.

For $J \subseteq \{1, \dots, t\}$, we set

$$E_J^0 := \bigcap_{i \in J} E_i \setminus \bigcup_{i \notin J} E_i.$$

Let m be in \mathbb{N}^* and let S be a nonempty constructible subset of $J_m(V)$. For $\alpha = (\alpha_i)_{i \in J} \in (\mathbb{N}^*)^{|J|}$, we denote by J_α the subset of elements $i \in J$ such that $\alpha_i \geq m + 1$ and we set

$$T_\alpha := \{\nu \in \pi_{m,0}^{-1}(E_J^0) \cap \gamma_m^{-1}(S) \mid F_{E_i}(\nu) = \alpha_i, i \in J \setminus J_\alpha\}.$$

For $q \in \mathbb{N} \cup \{\infty\}$, with $q \geq m + 1$, we set

$$S_\alpha^q := \{\nu \in \pi_{q,0}^{-1}(E_J^0) \cap (\gamma_m \circ \pi_{q,m})^{-1}(S) \mid F_{E_i}(\nu) = \alpha_i, i \in J\}.$$

NILPOTENT BICONE

Lemma 27. Fix $J \subseteq \{1, \dots, t\}$, $q \geq m + 1$ and $\alpha = (\alpha_i)_{i \in J} \in (\mathbb{N}^*)^{|J|}$. Suppose that S_α^q is a nonempty set. Then S_α^q is a locally trivial fibration over T_α whose fibre is isomorphic to

$$\mathbb{C}^{(q-m)(N-|J_\alpha|)} \times (\mathbb{C}^*)^{|J_\alpha|} \times \mathbb{C}^{q|J_\alpha| - \sum_{i \in J_\alpha} \alpha_i}.$$

Proof. We follow the proof of [Cr, Prop. 2.5]. Since the divisor $\gamma^{-1}(X) = \sum_{i=1}^t a_i E_i$ on Y has only simple normal crossing, for any y in Y , there exists a neighbourhood U of y in Y with global coordinates z_1, \dots, z_N on U for which a local defining equation for $\gamma^{-1}(X)$ is given by

$$g = z_1^{\alpha_1} \cdots z_{j_y}^{\alpha_{j_y}}, \quad (4)$$

for some $j_y \leq N$. We cover $Y = \bigcup U$ by finitely many charts on which $\gamma^{-1}(X)$ has a local equation of the form (4), and we lift to cover $J_q(Y) = \bigcup \pi_{q,0}^{-1}(U)$. Hence, S_α^q is covered by the subsets

$$U_\alpha^q := \bigcap_{i \in J} F_{E_i}^{-1}(\alpha_i) \cap \pi_{q,0}^{-1}(U) \cap \pi_{q,m}^{-1}(\gamma_m^{-1}(S)).$$

As $\alpha_i \geq 1$, the subset U_α^q is contained in $\pi_{q,0}^{-1}(E_j^0)$. Thus, the subsets U_α^q are open subsets of S_α^q . Let U_α^q be such an open subset of S_α^q which is nonempty. If J is not contained in $\{1, \dots, j_y\}$ then $E_j^0 \cap U$ is empty, and so U_α^q . So J is contained in $\{1, \dots, j_y\}$.

For $\nu \in \pi_{q,0}^{-1}(U)$, we can view ν as an N -tuple $(f_1(z), \dots, f_N(z))$ of polynomials of degree at most q with zero constant term. We continue to argue as in the proof of [Cr, Prop. 2.5]. We see that $\nu \in F_{E_i}^{-1}(\alpha_i)$ if and only if the truncation of $f_i(z)$ to degree α_i is of the form $c_{\alpha_i} z^{\alpha_i}$ where c_{α_i} is different from 0. Then we obtain $N - |J|$ polynomials of degree q with zero constant term, and, for each $j \in J$, a polynomial of the form

$$f_j(z) = 0 + \cdots + 0 + c_{\alpha_j} z^{\alpha_j} + c_{\alpha_{j+1}} z^{\alpha_{j+1}} + \cdots + c_{\alpha_q} z^{\alpha_q},$$

with $c_{\alpha_j} \in \mathbb{C}^*$ and $c_k \in \mathbb{C}$ for $k > j$. So, when we cut the m first terms of each polynomial $f_j(z)$, the space of all such N -tuples so obtained is isomorphic to

$$\mathbb{C}^{(q-m)(N-|J_\alpha|)} \times (\mathbb{C}^*)^{|J_\alpha|} \times \mathbb{C}^{q|J_\alpha| - \sum_{i \in J_\alpha} \alpha_i}.$$

As a consequence, U_α^q is isomorphic to

$$(\pi_{m,0}^{-1}(U) \cap T_\alpha) \times \mathbb{C}^{(q-m)(N-|J_\alpha|)} \times (\mathbb{C}^*)^{|J_\alpha|} \times \mathbb{C}^{q|J_\alpha| - \sum_{i \in J_\alpha} \alpha_i},$$

whence the lemma. \square

Let S be a constructible subset of $J_m(V)$. For $p \geq m + 1$, we set

$$S_{\infty,m,p} := \{\nu \in (\pi_{\infty,m}^V)^{-1}(S) \mid F_X(\nu) \geq p\}.$$

Then, for $k > 0$, we set

$$\mathbf{I}_{m,p,k} := h \left(\int_{S_{\infty,m,p}} \mathbb{L}^{-kF_Z(\nu)} d\mu_V(\nu) \right).$$

Proposition 28. *Let m, k be in \mathbb{N}^* and let S be a constructible subset of $J_m(V)$. We suppose that the image by $\pi_{m,0}^V$ of S is dense in X and we suppose that S is \mathbf{K} -invariant.*

- (i) *We suppose that the image by $\pi_{m,0}^V$ of any irreducible component of maximal dimension of S is dense in X . Then the highest degree terms of $\mathbf{I}_{m,p,k}$ do not depend on k for p big enough.*
- (ii) *We suppose that the highest degree terms of $\mathbf{I}_{m,p,k}$ do not depend on k for p big enough. Then, for p big enough, the highest degree terms of $h \circ \mu_V(S_{\infty,m,p})$ only depend on the irreducible components of S whose image by $\pi_{m,0}^V$ is dense in X .*

Proof. We apply the transformation rule for the motivic integrals to $\mathbf{I}_{m,p,k}$. Namely, by [DL2, Lemma 3.3], we have

$$\mathbf{I}_{m,p,k} = h \left(\int_{\gamma_{\infty}^{-1}(S_{\infty,m,p})} \mathbb{L}^{-F_W(\nu) - k \sum_{i=2}^t c_i F_{E_i}(\nu)} d\mu_Y(\nu) \right). \quad (5)$$

For $J \subseteq \{1, \dots, t\}$ and $\alpha = (\alpha_i)_{i \in J} \in (\mathbb{N}^*)^{|J|}$, S_{α}^{∞} is the inverse image by $\pi_{\infty,q}$ of S_{α}^q for any $q \geq m+1$. So, by Proposition 25(1), $h \circ \mu_Y(S_{\alpha}^{\infty}) = h([S_{\alpha}^q])(uv)^{-N(q+1)}$. In addition, by Lemma 27, the subset S_{α}^q is a locally trivial fibration over T_{α} whose fibre is isomorphic to

$$\mathbb{C}^{(q-m)(N-|J_{\alpha}|)} \times (\mathbb{C}^*)^{|J_{\alpha}|} \times \mathbb{C}^{q|J_{\alpha}| - \sum_{i \in J_{\alpha}} \alpha_i}.$$

Then we deduce from [Cr, Theorem 3.2(iii)] the relation

$$\begin{aligned} h \circ \mu_Y(S_{\alpha}^{\infty}) &= h([T_{\alpha}])(uv-1)^{|J_{\alpha}|} \\ &= (uv)^{-|J_{\alpha}| - (N-|J_{\alpha}|)(m+1) - \sum_{i \in J_{\alpha}} \alpha_i}. \end{aligned} \quad (6)$$

As the subset $\gamma^{-1}(X)$ is equal to the union of E_1, \dots, E_t , the subset $\gamma_{\infty}^{-1}(S_{\infty,m,p})$ is given by the equality

$$\gamma_{\infty}^{-1}(S_{\infty,m,p}) = \bigcup_{\substack{J \subseteq \{1, \dots, t\} \\ J \neq \emptyset}} \bigcup_{\substack{(\alpha_i, i \in J) \in (\mathbb{N} \setminus \{0\})^{|J|} \\ \sum_{i \in J} \alpha_i \alpha_i \geq p}} S_{\alpha}^{\infty}$$

for any p . We set

$$M := 1 + m \sum_{i=1}^t a_i.$$

In particular, when $p \geq M$, J_{α} is not empty as soon as $\sum_{i \in J} a_i \alpha_i \geq p$. For any $p \geq M$ and any $k \in \mathbb{N}$, we have the equality

$$\mathbf{I}_{m,p,k} = \sum_{\substack{J \subseteq \{1, \dots, t\} \\ J \neq \emptyset}} \sum_{\substack{\alpha_i, i \in J \\ \sum_{i \in J} a_i \alpha_i \geq p, \alpha_i \geq 1}} \mathbf{S}_{J, \alpha, k}$$

NILPOTENT BICONE

where

$$\begin{aligned} \mathbf{S}_{J,\alpha,k} &:= h([T_\alpha])(uv)^{-(N-|J_\alpha|)(m+1)+\sum_{i \in J \setminus J_\alpha} \alpha_i} \\ &\quad \times (uv-1)^{|J_\alpha|} (uv)^{-|J_\alpha|} (uv)^{-\sum_{i \in J} \alpha_i (b_i+1) - k \sum_{i \in J \cap \{2, \dots, t\}} c_i \alpha_i}. \end{aligned} \quad (7)$$

For J nonempty subset in $\{1, \dots, t\}$, we set

$$\mathbf{S}'_{J,p,k} = \sum_{\substack{\alpha_i, i \in J \\ \sum_{i \in J} \alpha_i \alpha_i \geq p, \alpha_i \geq 1}} \mathbf{S}_{J,\alpha,k}.$$

Thus $\mathbf{I}_{m,p,k}$ is the sum of the $\mathbf{S}'_{J,p,k}$. For any J , the highest degree terms in $\mathbf{S}'_{J,p,k}$ are the terms $\mathbf{S}_{J,\alpha,p,k}$ for which the number

$$\dim T_\alpha - (N - |J_\alpha|)(m+1) + \sum_{i \in J \setminus J_\alpha} \alpha_i - \sum_{i \in J} \alpha_i (b_i + 1) - k \sum_{i \in J \cap \{2, \dots, t\}} c_i \alpha_i$$

is maximal. In particular, for $J = \{1\}$, the highest degree term of $\mathbf{S}'_{J,p,k}$ does not depend on k .

(i) By hypothesis, the image by $\pi_{m,0}$ of any irreducible component of maximal dimension of $\gamma_m^{-1}(S)$ is dense in E_1 . Therefore, for $J \neq \{1\}$, the degree of

$$h([\pi_{m,0}^{-1}(E_J^0) \cap \gamma_m^{-1}(S)])$$

is strictly smaller than the degree of the same expression with $J = \{1\}$. As a consequence, when $J \neq \{1\}$, for $\alpha = (\alpha_i)_{i \in J} \in (\mathbb{N}^*)^{|J|}$, the degree of $h([T_\alpha])$ is strictly smaller than the degree of

$$h([\pi_{m,0}^{-1}(E_{\{1\}}^0) \cap \gamma_m^{-1}(S)]).$$

Hence by Lemma 26 and relation (7), when p is big enough, the highest degree term of $\mathbf{I}_{m,p,k}$ is the highest degree term of $\mathbf{S}'_{\{1\},p,k}$. In particular, it does not depend on k .

(ii) Denote by \tilde{S} the union of irreducible components of S whose image by $\pi_{m,0}^V$ is dense in X . By hypothesis, \tilde{S} is a nonempty set. We set

$$\tilde{S}_{\infty,m,p} := \{\nu \in (\pi_{\infty,m}^V)^{-1}(\tilde{S}) \mid F_X(\nu) \geq p\}$$

and

$$\tilde{\mathbf{I}}_{m,p,k} := h\left(\int_{\tilde{S}_{\infty,m,p}} \mathbb{L}^{-kF_Z(\nu)} d\mu_V(\nu)\right)$$

for $k > 0$. It suffices to prove that the highest degree terms of $\mathbf{I}_{m,p,k}$ and $\tilde{\mathbf{I}}_{m,p,k}$ are the same for p big enough. Indeed, if so, we obtain the expected result with $k = 0$ in $\mathbf{I}_{m,p,k}$ and $\tilde{\mathbf{I}}_{m,p,k}$. By (i), the highest degree term of $\tilde{\mathbf{I}}_{m,p,k}$ does not depend on k for p big enough. In addition, by hypothesis, the highest degree term of $\mathbf{I}_{m,p,k}$ does not depend on k for p big enough. Fix p big enough such that the highest degree terms of both $\tilde{\mathbf{I}}_{m,p,k}$ and $\mathbf{I}_{m,p,k}$ do not depend on k . As S and \tilde{S}

are \mathbf{K} -invariant, the function $F_Z(\nu)$ is positive for any ν in $S_{\infty,m,p} \setminus \tilde{S}_{\infty,m,p}$ since \mathbf{K} has finitely many orbits in X . So we get

$$\lim_{k \rightarrow +\infty} h \left(\int_{S_{\infty,m,p} \setminus \tilde{S}_{\infty,m,p}} \mathbb{L}^{-kF_Z(\nu)} d\mu_V(\nu) \right) = 0.$$

Hence for k big enough, the highest degree terms of $\mathbf{I}_{m,p,k}$ and $\tilde{\mathbf{I}}_{m,p,k}$ are the same. \square

Let m, p be in \mathbb{N} with $p \geq m + 1$, and let S be a constructible subset of $J_m(V)$. In view of Proposition 28, we wish to investigate the highest degree term of $h \circ \mu_V(S_{\infty,m,p})$. For q in $\mathbb{N} \cup \{\infty\}$ and $q \geq n$, we denote by $\pi_{q,n}^X$ the canonical morphism from $J_q(X)$ to $J_n(X)$. The following lemma is easy and helpful for Lemma 30.

Lemma 29.

- (i) If $\nu \in J_q(X)$ for $q \in \mathbb{N} \cup \{\infty\}$, then $\pi_{q,n}^V(\nu) = \pi_{q,n}^X(\nu)$ for any $n \leq q$.
- (ii) For $\nu \in J_{\infty}(V)$ and $q \in \mathbb{N}^*$, $\pi_{\infty,q-1}^V(\nu) \in J_{q-1}(X)$ if and only if $F_X(\nu) \geq q$.

We suppose that S satisfies the following conditions:

- (1) S is contained in $J_m(X)$;
- (2) the image by $\pi_{m,0}^V$ of S is dense in X ; and
- (3) for any $\nu \in S$, the fibre $(\pi_{p-1,m}^X)^{-1}(\nu)$ is a nonempty set.

Denote by \tilde{S} the union of irreducible components of S whose image by $\pi_{m,0}^V$ is dense in X . By conditions (1) and (2), \tilde{S} is a nonempty constructible subset of $J_m(X)$. Let d and \tilde{d} be the dimensions of S and \tilde{S} respectively, and let c and \tilde{c} be the number of irreducible components of maximal dimensions of S and \tilde{S} respectively.

Denote by X_{reg} the smooth part of X . Notice that when S is contained in X_{reg} , condition (3) is automatically satisfied.

Lemma 30.

- (i) Let T be an irreducible component of S whose image by $\pi_{m,0}^V$ is dense in X . Then $(\pi_{p-1,m}^X)^{-1}(T)$ has dimension $\dim T + (N-r)(p-1-m)$.
- (ii) The highest degree term of $h \circ \mu_V(\tilde{S}_{\infty,m,p})$ is $\tilde{c}(uv)^{\tilde{d} - (N-r)(m+1) - pr}$.
- (iii) The degree of $h \circ \mu_V(S_{\infty,m,p})$ is at least $d - (N-r)(m+1) - pr$.
- (iv) Suppose that the highest degree term of $h \circ \mu_V(S_{\infty,m,p})$ is

$$\tilde{c}(uv)^{\tilde{d} - (N-r)(m+1) - pr}.$$

Then $d = \tilde{d}$ and $c = \tilde{c}$.

Proof. (i) By hypothesis, the preimage of $T \cap (\pi_{m,0}^X)^{-1}(X_{\text{reg}})$ by $\pi_{p-1,m}^X$ is a locally trivial fibration over $T \cap (\pi_{m,0}^X)^{-1}(X_{\text{reg}})$ with fibre $\mathbb{A}^{(N-r)(p-1-m)}$, whence (i).

(ii) Denote by $S^{(p-1)}$ and $\tilde{S}^{(p-1)}$ the preimage by $\pi_{p-1,m}^X$ of S and \tilde{S} respectively. Using Lemma 29, we can readily check that the relation $\tilde{S}_{\infty,m,p} =$

NILPOTENT BICONE

$(\pi_{\infty, p-1}^V)^{-1}(\tilde{S}^{(p-1)})$ holds, since S is contained in $J_m(X)$. Hence, by Proposition 25(1), we have

$$h \circ \mu_V(\tilde{S}_{\infty, m, p}) = h([\tilde{S}^{(p-1)}])(uv)^{-Np}. \quad (8)$$

By definition of \tilde{S} , the subsets

$$(\pi_{\infty, p-1}^X)^{-1}(\tilde{S}^{(p-1)}) \quad \text{and} \quad (\pi_{\infty, m}^X)^{-1}(\tilde{S})$$

are stable sets with respect to μ_X at levels $p-1$ and m , respectively. Then, by Proposition 25(1), applied to μ_X , we get

$$\begin{aligned} h \circ \mu_X((\pi_{\infty, p-1}^X)^{-1}(\tilde{S}^{(p-1)})) &= h([\tilde{S}^{(p-1)}])(uv)^{-(N-r)p}, \\ h \circ \mu_X((\pi_{\infty, m}^X)^{-1}(\tilde{S})) &= h([\tilde{S}]) (uv)^{-(N-r)(m+1)}. \end{aligned}$$

As $(\pi_{\infty, p-1}^X)^{-1}(\tilde{S}^{(p-1)})$ and $(\pi_{\infty, m}^X)^{-1}(\tilde{S})$ coincide, we deduce

$$h([\tilde{S}^{(p-1)}]) = h([\tilde{S}]) (uv)^{(N-r)(p-1-m)}.$$

But the highest degree term of $h([\tilde{S}])$ is $\tilde{c}(uv)^{\tilde{d}}$, whence (ii) by relation (8).

(iii) Analogously to (ii), we can write

$$h \circ \mu_V(S_{\infty, m, p}) = h([S^{(p-1)}])(uv)^{-Np}. \quad (9)$$

Let T be an irreducible component of S of dimension d . As S satisfies condition (3), we have $\pi_{p-1, m}^X((\pi_{p-1, m}^X)^{-1}(T)) = T$. So there is an irreducible component T' of $(\pi_{p-1, m}^X)^{-1}(T)$ whose image by $\pi_{p-1, m}^X$ is dense in T . As the generic fibre of $\pi_{p-1, m}^X : J_{p-1}(X) \rightarrow J_m(X)$ has dimension $(N-r)(p-1-m)$, the dimension of T' is at least $d + (N-r)(p-1-m)$. Hence, by relation (9), the degree of $h \circ \mu_V(S_{\infty, m, p})$ is at least $d - (N-r)(m+1) - pr$.

(iv) By relation (9), the hypothesis of (iv) means that the highest degree term of $h([S^{(p-1)}])$ is $\tilde{c}(uv)^{\tilde{d} + (N-r)(p-1-m)}$. In particular, $S^{(p-1)}$ has dimension $\tilde{d} + (N-r)(p-1-m)$ and the number of its irreducible components of maximal dimension is \tilde{c} . If $d > \tilde{d}$, then (iii) gives a contradiction. Hence $d = \tilde{d}$. So any irreducible component of maximal dimension of \tilde{S} is an irreducible component of maximal dimension of S . Hence, $c \geq \tilde{c}$. It remains to prove $c \leq \tilde{c}$. Let T_1, \dots, T_q be the irreducible components of S . Then

$$S^{(p-1)} = (\pi_{p-1, m}^X)^{-1}(T_1) \cup \dots \cup (\pi_{p-1, m}^X)^{-1}(T_q).$$

By condition (3), $\pi_{p-1, m}^X((\pi_{p-1, m}^X)^{-1}(T_i)) = T_i$ for any $i = 1, \dots, q$. Consequently,

$$\pi_{p-1, m}^X(T_i) \not\subseteq \pi_{p-1, m}^X(T_j),$$

for any $1 \leq i, j \leq q$ with $i \neq j$. Therefore there is an injection from the set of irreducible components of S into the set of irreducible components of $S^{(p-1)}$. Moreover, by (i), any irreducible component of S whose image by $\pi_{m, 0}^V$ is dense in X provides an irreducible component of $S^{(p-1)}$ of maximal dimension, since $d = \tilde{d}$. So $\tilde{c} \geq c$, since \tilde{c} is the number of the irreducible components of maximal dimension of $S^{(p-1)}$. \square

3.4. Key proposition

We keep the notations of the previous subsection. We explicitly construct in this subsection a sequence of subsets in $J_m(X)$ for $m \geq 1$, to which we apply the results of Subsection 3.3. Here, for q in $\mathbb{N} \cup \{\infty\}$ and $q \geq m$, $\pi_{q,m}$ refers to the canonical projection from $J_q(V)$ to $J_m(V)$. As V is a vector space, there is a canonical injection from $J_m(V)$ into $J_{m+1}(V)$. Furthermore, this injection is a closed immersion. The first projection from $V \times V$ to V is denoted by ϖ_1 . Let T be a closed bicone of $X \times V$ satisfying the following two conditions:

- (1) T is \mathbf{K} -invariant under the diagonal action of \mathbf{K} in $V \times V$; and
- (2) for any (x, y) in T , y is a tangent vector of X at x .

We define by induction on m a subset C_m of $J_m(V)$.

For $m = 1$, we denote by C_1 the image of T by the map $(x, y) \mapsto \nu_{x,y,1}$.

Let us suppose that the subset C_m of $J_m(V)$ is defined for some $m \geq 1$. Let C'_{m+1} be the image of C_m by the canonical injection from $J_m(V)$ into $J_{m+1}(V)$. For any $p \geq m + 1$, we set

$$C'_{\infty, m+1, p} := \{\nu \in \pi_{\infty, m+1}^{-1}(C'_{m+1}) \mid F_X(\nu) \geq p\}.$$

Then we set

$$C_{m+1} := \bigcap_{p \geq m+1} \overline{\pi_{\infty, m+1}(C'_{\infty, m+1, p})},$$

where \overline{C} denotes the closure of the subset C in $J_{m+1}(V)$. Thus, C_{m+1} is a closed subset of $J_{m+1}(V)$.

For $m \geq 1$, we set

$$D_m := \{(x, y) \in X \times V \mid \nu_{x,y,m} \in C_m\}.$$

Lemma 31. *Let m be in \mathbb{N}^* .*

- (i) *If $m \geq 2$, then C_m is the closure of $\pi_{\infty, m}(C'_{\infty, m, p})$ in $J_m(V)$ for p big enough.*
- (ii) *The subset C_m is the image of D_m by the map $(x, y) \mapsto \nu_{x,y,m}$. Moreover, C_m is contained in $J_m(X)$.*
- (iii) *If $m \geq 2$, then for $p \geq m + 1$ and for $\nu \in \pi_{\infty, m}(C'_{\infty, m, p})$, the fibre at ν of the canonical morphism from $J_{p-1}(X)$ to $J_m(X)$ is a nonempty set.*
- (iv) *The set C_m is invariant under the action of \mathbf{K} in $J_m(V)$.*
- (v) *If $\varpi_1(T)$ is equal to X , then the image by $\pi_{m,0}$ of C_m is equal to X .*

Proof. (i) By definition, the sequence

$$\pi_{\infty, m}(C'_{\infty, m, m}), \pi_{\infty, m}(C'_{\infty, m, m+1}), \dots$$

is weakly decreasing. Therefore, by Noetherianity, the weakly decreasing sequence

$$\overline{\pi_{\infty, m}(C'_{\infty, m, m})}, \overline{\pi_{\infty, m}(C'_{\infty, m, m+1})}, \dots$$

of closed subsets in $J_m(V)$ is stationary. Hence, for p big enough, C_m is the closure of $\pi_{\infty, m}(C'_{\infty, m, p})$ in $J_m(V)$.

NILPOTENT BICONE

(ii) It suffices to prove that C_m is contained in the image of D_m by the map $(x, y) \mapsto \nu_{x,y,m}$ and that C_m belongs to $J_m(X)$. We prove the statements by induction on m . By definition, C_1 is the image of T by the map $(x, y) \mapsto \nu_{x,y,1}$ so C_1 is contained in $J_1(X)$ by condition (2). So D_1 is equal to T , whence the two statements for $m = 1$. Let us suppose that C_m is the image of D_m by the map $(x, y) \mapsto \nu_{x,y,m}$ and let us suppose that C_m is contained in $J_m(X)$ for some $m \geq 1$. Then C'_{m+1} is the image of D_m by the map $(x, y) \mapsto \nu_{x,y,m+1}$. By definition, (x, y) belongs to D_{m+1} if and only if $\nu_{x,y,m+1}$ belongs to C_{m+1} . As C_{m+1} is contained in C'_{m+1} , we deduce that C_{m+1} is contained in the image of D_{m+1} by the map $(x, y) \mapsto \nu_{x,y,m+1}$. In addition, for $p \geq m + 2$, $\pi_{\infty,m+1}(C'_{\infty,m+1,p})$ is contained in $J_{m+1}(X)$, by definition of $C'_{\infty,m+1,p}$. Hence, C_{m+1} is contained in $J_{m+1}(X)$ since $J_{m+1}(X)$ is closed in $J_{m+1}(V)$.

(iii) Let ν be in $\pi_{\infty,m}(C'_{\infty,m,p})$. Then, there is ν' in $C'_{\infty,m,p}$ such that $\pi_{\infty,m}(\nu') = \nu$. As ν' belongs to $C'_{\infty,m,p}$, we have $F_X(\nu') \geq p$. Hence, by Lemma 29(ii), $\pi_{\infty,p-1}(\nu')$ belongs to $J_{p-1}(X)$. In addition, since $m \geq 2$, $\pi_{\infty,m}(C'_{\infty,m,p})$ is contained in C_m . So, by (ii), ν belongs to $J_m(X)$. Therefore, by Lemma 29(i), $\pi_{\infty,p-1}(\nu')$ is in the fibre at ν of the canonical morphism from $J_{p-1}(X)$ to $J_m(X)$.

(iv) We prove the statement by induction on m . As T is \mathbf{K} -invariant by condition (1), C_1 is \mathbf{K} -invariant. We suppose $m \geq 2$ and we suppose that C_m is \mathbf{K} -invariant. Then C'_{m+1} is \mathbf{K} -invariant. So, $\pi_{\infty,m+1}^{-1}(C'_{m+1})$ is \mathbf{K} -invariant. As X is \mathbf{K} -invariant, the function F_X is \mathbf{K} -invariant. So for any $p \geq m + 1$, $C'_{\infty,m+1,p}$ is \mathbf{K} -invariant. Hence, $\pi_{\infty,m+1}(C'_{\infty,m+1,p})$ and C_{m+1} are \mathbf{K} -invariant.

(v) We suppose that $\varpi_1(T)$ is equal to X . Then, for any x in X , $(x, 0)$ belongs to T since T is a closed bicone. As a consequence, for any $m \in \mathbb{N}^*$, C_m contains $\nu_{x,0,m}$. Hence, $\pi_{m,0}(C_m)$ is equal to X . \square

Proposition 32.

(i) For m big enough, we have

$$D_m = \{(x, y) \in T \mid x + ty \in X, \forall t \in \mathbb{C}\}.$$

(ii) We suppose that the image by ϖ_1 of any irreducible component of maximal dimension of T is equal to X . Then, for $m \geq 1$, the image by ϖ_1 of any irreducible component of maximal dimension of D_m is dense in X .

Proof. (i) Let m be in \mathbb{N}^* such that $m - 1$ is strictly bigger than the degree of any element of a generating family of the ideal of definition of X in the algebra of polynomial functions on V . Let (x, y) be in T . If $x + ty$ belongs to X for any t in \mathbb{C} , then $F_X(\nu_{x,y})$ is equal to ∞ . In particular, C_m contains $\nu_{x,y,m}$ and D_m contains (x, y) . Conversely, let us suppose that D_m contains (x, y) . Then T contains (x, y) since T contains D_i for any $i \in \mathbb{N}^*$. Moreover, for any φ in the ideal of definition of X , the function $t \mapsto \varphi(x + ty)$ is divisible by t^m . So, by the choice of m , $\varphi(x + ty)$ is equal to 0, for any $t \in \mathbb{C}$. In other words, $x + ty$ belongs to X for any $t \in \mathbb{C}$.

(ii) By Lemma 31(ii), the statement is equivalent to the following statement:

For any $m \in \mathbb{N}^$, the image by $\pi_{m,0}$ of any irreducible component of maximal dimension of C_m is dense in X .*

We prove this statement by induction on m . It is true for $m = 1$ by hypothesis and by definition of C_1 . Suppose $m \geq 2$ and suppose the statement true for $m - 1$. Let \tilde{C}_m be the union of irreducible components of C_m whose image by $\pi_{m,0}$ is dense in X . By Lemma 31(v), \tilde{C}_m is not empty. Let d and \tilde{d} be the dimensions of C_m and \tilde{C}_m respectively, and let c and \tilde{c} be the number of their irreducible components of maximal dimension. It is enough to prove the equalities $d = \tilde{d}$ and $c = \tilde{c}$. In fact, in this case, any irreducible component of maximal dimension of C_m is an irreducible component of \tilde{C}_m . By Lemma 31(ii), (iii) and (iv), the conditions of Lemma 30 are satisfied.

For $p \geq m + 1$, we set

$$\tilde{C}_{\infty,m,p} := \{\nu \in \pi_{\infty,m}^{-1}(\tilde{C}_m) \mid F_X(\nu) \geq p\}. \quad (10)$$

The image by $\pi_{m,0}$ of any irreducible component of \tilde{C}_m is dense in X . Hence, by Proposition 28(i), the highest degree term of the element

$$\tilde{\mathbf{I}}_{m,p,k} := h \left(\int_{\tilde{C}_{\infty,m,p}} \mathbb{L}^{-kF_Z(\nu)} d\mu_V(\nu) \right)$$

of $\mathbb{Z}[u, v][[u^{-1}, v^{-1}]]$ does not depend on k , for p big enough. On the other hand, by induction hypothesis, the image by $\pi_{\infty,m-1}$ of any irreducible component of maximal dimension of C_{m-1} is dense in X , so the image by $\pi_{\infty,m}$ of any irreducible component of maximal dimension of C'_m is dense in X . Hence, by Proposition 28(i), the highest degree term of the element

$$\mathbf{I}_{m,p,k} := h \left(\int_{C'_{\infty,m,p}} \mathbb{L}^{-kF_Z(\nu)} d\mu_V(\nu) \right)$$

of $\mathbb{Z}[u, v][[u^{-1}, v^{-1}]]$ does not depend on k , for p big enough. Finally, applying besides Lemma 31(i), we may choose p big enough such that the following conditions hold:

- (1) C_m is the closure of $\pi_{\infty,m}(C'_{\infty,m,p})$; and
- (2) the highest degree terms of $\mathbf{I}_{m,p,k}$ and $\tilde{\mathbf{I}}_{m,p,k}$ do not depend on k .

Using condition (1), we can easily check that the relation

$$C'_{\infty,m,p} = \{\nu \in \pi_{\infty,m}^{-1}(\pi_{\infty,m}(C'_{\infty,m,p})) \mid F_X(\nu) \geq p\} \quad (11)$$

holds. By Lemma 31(v), $\pi_{m,0}(C_m)$ is equal to X . So, by condition (1), the image of $\pi_{\infty,m}(C'_{\infty,m,p})$ by $\pi_{m,0}$ is dense in X . Then, since condition (2) holds, it results from Proposition 28(ii), that the highest degree term of $\mathbf{I}_{m,p,k}$ only depends on the irreducible components of $\pi_{\infty,m}(C'_{\infty,m,p})$ whose image by $\pi_{m,0}$ is dense in X .

In other words, the highest degree terms of $\mathbf{I}_{m,p,k}$ and $\tilde{\mathbf{I}}_{m,p,k}$ are the same.

By condition (2) with $k = 0$ in $\tilde{\mathbf{I}}_{m,p,k}$ and Lemma 30(ii), we deduce that the highest degree term of $\tilde{\mathbf{I}}_{m,p,k}$ is $\tilde{c}(uv)^{\tilde{d} - (N-r)(m+1) - pr}$. So the highest degree term of $\mathbf{I}_{m,p,k}$ is $\tilde{c}(uv)^{\tilde{d} - (N-r)(m+1) - pr}$, too. Hence, by condition (2) with $k = 0$ in $\mathbf{I}_{m,p,k}$, we deduce that the highest degree term of $h \circ \mu_V(C'_{\infty,m,p})$ is $\tilde{c}(uv)^{\tilde{d} - (N-r)(m+1) - pr}$. Then, by Lemma 30(iv), the two equalities $d = \tilde{d}$ and $c = \tilde{c}$ hold. \square

4. Dimension of the nilpotent bicone via motivic integration

In this section we prove the main result of this paper. As in Section 2, we suppose that \mathfrak{g} is simple and we adopt the notations of the previous sections. The aim of this section is the following theorem.

Theorem 33.

- (i) *The principal bicone of \mathfrak{g} is a reduced complete intersection of dimension $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g} + 1)$.*
- (ii) *The subscheme $\mathcal{Y}_{\mathfrak{g}}$ is a reduced complete intersection of dimension $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g}) + 2$. Moreover, any irreducible component of $\mathcal{Y}_{\mathfrak{g}}$ is the intersection of $\mathcal{Y}_{\mathfrak{g}}$ with an irreducible component of $\mathcal{X}_{\mathfrak{g}}$.*
- (iii) *The subscheme $\mathcal{Z}_{\mathfrak{g}}$ is a reduced complete intersection of dimension $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g}) + 1$.*
- (iv) *The nilpotent bicone $\mathcal{N}_{\mathfrak{g}}$ is a complete intersection of dimension $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g})$.*

Let us give a brief description of our approach to prove Theorem 33. We plan to apply Corollary 23. Namely, we intend to prove that any irreducible component of maximal dimension of $\mathcal{X}_{\mathfrak{g}}$ satisfies conditions (1) and (2) of Corollary 23. Then we will deduce Theorem 33(i), from Lemma 18(i). Next, the statements (ii), (iii) and (iv) of Theorem 33 will be mostly consequences of statement (i). The main point is therefore to study the images by ϖ_1 and ϖ_2 of the irreducible components of maximal dimension of $\mathcal{X}_{\mathfrak{g}}$, $\mathcal{Y}_{\mathfrak{g}}$, $\mathcal{Z}_{\mathfrak{g}}$ and $\mathcal{N}_{\mathfrak{g}}$. To process, we consider jet schemes of $\mathfrak{X}_{\mathfrak{g}}$ and $\mathfrak{N}_{\mathfrak{g}}$ and we use the results of Section 3 about motivic integration. In fact, we remark that an element $(x, y) \in \mathfrak{g} \times \mathfrak{g}$ belongs to $\mathcal{X}_{\mathfrak{g}}$ or $\mathcal{N}_{\mathfrak{g}}$ if and only if the arc $t \mapsto x + ty$ is an element of $J_{\infty}(\mathfrak{X}_{\mathfrak{g}})$ or $J_{\infty}(\mathfrak{N}_{\mathfrak{g}})$ respectively. Thus $\mathcal{X}_{\mathfrak{g}}$ and $\mathcal{N}_{\mathfrak{g}}$ can be identified to subsets of $J_{\infty}(\mathfrak{X}_{\mathfrak{g}})$ and $J_{\infty}(\mathfrak{N}_{\mathfrak{g}})$ respectively. As $\mathfrak{X}_{\mathfrak{g}}$ and $\mathfrak{N}_{\mathfrak{g}}$ are strictly contained in \mathfrak{g} , the motivic measure with respect to \mathfrak{g} of the so obtained subsets is zero by Proposition 25(2). So we cannot expect to obtain any information from these measures. That is why we are instead going to make use of the subtler construction as described in Subsection 3.4. We will obtain in this way subsets whose Hodge realization of the motivic measure with respect to \mathfrak{g} provide all the information we need about the varieties $\mathcal{X}_{\mathfrak{g}}$ and $\mathcal{N}_{\mathfrak{g}}$.

4.1. Two lemmas

In this subsection we give two lemmas useful for Subsection 4.2. We denote by θ the map

$$\mathbb{C} \times \mathfrak{g} \times \mathfrak{g} \longrightarrow \mathfrak{g} \times \mathfrak{g}, \quad (t, x, y) \mapsto (x, y + tx).$$

Lemma 34. *Let \mathcal{X} be a G -invariant irreducible closed bicone of $\mathfrak{g} \times \mathfrak{g}$. Suppose that $\varpi_1(\mathcal{X}) = \mathfrak{X}_{\mathfrak{g}}$ and that $\theta(\mathbb{C} \times \mathcal{X})$ is contained in \mathcal{X} . If \mathcal{X}_1 and \mathcal{X}_2 are two irreducible components of the nullvariety of $p_{1,1,1}$ in \mathcal{X} such that $\varpi_1(\mathcal{X}_1) = \varpi_1(\mathcal{X}_2) = \mathfrak{X}_{\mathfrak{g}}$, then $\mathcal{X}_1 = \mathcal{X}_2$.*

Proof. By hypothesis, we can suppose that \mathcal{X}_1 and \mathcal{X}_2 have codimension 1 in \mathcal{X} since \mathcal{X} is irreducible. If (x, y) belongs to the nullvariety of $p_{1,1,1}$ in \mathcal{X} , with x semisimple, then $p_{1,1,1}(x, x + y) \neq 0$, since $\langle x, x \rangle \neq 0$ by Lemma 15(i). By hypothesis, for $i = 1, 2$, $\theta(\mathbb{C} \times \mathcal{X}_i)$ is an irreducible constructible subset of \mathcal{X} which

strictly contains \mathcal{X}_i . Hence for $i = 1, 2$, $\theta(\mathbb{C} \times \mathcal{X}_i)$ contains a dense open subset of \mathcal{X} . As a result, for any (t, x, y) in a certain dense open subset of $\mathbb{C} \times \mathcal{X}_1$, x is semisimple and there exists (t', x', y') in $\mathbb{C} \times \mathcal{X}_2$ such that $\theta(t', x', y') = \theta(t, x, y)$. From this equality we deduce the equalities

$$x' = x, \quad y' - y = (t - t')x.$$

But $\langle x, x \rangle \neq 0$ since x is a semisimple element of $\mathfrak{X}_{\mathfrak{g}}$. On the other hand, $\langle x, y \rangle = \langle x, y' \rangle = 0$. So $t' = t$ and $(x, y) = (x', y')$. As a consequence, $\mathcal{X}_1 = \mathcal{X}_2$. \square

Let $T\mathfrak{X}_{\mathfrak{g}}$ be the total tangent space of $\mathfrak{X}_{\mathfrak{g}}$ and let $T\mathfrak{N}_{\mathfrak{g}}$ be the total tangent space of $\mathfrak{N}_{\mathfrak{g}}$. They are closed subsets of $\mathfrak{g} \times \mathfrak{g}$. Let $T'\mathfrak{X}_{\mathfrak{g}}$ be the nullvariety of $p_{1,1,1}$ in $T\mathfrak{X}_{\mathfrak{g}}$ and let $T''\mathfrak{X}_{\mathfrak{g}}$ be the nullvariety of $p_{1,0,2}$ in $T'\mathfrak{X}_{\mathfrak{g}}$.

Lemma 35.

- (i) *The subsets $T\mathfrak{X}_{\mathfrak{g}}$ and $T\mathfrak{N}_{\mathfrak{g}}$ are G -invariant closed irreducible bicones, their images by ϖ_1 are equal to $\mathfrak{X}_{\mathfrak{g}}$ and $\mathfrak{N}_{\mathfrak{g}}$ respectively, and they have dimension $4(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g}) + 2$ and $4(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g})$ respectively.*
- (ii) *The intersection of $\varpi_1^{-1}(\mathfrak{N}_{\mathfrak{g}})$ and $T'\mathfrak{X}_{\mathfrak{g}}$ is equal to $T\mathfrak{N}_{\mathfrak{g}}$.*
- (iii) *The subset $T'\mathfrak{X}_{\mathfrak{g}}$ is irreducible of dimension $4(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g}) + 1$. Moreover, $\varpi_1(T'\mathfrak{X}_{\mathfrak{g}})$ is equal to $\mathfrak{X}_{\mathfrak{g}}$.*
- (iv) *The image of any irreducible component of $T''\mathfrak{X}_{\mathfrak{g}}$ by ϖ_1 is equal to $\mathfrak{X}_{\mathfrak{g}}$.*

Proof. (i) As $\mathfrak{X}_{\mathfrak{g}}$ and $\mathfrak{N}_{\mathfrak{g}}$ are G -invariant closed cones, $T\mathfrak{X}_{\mathfrak{g}}$ and $T\mathfrak{N}_{\mathfrak{g}}$ are G -invariant closed bicones. By Theorem 24 and Remark 7, $T\mathfrak{X}_{\mathfrak{g}}$ is irreducible and has dimension $4(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g}) + 2$. Likewise, by Theorem 24 and Remark 7, $T\mathfrak{N}_{\mathfrak{g}}$ is irreducible and has dimension $4(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g})$.

(ii) For x in $G.e$ and y in \mathfrak{g} , (x, y) belongs to $T\mathfrak{N}_{\mathfrak{g}}$ if and only if y belongs to $[x, \mathfrak{g}]$. In particular, for any x in $G.e$ and any y in \mathfrak{g} such that (x, y) belongs to $T\mathfrak{N}_{\mathfrak{g}}$, (x, y) belongs to $T'\mathfrak{X}_{\mathfrak{g}}$ since x is orthogonal to $[x, \mathfrak{g}]$. By (i), the intersection of $T\mathfrak{N}_{\mathfrak{g}}$ and $\varpi_1^{-1}(G.e)$ is dense in $T\mathfrak{N}_{\mathfrak{g}}$. So $T\mathfrak{N}_{\mathfrak{g}} \subset \varpi_1^{-1}(\mathfrak{N}_{\mathfrak{g}}) \cap T'\mathfrak{X}_{\mathfrak{g}}$. Let us prove the other inclusion. As $\mathfrak{X}_{\mathfrak{g}}$ is the nullvariety in \mathfrak{g} of $q_2, \dots, q_{\text{rk } \mathfrak{g}}$, for any x in $\mathfrak{X}_{\mathfrak{g}}$, the subspace of elements y of \mathfrak{g} such that (x, y) belongs to $T\mathfrak{X}_{\mathfrak{g}}$ is the intersection of the kernels of the differentials at x of $q_2, \dots, q_{\text{rk } \mathfrak{g}}$. By definition of the q_i (see Subsection 2.1), for any x in $\mathfrak{N}_{\mathfrak{g}}$, the subset of elements y of \mathfrak{g} such that (x, y) belongs to $T\mathfrak{X}_{\mathfrak{g}}$ is the intersection of the kernels of the differentials at x of $p_2, \dots, p_{\text{rk } \mathfrak{g}}$. Therefore, the intersection of $T'\mathfrak{X}_{\mathfrak{g}}$ and $\varpi_1^{-1}(\mathfrak{N}_{\mathfrak{g}})$ is contained in $T\mathfrak{N}_{\mathfrak{g}}$, since $\mathfrak{N}_{\mathfrak{g}}$ is the nullvariety of $p_1, \dots, p_{\text{rk } \mathfrak{g}}$ in \mathfrak{g} .

(iii) As $\mathfrak{X}_{\mathfrak{g}}$ is a cone, $T\mathfrak{X}_{\mathfrak{g}}$ contains the diagonal of $\mathfrak{X}_{\mathfrak{g}} \times \mathfrak{X}_{\mathfrak{g}}$. So $T\mathfrak{X}_{\mathfrak{g}}$ strictly contains $T'\mathfrak{X}_{\mathfrak{g}}$. Then, by (i), $T'\mathfrak{X}_{\mathfrak{g}}$ has dimension $4(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g}) + 1$. Moreover, $T'\mathfrak{X}_{\mathfrak{g}}$ is an equidimensional G -invariant closed bicone, since $p_{1,1,1}$ is bihomogeneous and G -invariant. Hence, by (ii) and Lemma 3, the image by ϖ_1 of any irreducible component of $T'\mathfrak{X}_{\mathfrak{g}}$ is equal to $\mathfrak{X}_{\mathfrak{g}}$. As $\mathfrak{X}_{\mathfrak{g}}$ is a cone, for any t in \mathbb{C} and any (x, y) in $T\mathfrak{X}_{\mathfrak{g}}$, $(x, y + tx)$ belongs to $T\mathfrak{X}_{\mathfrak{g}}$. Therefore, by Lemma 34, $T'\mathfrak{X}_{\mathfrak{g}}$ is irreducible.

(iv) By (iii), $T''\mathfrak{X}_{\mathfrak{g}}$ is equidimensional of dimension at least $4(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g})$. Moreover, since $p_{1,0,2}$ is bihomogeneous and G -invariant, $T''\mathfrak{X}_{\mathfrak{g}}$ is a G -invariant closed bicone. As (e, h) belongs to $T'\mathfrak{X}_{\mathfrak{g}}$, $T''\mathfrak{X}_{\mathfrak{g}}$ is strictly contained in $T'\mathfrak{X}_{\mathfrak{g}}$ and $T''\mathfrak{X}_{\mathfrak{g}}$ is equidimensional of dimension $4(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g})$. In addition, as (e, h) belongs to

$T\mathfrak{N}_{\mathfrak{g}} \setminus T''\mathfrak{X}_{\mathfrak{g}}$ we deduce, from (ii) and Lemma 15(i), that the intersection of $T''\mathfrak{X}_{\mathfrak{g}}$ and $\varpi_1^{-1}(\mathfrak{N}_{\mathfrak{g}})$ is equidimensional of dimension $4(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g}) - 1$. Hence, by Lemma 3, the image of any irreducible component of $T''\mathfrak{X}_{\mathfrak{g}}$ is equal to $\mathfrak{X}_{\mathfrak{g}}$. \square

4.2. Proof of Theorem 33

In this subsection we apply Proposition 32 to suitable V , X , T and \mathbf{K} in order to prove Theorem 33.

Proposition 36.

- (i) *Let \mathcal{X} be an irreducible component of maximal dimension of $\mathcal{X}_{\mathfrak{g}}$ (resp., $\mathcal{Y}_{\mathfrak{g}}$, $\mathcal{Z}_{\mathfrak{g}}$). Then $\varpi_1(\mathcal{X})$ is equal to $\mathfrak{X}_{\mathfrak{g}}$.*
- (ii) *Let \mathcal{X} be an irreducible component of maximal dimension of $\mathcal{N}_{\mathfrak{g}}$, then $\varpi_1(\mathcal{X})$ is equal to $\mathfrak{N}_{\mathfrak{g}}$.*

Proof. (i) Let \mathbf{K} be the subgroup of $\text{GL}(\mathfrak{g})$ generated by G and its homotheties. By Corollary 16, $\mathfrak{X}_{\mathfrak{g}}$ is a \mathbf{K} -invariant irreducible closed normal cone and it is a complete intersection in \mathfrak{g} with rational singularities. Moreover, \mathbf{K} has finitely many orbits in $\mathfrak{X}_{\mathfrak{g}}$. Let T be the bicone $T\mathfrak{X}_{\mathfrak{g}}$ (resp., $T'\mathfrak{X}_{\mathfrak{g}}$, $T''\mathfrak{X}_{\mathfrak{g}}$) of $\mathfrak{X}_{\mathfrak{g}} \times \mathfrak{g}$. By Lemma 35(i) (resp., (iii), (iv)), the image by ϖ_1 of any irreducible component of T is equal to $\mathfrak{X}_{\mathfrak{g}}$. Then, by Proposition 32(i), applied to $V = \mathfrak{g}$, $X = \mathfrak{X}_{\mathfrak{g}}$, T and \mathbf{K} , D_m is equal to $\mathcal{X}_{\mathfrak{g}}$ (resp., $\mathcal{Y}_{\mathfrak{g}}$, $\mathcal{Z}_{\mathfrak{g}}$) for m big enough. Hence, by Proposition 32(ii), $\varpi_1(\mathcal{X}) = \mathfrak{X}_{\mathfrak{g}}$ since $\varpi_1(\mathcal{X})$ is closed by Lemma 3.

(ii) Let \mathbf{K} be the subgroup G of $\text{GL}(\mathfrak{g})$. The cone $\mathfrak{N}_{\mathfrak{g}}$ is an irreducible closed normal cone and \mathbf{K} has finitely many orbits in $\mathfrak{N}_{\mathfrak{g}}$. Moreover, it is a complete intersection in \mathfrak{g} and by [He, Theorem A], $\mathfrak{N}_{\mathfrak{g}}$ has rational singularities. Let T be the bicone $T\mathfrak{N}_{\mathfrak{g}}$ of $\mathfrak{N}_{\mathfrak{g}} \times \mathfrak{g}$. By Lemma 35(i), T is irreducible and its image by ϖ_1 is equal to $\mathfrak{N}_{\mathfrak{g}}$. Then, by Proposition 32(i), applied to $V = \mathfrak{g}$, $X = \mathfrak{N}_{\mathfrak{g}}$, T and \mathbf{K} , D_m is equal to $\mathcal{N}_{\mathfrak{g}}$ for m big enough. Hence, by Proposition 32(ii), $\varpi_1(\mathcal{X}) = \mathfrak{N}_{\mathfrak{g}}$ since $\varpi_1(\mathcal{X})$ is closed by Lemma 3. \square

Let d be the dimension of $\mathcal{X}_{\mathfrak{g}}$.

Proposition 37. *Let \mathcal{X} be an irreducible component of dimension d of $\mathcal{X}_{\mathfrak{g}}$.*

- (i) *The dimension of $\mathcal{Y}_{\mathfrak{g}}$ is $d - 1$.*
- (ii) *The intersection of \mathcal{X} and $\mathcal{Y}_{\mathfrak{g}}$ is irreducible and has dimension $d - 1$. Moreover, this intersection is stable under the involution $(x, y) \mapsto (y, x)$.*
- (iii) *The intersection of \mathcal{X} and $\mathcal{Z}_{\mathfrak{g}}$ is an union of irreducible components of $\mathcal{Z}_{\mathfrak{g}}$ of maximal dimension. Moreover, $\mathcal{Z}_{\mathfrak{g}}$ has dimension $d - 2$.*
- (iv) *The intersection of $\mathcal{N}_{\mathfrak{g}}$ and \mathcal{X} is equidimensional of dimension $d - 3$.*
- (v) *The dimension of $\mathcal{N}_{\mathfrak{g}}$ is equal to $d - 3$.*

Proof. (i) Let \mathcal{X}' be an irreducible component of $\mathcal{Y}_{\mathfrak{g}}$ of maximal dimension. As $\mathcal{Y}_{\mathfrak{g}}$ is contained in $\mathcal{X}_{\mathfrak{g}}$, \mathcal{X}' is contained in an irreducible component \mathcal{X}'' of $\mathcal{X}_{\mathfrak{g}}$. Moreover, \mathcal{X}' is an irreducible component of the nullvariety of $p_{1,1,1}$ in \mathcal{X}'' . So \mathcal{X}' has codimension at most 1 in \mathcal{X}'' . By Proposition 36(i), $\varpi_1(\mathcal{X}')$ is equal to $\mathfrak{X}_{\mathfrak{g}}$. Hence \mathcal{X}' contains $(h, 0)$ since it is a closed bicone. As \mathcal{X}'' is an irreducible component of $\mathcal{X}_{\mathfrak{g}}$, it is invariant under the action of $\text{GL}_2(\mathbb{C})$. Hence \mathcal{X}'' contains (h, h) and so strictly contains \mathcal{X}' . So \mathcal{X}' has dimension $\dim \mathcal{X}'' - 1$. In particular,

$\dim \mathcal{X}'$ is at most $d - 1$. As any irreducible component of the nullvariety of $p_{1,1,1}$ in \mathcal{X} is contained in $\mathcal{Y}_{\mathfrak{g}}$, $\mathcal{Y}_{\mathfrak{g}}$ has dimension $d - 1$.

(ii) As $\mathcal{Y}_{\mathfrak{g}}$ is stable under the involution $(x, y) \mapsto (y, x)$ and \mathcal{X} is stable under the action of $\mathrm{GL}_2(\mathbb{C})$, their intersection is stable under the involution $(x, y) \mapsto (y, x)$. By (i), any irreducible component of the intersection of \mathcal{X} and $\mathcal{Y}_{\mathfrak{g}}$ has dimension $d - 1$. Hence by (i), Proposition 36(i) and Lemma 34, the intersection of $\mathcal{Y}_{\mathfrak{g}}$ and \mathcal{X} is irreducible since $\theta(\mathbb{C} \times \mathcal{X})$ is contained in \mathcal{X} .

(iii) Let \mathcal{Y} be the intersection of $\mathcal{Y}_{\mathfrak{g}}$ and \mathcal{X} and let \mathcal{Z} be the intersection of \mathcal{X} and $\mathcal{Z}_{\mathfrak{g}}$. By definition, \mathcal{Z} is the nullvariety of $p_{1,0,2}$ in \mathcal{Y} . Moreover, by (ii) and Proposition 36(i), $\varpi_2(\mathcal{Y})$ is equal to $\mathfrak{X}_{\mathfrak{g}}$. Hence by (i) and (ii), any irreducible component of \mathcal{Z} has dimension $d - 2$. Let \mathcal{Z}' be an irreducible component of $\mathcal{Z}_{\mathfrak{g}}$ of maximal dimension. Then $\dim \mathcal{Z}' \geq d - 2$. Suppose $\dim \mathcal{Z}' > d - 2$. As $\mathcal{Z}_{\mathfrak{g}}$ is contained in $\mathcal{Y}_{\mathfrak{g}}$, \mathcal{Z}' is an irreducible component of maximal dimension of $\mathcal{Y}_{\mathfrak{g}}$ by (i). So, by Proposition 36(i), $\varpi_2(\mathcal{Z}')$ is equal to $\mathfrak{X}_{\mathfrak{g}}$. This is impossible since $\mathcal{Z}' \subset \mathcal{Z}_{\mathfrak{g}}$. So $\dim \mathcal{Z}_{\mathfrak{g}} = d - 2$ and \mathcal{Z} is a union of irreducible components of dimension $d - 2$ of $\mathcal{Z}_{\mathfrak{g}}$.

(iv) Let \mathcal{Y} and \mathcal{Z} be as in (iii). As $\mathcal{N}_{\mathfrak{g}}$ is contained in $\mathcal{Z}_{\mathfrak{g}}$, the intersection of \mathcal{X} and $\mathcal{N}_{\mathfrak{g}}$ is contained in \mathcal{Z} . By (iii) and Proposition 36(i), any irreducible component of \mathcal{Z} is not contained in $\mathcal{N}_{\mathfrak{g}}$ and has dimension $d - 2$. The intersection of \mathcal{X} and $\mathcal{N}_{\mathfrak{g}}$ is the nullvariety of $p_{1,2,0}$ in \mathcal{Z} . Hence this intersection is equidimensional of dimension $d - 3$.

(v) As $\mathcal{N}_{\mathfrak{g}}$ is contained in $\mathcal{Z}_{\mathfrak{g}}$, any irreducible component of $\mathcal{N}_{\mathfrak{g}}$ is contained in an irreducible component of $\mathcal{Z}_{\mathfrak{g}}$. Moreover, by (iii) and Proposition 36(i), any irreducible component of dimension $d - 2$ of $\mathcal{Z}_{\mathfrak{g}}$ is not contained in $\mathcal{N}_{\mathfrak{g}}$. Hence the dimension of $\mathcal{N}_{\mathfrak{g}}$ is at most $d - 3$. So, by (iv), $\mathcal{N}_{\mathfrak{g}}$ has dimension $d - 3$. \square

Corollary 38.

- (i) *The subscheme $\mathcal{X}_{\mathfrak{g}}$ is equidimensional of dimension $3(\mathrm{b}_{\mathfrak{g}} - \mathrm{rk} \mathfrak{g} + 1)$. Moreover, any irreducible component of $\mathcal{X}_{\mathfrak{g}}$ has a nonempty intersection with the intersection of $\Omega_{\mathfrak{g}}$ and $\mathcal{Z}_{\mathfrak{g}}$.*
- (ii) *The subscheme $\mathcal{Y}_{\mathfrak{g}}$ is equidimensional of dimension $3(\mathrm{b}_{\mathfrak{g}} - \mathrm{rk} \mathfrak{g}) + 2$. Moreover, any irreducible component of $\mathcal{Y}_{\mathfrak{g}}$ has a nonempty intersection with $\Omega_{\mathfrak{g}}$.*
- (iii) *The subscheme $\mathcal{Z}_{\mathfrak{g}}$ is equidimensional of dimension $3(\mathrm{b}_{\mathfrak{g}} - \mathrm{rk} \mathfrak{g}) + 1$. Moreover, any irreducible component of $\mathcal{Z}_{\mathfrak{g}}$ has a nonempty intersection with $\Omega_{\mathfrak{g}}$.*

Proof. (i) Let \mathcal{X} be an irreducible component of dimension d of $\mathcal{X}_{\mathfrak{g}}$. Let \mathcal{Z} and \mathcal{N} be the intersections of \mathcal{X} with $\mathcal{Z}_{\mathfrak{g}}$ and $\mathcal{N}_{\mathfrak{g}}$ respectively. By Proposition 37(iii) and (iv), \mathcal{Z} and \mathcal{N} are equidimensional of dimensions $d - 2$ and $d - 3$ respectively. Hence, by Proposition 37(v), \mathcal{N} is a union of irreducible components of maximal dimension of $\mathcal{N}_{\mathfrak{g}}$. As any irreducible component of $\mathcal{N}_{\mathfrak{g}}$ is stable under the action of $\mathrm{GL}_2(\mathbb{C})$, the image by ϖ_2 of any irreducible component of \mathcal{N} is equal to $\mathfrak{N}_{\mathfrak{g}}$ by Proposition 36(ii). Moreover, each irreducible component of \mathcal{Z} contains an irreducible component of \mathcal{N} since \mathcal{N} is the nullvariety of $p_{1,2,0}$ in \mathcal{Z} . Hence, the image by ϖ_2 of any irreducible component of \mathcal{Z} is equal to $\mathfrak{N}_{\mathfrak{g}}$. By Proposition 37(iii), each irreducible component of \mathcal{Z} is an irreducible component of $\mathcal{Z}_{\mathfrak{g}}$ of max-

NILPOTENT BICONE

imal dimension. Hence, by Proposition 36(i), the image by ϖ_1 of any irreducible component of \mathcal{Z} is equal to $\mathfrak{X}_{\mathfrak{g}}$. So \mathcal{Z} satisfies conditions (1) and (2) of Corollary 23. Hence, by Corollary 23, \mathcal{Z} has a nonempty intersection with $\Omega_{\mathfrak{g}}$ and \mathcal{X} has dimension $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g} + 1)$. But, by Lemma 18(i), any irreducible component of $\mathcal{X}_{\mathfrak{g}}$ has dimension at least $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g} + 1)$. Hence $\mathcal{X}_{\mathfrak{g}}$ is equidimensional of dimension $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g} + 1)$.

(ii) By Proposition 37(ii) and Lemma 19(ii), $\mathcal{Y}_{\mathfrak{g}}$ is equidimensional of dimension $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g}) + 2$. Moreover, any irreducible component of $\mathcal{Y}_{\mathfrak{g}}$ is the intersection of $\mathcal{Y}_{\mathfrak{g}}$ and an irreducible component of $\mathcal{X}_{\mathfrak{g}}$. Hence by (i) and Corollary 23, any irreducible component of $\mathcal{Y}_{\mathfrak{g}}$ has a nonempty intersection with $\Omega_{\mathfrak{g}}$.

(iii) By Proposition 37(ii) and Lemma 19, $\mathcal{Z}_{\mathfrak{g}}$ is equidimensional of dimension $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g}) + 1$. Let \mathcal{Z} be an irreducible component of $\mathcal{Z}_{\mathfrak{g}}$. By Proposition 36(i), $\varpi_1(\mathcal{Z})$ is equal to $\mathfrak{X}_{\mathfrak{g}}$. Hence, by (i) and Proposition 37(v), the nullvariety of $p_{1,2,0}$ in \mathcal{Z} is a union of irreducible components of $\mathcal{N}_{\mathfrak{g}}$ of maximal dimension. Then, by Proposition 36(ii), $\varpi_2(\mathcal{Z})$ is equal to $\mathfrak{N}_{\mathfrak{g}}$ since any irreducible component of $\mathcal{N}_{\mathfrak{g}}$ is stable under the involution $(x, y) \mapsto (y, x)$. So, by Lemma 22, \mathcal{Z} has a nonempty intersection with $\Omega_{\mathfrak{g}}$. \square

We can now give the proof of Theorem 33.

Proof. (i) By Corollary 38(i), Lemma 18(i) and Lemma 20(ii), $\mathcal{X}_{\mathfrak{g}}$ is a complete intersection of dimension $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g} + 1)$ and any irreducible component of $\mathcal{X}_{\mathfrak{g}}$ contains a smooth point. So $\mathcal{X}_{\mathfrak{g}}$ is generically reduced and, by [Ma, Chap. 8, §23], the scheme $\mathcal{X}_{\mathfrak{g}}$ is reduced.

(ii) By Corollary 38(ii), Lemma 19(ii) and Lemma 20(ii), $\mathcal{Y}_{\mathfrak{g}}$ is a complete intersection of dimension $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g}) + 2$ and any irreducible component of $\mathcal{Y}_{\mathfrak{g}}$ contains a smooth point. So $\mathcal{Y}_{\mathfrak{g}}$ is generically reduced and, by [Ma, Chap. 8, §23], the scheme $\mathcal{Y}_{\mathfrak{g}}$ is reduced. Moreover, by Proposition 37(ii), any irreducible component of $\mathcal{Y}_{\mathfrak{g}}$ is the intersection of $\mathcal{Y}_{\mathfrak{g}}$ and an irreducible component of $\mathcal{X}_{\mathfrak{g}}$.

(iii) By Corollary 38(iii), Lemma 18(i) and Lemma 20(ii), $\mathcal{Z}_{\mathfrak{g}}$ is a complete intersection of dimension $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g}) + 1$ and any irreducible component of $\mathcal{Z}_{\mathfrak{g}}$ contains a smooth point. So $\mathcal{Z}_{\mathfrak{g}}$ is generically reduced and by [Ma, Chap. 8, §23], the scheme $\mathcal{Z}_{\mathfrak{g}}$ is reduced.

(iv) As $\mathcal{N}_{\mathfrak{g}}$ is the nullvariety of $\mathfrak{b}_{\mathfrak{g}} + \text{rk } \mathfrak{g}$ polynomial functions, any irreducible component of $\mathcal{N}_{\mathfrak{g}}$ has dimension at least $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g})$. Hence, by Corollary 38(i) and Proposition 37(v), $\mathcal{N}_{\mathfrak{g}}$ is a complete intersection of dimension $3(\mathfrak{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g})$. \square

We deduce Theorem 2 from Theorem 33(i) and Proposition 36(i) and we deduce the main part of Theorem 1 from Theorem 33(iv) and Proposition 36(ii). To complete the proof of Theorem 1, it only remains to show that $\mathcal{N}_{\mathfrak{g}}$ is a nonreduced scheme, which we will do in Section 6.

5. Applications to invariant theory

In this section we present various applications of previous results to invariant theory. In this section, \mathfrak{g} is supposed to be simple again.

5.1. Properties of the algebra of $\text{pol}_2\mathbf{S}(\mathfrak{g})^{\mathfrak{g}}$

We denote by $\text{pol}_2\mathbf{S}(\mathfrak{g})^{\mathfrak{g}}$ the subalgebra of $\mathbf{S}(\mathfrak{g}) \otimes_{\mathbb{C}} \mathbf{S}(\mathfrak{g})$ generated by the 2-order polarizations of elements of $\mathbf{S}(\mathfrak{g})^{\mathfrak{g}}$. Since the polynomials $p_1, \dots, p_{\text{rk } \mathfrak{g}}$ generate $\mathbf{S}(\mathfrak{g})^{\mathfrak{g}}$, the polynomials $p_{i,m,n}$, where $i = 1, \dots, \text{rk } \mathfrak{g}$, $m+n = d_i$ generate $\text{pol}_2\mathbf{S}(\mathfrak{g})^{\mathfrak{g}}$ as a graded algebra. Besides, the morphism σ , introduced in Subsection 1.4, is the morphism of affine varieties whose comorphism is the canonical injection from $\text{pol}_2\mathbf{S}(\mathfrak{g})^{\mathfrak{g}}$ into $\mathbf{S}(\mathfrak{g}) \otimes_{\mathbb{C}} \mathbf{S}(\mathfrak{g})$. Then, by Proposition 13, we can state:

Proposition 39. *The subalgebra $\text{pol}_2\mathbf{S}(\mathfrak{g})^{\mathfrak{g}}$ is a polynomial algebra in $\text{b}_{\mathfrak{g}} + \text{rk } \mathfrak{g}$ variables.*

We deduce from Proposition 39:

Theorem 40. *The morphism σ is faithfully flat. Equivalently, the extension $\mathbf{S}(\mathfrak{g}) \otimes_{\mathbb{C}} \mathbf{S}(\mathfrak{g})$ of $\text{pol}_2\mathbf{S}(\mathfrak{g})^{\mathfrak{g}}$ is faithfully flat.*

Proof. As $\text{pol}_2\mathbf{S}(\mathfrak{g})^{\mathfrak{g}}$ is generated by homogeneous functions, the fibre at 0 of the morphism σ has maximal dimension. On the other hand, by Proposition 39, $\text{pol}_2\mathbf{S}(\mathfrak{g})^{\mathfrak{g}}$ is a polynomial algebra in $\text{b}_{\mathfrak{g}} + \text{rk } \mathfrak{g}$ variables. So σ is an equidimensional morphism and, by [Ma, Chap. 8, Theorem 21.3], σ is a flat morphism. In particular, by [Ha, (Chap. III, Exer. 9.4)], it is an open morphism whose image contains 0. So σ is surjective. Hence σ is faithfully flat, according to [Ma, Chap. 3, Theorem 7.2]. \square

5.2. The nullcone

The nullcone $\mathcal{V}_{\mathfrak{g}}$ of the G -module $\mathfrak{g} \times \mathfrak{g}$ is the nullvariety in $\mathfrak{g} \times \mathfrak{g}$ of the augmentation ideal of the ideal of $\mathbf{S}(\mathfrak{g}) \otimes_{\mathbb{C}} \mathbf{S}(\mathfrak{g})$ generated by $(\mathbf{S}(\mathfrak{g}) \otimes_{\mathbb{C}} \mathbf{S}(\mathfrak{g}))^{\mathfrak{g}}$. The nullcone plays a leading part in invariant theory. It is studied in [Ri5], [Po], [LMP], and recently in [KrW1] and [KrW2]. By [KrW1], $\mathcal{V}_{\mathfrak{g}}$ is irreducible and has dimension $3(\text{b}_{\mathfrak{g}} - \text{rk } \mathfrak{g})$. Furthermore, Kraft and Wallach conjecture in [KrW2] that $\mathcal{V}_{\mathfrak{g}}$ is an irreducible component of $\mathcal{N}_{\mathfrak{g}}$. Clearly, Theorem 1 confirms that conjecture. Thereby, we claim:

Theorem 41. *The nullcone $\mathcal{V}_{\mathfrak{g}}$ is an irreducible component of the nilpotent bicone $\mathcal{N}_{\mathfrak{g}}$.*

Let k be in \mathbb{N}^* and let us denote by \mathfrak{g}^k the k th Cartesian power of \mathfrak{g} . We extend the previous notions to \mathfrak{g}^k . Let $\mathbf{S}(\mathfrak{g}^k)$ be the symmetric algebra of \mathfrak{g}^k and let $\mathbf{S}(\mathfrak{g}^k)^{\mathfrak{g}}$ be the subalgebra of its invariant elements under the diagonal action of G in \mathfrak{g}^k . Then we denote by $\mathcal{V}_{\mathfrak{g}}^{(k)}$ the nullvariety in \mathfrak{g}^k of the augmentation ideal $\mathbf{S}(\mathfrak{g}^k)_{+}^{\mathfrak{g}}$ of $\mathbf{S}(\mathfrak{g}^k)^{\mathfrak{g}}$. For φ in $\mathbf{S}(\mathfrak{g})$, the k -order polarizations $\varphi_{i_1, \dots, i_k}$ of φ are defined by the relation:

$$\varphi(t_1x_1 + \dots + t_kx_k) = \sum_{(i_1, \dots, i_k) \in \mathbb{N}^k} t_1^{i_1} \dots t_k^{i_k} \varphi_{i_1, \dots, i_k}(x_1, \dots, x_k),$$

for (x_1, \dots, x_k) in \mathfrak{g}^k and (t_1, \dots, t_k) in \mathbb{C}^k . Then, we denote by $\text{pol}_k\mathbf{S}(\mathfrak{g})^{\mathfrak{g}}$ the subalgebra of $\mathbf{S}(\mathfrak{g}^k)$ generated by the k -order polarizations of elements in $\mathbf{S}(\mathfrak{g})^{\mathfrak{g}}$. Then $\text{pol}_k\mathbf{S}(\mathfrak{g})^{\mathfrak{g}}$ is a homogeneous subalgebra of $\mathbf{S}(\mathfrak{g}^k)$ and its augmentation ideal $\text{pol}_k\mathbf{S}(\mathfrak{g})_{+}^{\mathfrak{g}}$ is generated by the k -order polarizations of the elements of $\mathbf{S}(\mathfrak{g})_{+}^{\mathfrak{g}}$. We

NILPOTENT BICONE

denote by $\mathcal{N}_{\mathfrak{g}}^{(k)}$ the nullvariety in \mathfrak{g}^k of the ideal $\text{pol}_k \mathcal{S}(\mathfrak{g})_{+}^{\mathfrak{g}}$. Obviously, $\mathcal{V}_{\mathfrak{g}}^{(1)} = \mathcal{N}_{\mathfrak{g}}^{(1)} = \mathfrak{N}_{\mathfrak{g}}$, $\mathcal{V}_{\mathfrak{g}}^{(2)} = \mathcal{V}_{\mathfrak{g}}$ and $\mathcal{N}_{\mathfrak{g}}^{(2)} = \mathcal{N}_{\mathfrak{g}}$.

The *index of polarization of \mathfrak{g}* , denoted by $\text{pol ind}(\mathfrak{g})$, is defined in [LMP] as being the upper bound of $k \in \mathbb{N}^*$ such that $\mathcal{V}_{\mathfrak{g}}^{(k)} = \mathcal{N}_{\mathfrak{g}}^{(k)}$. The inequality $\text{pol ind}(\mathfrak{g}) \geq 1$ obviously always holds. We next wish to establish an equality.

Recall that (e, h, f) is a principal \mathfrak{sl}_2 -triple of \mathfrak{g} .

Lemma 42. *Let x be in \mathfrak{g} satisfying $(\text{ad } x)^2(e) = 0$. Then $(e, [x, e])$ belongs to $\mathcal{N}_{\mathfrak{g}}$.*

Proof. Since $(\text{ad } x)^2(e) = 0$, we have $\exp(\text{tad } x)(e) = e + t[x, e]$, for any t in \mathbb{C} . As $\mathfrak{N}_{\mathfrak{g}}$ is a closed cone in \mathfrak{g} , the element $se + t[x, e]$ belongs to $\mathfrak{N}_{\mathfrak{g}}$ for any (s, t) in \mathbb{C}^2 . So $(e, [x, e])$ belongs to $\mathcal{N}_{\mathfrak{g}}$. \square

By the Hilbert–Mumford criterion [Kr](Kap. II), an element (x, y) of $\mathfrak{g} \times \mathfrak{g}$ belongs to the nullcone $\mathcal{V}_{\mathfrak{g}}$ if and only if there is a one-parameter subgroup $\lambda : \mathbb{C}^* \rightarrow G$ such that $\lim_{t \rightarrow 0} \lambda(t)(x, y) = 0$. In particular, if (x, y) belongs to the nullcone, then the subalgebra L generated by x and y is contained in the nilpotent cone. Hence L is nilpotent and so contained in a Borel subalgebra of \mathfrak{g} . Conversely, if x and y belong to the nilpotent radical of a common Borel subalgebra, then the previous criterion applies to (x, y) . To summarize, we have actually shown:

Lemma 43. *The element (x, y) of $\mathfrak{g} \times \mathfrak{g}$ belongs to the nullcone $\mathcal{V}_{\mathfrak{g}}$ if and only if x and y belong to the nilpotent radical of a common Borel subalgebra.*

The following proposition has already been noticed in [LMP, Theorem 3.16]. We provide here a shorter proof.

Proposition 44. *We have*

$$\text{pol ind}(\mathfrak{g}) = \begin{cases} \infty & \text{if } \mathfrak{g} \text{ is isomorphic to } \mathfrak{sl}_2(\mathbb{C}), \\ 1 & \text{otherwise.} \end{cases}$$

Proof. Suppose first $\dim \mathfrak{g} = 3$. The elements x in \mathfrak{g} such that the subspace generated by x and e is contained in the nilpotent cone of \mathfrak{g} are collinear to e . Hence $\mathcal{V}_{\mathfrak{g}}^{(2)} = \mathcal{N}_{\mathfrak{g}}$, and $\mathcal{N}_{\mathfrak{g}}$ is irreducible. Then for $k \geq 2$ and x_1, \dots, x_k in \mathfrak{g} , the subspace generated by x_1, \dots, x_k is contained in $\mathfrak{N}_{\mathfrak{g}}$ if and only if its dimension is smaller than 1. So $\mathcal{V}_{\mathfrak{g}}^{(k)}$ is equal to $\mathcal{N}_{\mathfrak{g}}^{(k)}$ for any k .

Suppose now $\dim \mathfrak{g} > 3$. As $d_1, \dots, d_{\text{rk } \mathfrak{g}}$ is a weakly increasing sequence, $-2(d_{\text{rk } \mathfrak{g}} - 1)$ is the smallest eigenvalue of $\text{ad } h$. Let v be a nonzero eigenvector of $\text{ad } h$ of eigenvalue $-2(d_{\text{rk } \mathfrak{g}} - 1)$ and set $v_0 = [v, e]$. Then $(\text{ad } v)^2(e)$ is an eigenvector of $\text{ad } h$ of eigenvalue $-4(d_{\text{rk } \mathfrak{g}} - 1) + 2$. As \mathfrak{g} is simple and $\dim \mathfrak{g} > 3$, we have $d_{\text{rk } \mathfrak{g}} > 2$ and $-4(d_{\text{rk } \mathfrak{g}} - 1) + 2 < -2(d_{\text{rk } \mathfrak{g}} - 1)$. So $(\text{ad } v)^2(e)$ is equal to 0. Hence, by Lemma 42, (e, v_0) belongs to $\mathcal{N}_{\mathfrak{g}}$. As v_0 is an eigenvector of $\text{ad } h$ of negative eigenvalue, \mathfrak{b} does not contain v_0 . Since \mathfrak{b} is the unique Borel subalgebra which contains e , we deduce from Lemma 43 that (e, v_0) does not belong to $\mathcal{V}_{\mathfrak{g}}$. So $\mathcal{V}_{\mathfrak{g}}$ is strictly contained in $\mathcal{N}_{\mathfrak{g}}$. As a result, for $k \geq 2$, $\mathcal{V}_{\mathfrak{g}}^{(k)}$ is strictly contained in $\mathcal{N}_{\mathfrak{g}}^{(k)}$ since $\mathcal{V}_{\mathfrak{g}} \times \{0_{\mathfrak{g}^{k-2}}\}$ and $\mathcal{N}_{\mathfrak{g}} \times \{0_{\mathfrak{g}^{k-2}}\}$ are the intersections of $\mathcal{V}_{\mathfrak{g}}^{(k)}$ and $\mathcal{N}_{\mathfrak{g}}^{(k)}$ with $\mathfrak{g} \times \mathfrak{g} \times \{0_{\mathfrak{g}^{k-2}}\}$ respectively. \square

Remark 8. The nullcone has a natural structure of scheme. This scheme is irreducible. Furthermore, it is reduced if and only if \mathfrak{g} has dimension 3. If $\dim \mathfrak{g} = 3$, $\text{pol}_2 \mathbb{S}(\mathfrak{g})^{\mathfrak{g}}$ is equal to $(\mathbb{S}(\mathfrak{g}) \otimes_{\mathbb{C}} \mathbb{S}(\mathfrak{g}))^{\mathfrak{g}}$ and the ideal in $\mathbb{S}(\mathfrak{g}) \otimes_{\mathbb{C}} \mathbb{S}(\mathfrak{g})$ generated by the augmentation ideal in $\mathbb{S}(\mathfrak{g})^{\mathfrak{g}}$ is prime. Suppose $\dim \mathfrak{g} > 3$. If this scheme were reduced, the extension $\mathbb{S}(\mathfrak{g}) \otimes_{\mathbb{C}} \mathbb{S}(\mathfrak{g})$ of $(\mathbb{S}(\mathfrak{g}) \otimes_{\mathbb{C}} \mathbb{S}(\mathfrak{g}))^{\mathfrak{g}}$ would be flat by arguments used in [Kos2]. But this extension is not equidimensional since the G -orbit in $\mathfrak{g} \times \mathfrak{g}$ in general position has dimension $2b_{\mathfrak{g}} - \text{rk } \mathfrak{g}$ and $2b_{\mathfrak{g}} - \text{rk } \mathfrak{g} > 3(b_{\mathfrak{g}} - \text{rk } \mathfrak{g})$.

6. Additional properties of the nilpotent bicone

In this section we give additional results about the nilpotent bicone $\mathcal{N}_{\mathfrak{g}}$. We assert in the first subsection that $\mathcal{N}_{\mathfrak{g}}$ is not reduced. Next, we discuss in the second subsection the number of its irreducible components. As before, we fix a principal \mathfrak{sl}_2 -triple (e, h, f) in \mathfrak{g} and we use the notations of the Introduction, Subsection 0.3.

6.1. Nonreducibility of $\mathcal{N}_{\mathfrak{g}}$

According to [Ma, §23], a Noetherian ring A is reduced if and only if A satisfies conditions (R_0) and (S_1) . By Theorem 33, the subscheme $\mathcal{N}_{\mathfrak{g}}$ of $\mathfrak{g} \times \mathfrak{g}$ defined by the ideal $\text{pol}_2 \mathbb{S}(\mathfrak{g})_+^{\mathfrak{g}}$ is a complete intersection in $\mathfrak{g} \times \mathfrak{g}$. Therefore the quotient ring $(\mathbb{S}(\mathfrak{g}) \otimes_{\mathbb{C}} \mathbb{S}(\mathfrak{g})) / \text{pol}_2 \mathbb{S}(\mathfrak{g})_+^{\mathfrak{g}}$ is Cohen–Macaulay, according to [Ma, (Chap. 8, Theorem 21.3)]. It is natural to ask if $(\mathbb{S}(\mathfrak{g}) \otimes_{\mathbb{C}} \mathbb{S}(\mathfrak{g})) / \text{pol}_2 \mathbb{S}(\mathfrak{g})_+^{\mathfrak{g}}$ is reduced. The following theorem answers this question negatively.

Theorem 45. *The scheme $\mathcal{N}_{\mathfrak{g}}$ is not reduced.*

Proof. In view of the preceding remarks, it suffices to find an irreducible component of $\mathcal{N}_{\mathfrak{g}}$ which does not contain any smooth point of $\mathcal{N}_{\mathfrak{g}}$. Hence, according to Proposition 13, Lemma 20(ii) and Theorem 41, it is enough to prove that the nullcone $\mathcal{V}_{\mathfrak{g}}$ has an empty intersection with the open subset $\Omega_{\mathfrak{g}}$. We suppose that this intersection is not empty and we expect a contradiction. Let (x, y) be in the intersection of $\mathcal{V}_{\mathfrak{g}}$ and $\Omega_{\mathfrak{g}}$. By Lemma 43, we can suppose that x and y belong to \mathfrak{u} . By Lemma 7(ii), for any (a, b) in $\mathbb{C}^2 \setminus \{(0, 0)\}$, the centralizer of $ax + by$ is contained in \mathfrak{u} . Therefore, it results from Lemma 4 that the subspace $\mathfrak{V}'(x, y)$ is contained in \mathfrak{u} . Moreover, by Lemma 8(iii), $\mathfrak{V}'(x, y)$ is equal to $\mathfrak{V}(x, y)$. But by Lemma 8(i), $\mathfrak{V}(x, y)$ has dimension $b_{\mathfrak{g}} > \dim \mathfrak{u}$, since (x, y) belongs to $\Omega_{\mathfrak{g}}$, whence the expected contradiction. \square

The proof of Theorem 1 is now completed by Theorem 45.

6.2. Irreducible components of $\mathcal{N}_{\mathfrak{g}}$

Denote by Φ the map

$$G \times \mathfrak{g} \times \mathfrak{g} \longrightarrow \mathfrak{g} \times \mathfrak{g}, \quad (g, x, y) \mapsto (g(x), g(y)).$$

For a subset X of $\mathfrak{g} \times \mathfrak{g}$, the closure in $\mathfrak{g} \times \mathfrak{g}$ of the image of $G \times X$ by Φ is denoted by X^{Φ} . For Y a subset of $\mathfrak{g} \times \mathfrak{g}$ and x in \mathfrak{g} , we denote by Y_x the subset of elements y in \mathfrak{g} such that (x, y) belongs to Y .

NILPOTENT BICONE

Lemma 46. *Let Y be a G -invariant closed bicone of $\mathfrak{g} \times \mathfrak{g}$ and let x_0 be a nilpotent element of \mathfrak{g} . Suppose that the image of any irreducible component of Y by ϖ_1 is equal to the closure in \mathfrak{g} of $G.x_0$. Suppose besides that there exists a subvariety T of G such that the map $g \mapsto g(x_0)$ is an isomorphism from T to an open subset of $G.x_0$ containing x_0 .*

- (i) *The subset Y is equal to $(\{x_0\} \times Y_{x_0})^\Phi$.*
- (ii) *If Z is an irreducible component of Y , then Z_{x_0} is irreducible.*
- (iii) *The map $X \mapsto (\{x_0\} \times X)^\Phi$ is a bijection from the set of irreducible components of Y_{x_0} to the set of irreducible components of Y .*

Proof. (i) By our hypothesis, for any irreducible component Z of Y , $\varpi_1(Z)$ is the closure of $G.x_0$. So $G.x_0$ is a dense open subset in $\varpi_1(Z)$. Then the subset Y' of elements of Y whose first component is in $G.x_0$ is a dense open subset in Y . As Y is a G -invariant bicone, Y_{x_0} is a cone and Y' is the image by Φ of $G \times \{x_0\} \times Y_{x_0}$. So Y is equal to $(\{x_0\} \times Y_{x_0})^\Phi$.

(ii) Let Z be an irreducible component of Y . Then Z is a G -invariant closed bicone. We denote by Ω the image of T by the map $g \mapsto g(x_0)$ and we denote by \tilde{Z} the subset of elements (x, y) in Z such that x belongs to Ω . By our hypothesis, \tilde{Z} is a nonempty open subset of Z . In particular, it is irreducible. Let τ be the inverse map of the map $g \mapsto g(x_0)$ from T to Ω . Then the map

$$\tilde{Z} \longrightarrow T \times Z_{x_0}, \quad (x, y) \mapsto (\tau(x), \tau(x)^{-1}(y)),$$

is an isomorphism. So Z_{x_0} is irreducible.

(iii) Let X be an irreducible component of Y_{x_0} . Then $(\{x_0\} \times X)^\Phi$ is irreducible as the closure of the image of an irreducible variety by a regular map. If Z is an irreducible component of Y which contains $(\{x_0\} \times X)^\Phi$ then Z_{x_0} contains X . So, by (ii), X is equal to Z_{x_0} and Z is equal to $(\{x_0\} \times X)^\Phi$ since $\Phi(G \times \{x_0\} \times X)$ is open in Z . We then deduce that the map $Z \mapsto Z_{x_0}$ is the inverse of the map $X \mapsto (\{x_0\} \times X)^\Phi$. \square

Let us recall that \mathbf{B}_- is the normalizer in G of the Borel subalgebra $\mathfrak{b}_- = \mathfrak{h} \oplus \mathfrak{u}_-$.

Lemma 47. *Let \mathbf{U} be the unipotent radical of the normalizer of \mathfrak{b} in G . Then there exists a closed subset Z of \mathbf{U} such that the map*

$$\mathbf{B}_- \times Z \rightarrow G.e, \quad (g, k) \mapsto gk(e),$$

is an isomorphism onto an open subset V of $G.e$ containing e .

Proof. As $\mathbf{B}_-\mathbf{U}$ is an open subset of G , its image by the map $g \mapsto g(e)$ from G to $G.e$ is an open subset V containing e . Moreover, the map

$$\mathbf{B}_- \times \mathbf{U}.e \rightarrow V, \quad (g, x) \mapsto g(x),$$

is an isomorphism since \mathbf{U} contains the stabilizer of e in G and $\mathbf{B}_- \cap \mathbf{U} = \{\mathbf{1}_{\mathfrak{g}}\}$. By [Pu, Chap. I, Part II, §3], there exists a complement \mathfrak{m} of $\mathfrak{g}(e)$ in \mathfrak{u} and a basis v_1, \dots, v_m of \mathfrak{m} such that the map

$$\mathbb{C}^m \rightarrow \mathbf{U}.e, \quad (t_1, \dots, t_m) \mapsto \exp(t_1 \operatorname{ad} v_1) \cdots \exp(t_m \operatorname{ad} v_m)(e),$$

is an isomorphism. Then the map

$$\mathbb{C}^m \rightarrow \mathbf{U}, \quad (t_1, \dots, t_m) \mapsto \exp(t_1 \operatorname{ad} v_1) \cdots \exp(t_m \operatorname{ad} v_m),$$

is proper and its image Z is closed in \mathbf{U} . Moreover, the map

$$\mathbf{B}_- \times Z \rightarrow V, \quad (g, k) \mapsto gk(e),$$

is an isomorphism. \square

Let $\mathcal{N}_{\mathfrak{g},e}$ be the subset of elements y of \mathfrak{g} such that (e, y) is in $\mathcal{N}_{\mathfrak{g}}$. From Theorem 1, Lemma 46(iii) and Lemma 47, we deduce the following result.

Corollary 48. *The map $X \mapsto (\{e\} \times X)^\Phi$ is a bijection from the set of irreducible components of $\mathcal{N}_{\mathfrak{g},e}$ onto the set of irreducible components of $\mathcal{N}_{\mathfrak{g}}$.*

According to Subsection 2.3, $\mathcal{N}_{\mathfrak{g},e}$ has a natural structure of scheme. The following result is also a corollary of Theorem 1.

Corollary 49. *The subset $\mathcal{N}_{\mathfrak{g},e}$ is a complete intersection of dimension $\mathfrak{b}_{\mathfrak{g}} - \operatorname{rk} \mathfrak{g}$.*

Proof. As $\mathcal{N}_{\mathfrak{g},e}$ is the nullvariety in \mathfrak{g} of $\mathfrak{b}_{\mathfrak{g}}$ polynomial functions, the dimension of any irreducible component of $\mathcal{N}_{\mathfrak{g},e}$ is at least $\mathfrak{b}_{\mathfrak{g}} - \operatorname{rk} \mathfrak{g}$. Let Y be an irreducible component of $\mathcal{N}_{\mathfrak{g}}$. By Theorem 1, since $G.e$ is an open dense subset of $\mathfrak{N}_{\mathfrak{g}}$, Y_e has dimension $\mathfrak{b}_{\mathfrak{g}} - \operatorname{rk} \mathfrak{g}$. On the other hand, by Corollary 48, Y_e is an irreducible component of $\mathcal{N}_{\mathfrak{g},e}$ and any irreducible component of $\mathcal{N}_{\mathfrak{g},e}$ is obtained in this way, whence the corollary. \square

Let $\mathfrak{p} = \mathfrak{l}_{\mathfrak{p}} \oplus \mathfrak{u}_{\mathfrak{p}}$ be a parabolic subalgebra of \mathfrak{g} containing \mathfrak{b} , where $\mathfrak{l}_{\mathfrak{p}}$ is the \mathfrak{h} -stable Levi subalgebra of \mathfrak{p} and where $\mathfrak{u}_{\mathfrak{p}}$ is the nilpotent radical of \mathfrak{p} . We denote by $\varpi_{\mathfrak{l}_{\mathfrak{p}}}$ and $\varpi_{\mathfrak{u}_{\mathfrak{p}}}$ the projections from \mathfrak{p} to $\mathfrak{l}_{\mathfrak{p}}$ and $\mathfrak{u}_{\mathfrak{p}}$ respectively.

Lemma 50. *The map $Y \mapsto Y + \mathfrak{u}_{\mathfrak{p}}$ is a one-to-one correspondence between the set of irreducible components of $\mathcal{N}_{\mathfrak{l}_{\mathfrak{p}}, \varpi_{\mathfrak{l}_{\mathfrak{p}}}(e)}$ and the set of irreducible components of $\mathcal{N}_{\mathfrak{g},e}$ contained in \mathfrak{p} .*

Proof. Let Y be an irreducible component of $\mathcal{N}_{\mathfrak{l}_{\mathfrak{p}}, \varpi_{\mathfrak{l}_{\mathfrak{p}}}(e)}$. Then the set $Y + \mathfrak{u}_{\mathfrak{p}}$ is contained in the intersection of $\mathcal{N}_{\mathfrak{g},e}$ and \mathfrak{p} . As e is a regular nilpotent element of \mathfrak{g} , $\varpi_{\mathfrak{l}_{\mathfrak{p}}}(e)$ is a regular nilpotent element of $\mathfrak{l}_{\mathfrak{p}}$. So by Corollary 49, Y has dimension $\mathfrak{b}_{\mathfrak{l}_{\mathfrak{p}}} - \operatorname{rk} \mathfrak{g}$. Then $Y + \mathfrak{u}_{\mathfrak{p}}$ has dimension

$$\mathfrak{b}_{\mathfrak{l}_{\mathfrak{p}}} - \operatorname{rk} \mathfrak{g} + \dim \mathfrak{u}_{\mathfrak{p}} = \mathfrak{b}_{\mathfrak{g}} - \operatorname{rk} \mathfrak{g}.$$

According to Corollary 49, we deduce that $Y + \mathfrak{u}_{\mathfrak{p}}$ is an irreducible component of $\mathcal{N}_{\mathfrak{g},e}$. Thus, the above map is well defined. It is clearly injective. Prove now that it is surjective.

Let Z be an irreducible component of $\mathcal{N}_{\mathfrak{g},e}$ contained in \mathfrak{p} . Then $\varpi_{\mathfrak{l}_{\mathfrak{p}}}(Z)$ is contained in an irreducible component of $\mathcal{N}_{\mathfrak{l}_{\mathfrak{p}}, \varpi_{\mathfrak{l}_{\mathfrak{p}}}(e)}$. Let Y be such an irreducible component. Then $Y + \mathfrak{u}_{\mathfrak{p}}$ is an irreducible subset of $\mathcal{N}_{\mathfrak{g},e}$ which contains Z . So, Z is equal to $Y + \mathfrak{u}_{\mathfrak{p}}$. \square

Let $N_{\mathfrak{g}}$ be the set of irreducible components of $\mathcal{N}_{\mathfrak{g},e}$. We denote by $\Upsilon_{\mathfrak{g}}$ the set of proper parabolic subalgebras strictly containing \mathfrak{b} . Let $N'_{\mathfrak{g}}$ be the subset of irreducible components of $\mathcal{N}_{\mathfrak{g},e}$ which are not contained in any element of $\Upsilon_{\mathfrak{g}}$.

NILPOTENT BICONE

Lemma 51. *If $\dim \mathfrak{g} = 3$, then $|N'_{\mathfrak{g}}| = 0$. Otherwise, $|N'_{\mathfrak{g}}| \geq 1$.*

Proof. The set $\Upsilon_{\mathfrak{g}}$ is nonempty if and only if $\dim \mathfrak{g} > 3$. Suppose now $\dim \mathfrak{g} > 3$. Let $\alpha^{\#}$ be the highest root of \mathcal{R}_+ . By Lemma 42, $\mathcal{N}_{\mathfrak{g},e}$ contains $[e, \mathfrak{g}^{-\alpha^{\#}}]$. So it suffices to prove that $[e, \mathfrak{g}^{-\alpha^{\#}}]$ is not contained in \mathfrak{p} for any \mathfrak{p} in $\Upsilon_{\mathfrak{g}}$. For α in \mathcal{R} , we fix a nonzero element x_{α} in \mathfrak{g}^{α} . The elements x_{α} may be chosen so that $e = \sum_{\alpha \in \Pi} x_{\alpha}$. Suppose that there is \mathfrak{p} in $\Upsilon_{\mathfrak{g}}$ such that $[x_{-\alpha^{\#}}, e]$ belongs to \mathfrak{p} . We can suppose that \mathfrak{p} is a maximal element of $\Upsilon_{\mathfrak{g}}$ corresponding to a simple root β . A short discussion shows that β satisfies the following two conditions:

- (1) for all α in $\Pi \setminus \{\beta\}$, $\alpha^{\#} - \alpha$ is not a root; and
- (2) $\alpha^{\#} - \beta$ is a root and $n_{\beta} = 1$, where n_{β} is the coordinate at β of $\alpha^{\#}$ in the basis Π .

A quick look at the classification of root systems shows that there is no simple root satisfying both conditions (1) and (2). As we obtain a contradiction, the result follows. \square

Proposition 52. *The number of irreducible components of $\mathcal{N}_{\mathfrak{g}}$ is equal to $|N_{\mathfrak{g}}|$ and $|N_{\mathfrak{g}}|$ satisfies the following recursive relation:*

$$|N_{\mathfrak{g}}| = |N'_{\mathfrak{g}}| + \left(\sum_{\mathfrak{p} \in \Upsilon_{\mathfrak{g}}} \prod_{\substack{\text{simple factor} \\ \text{of } [\mathfrak{t}_{\mathfrak{p}}, \mathfrak{t}_{\mathfrak{p}}]}} |N'_{\mathfrak{t}}| \right) + 1.$$

Proof. By Corollary 48, the number of irreducible components of $\mathcal{N}_{\mathfrak{g}}$ is $|N_{\mathfrak{g}}|$. If $\dim \mathfrak{g} = 3$, then $\mathcal{N}_{\mathfrak{g},e}$ is the line generated by e , $\Upsilon_{\mathfrak{g}} = \emptyset$, $|N'_{\mathfrak{g}}| = 0$ and $|N_{\mathfrak{g}}| = 1$, whence the expected relation. Suppose now $\dim \mathfrak{g} > 3$. We can write

$$N_{\mathfrak{g}} = N'_{\mathfrak{g}} \cup \left(\bigcup_{\mathfrak{p} \in \Upsilon_{\mathfrak{g}}} N_{\mathfrak{g}}^{(\mathfrak{p})} \right) \cup \{\mathfrak{u}\},$$

where, for \mathfrak{p} in $\Upsilon_{\mathfrak{g}}$, $N_{\mathfrak{g}}^{(\mathfrak{p})}$ is the set of irreducible components of $\mathcal{N}_{\mathfrak{g},e}$ contained in \mathfrak{p} and not contained in any element of $\Upsilon_{\mathfrak{g}}$ strictly contained in \mathfrak{p} . Thus the terms of the above union are pairwise disjoint and we have

$$|N_{\mathfrak{g}}| = |N'_{\mathfrak{g}}| + \sum_{\mathfrak{p} \in \Upsilon_{\mathfrak{g}}} |N_{\mathfrak{g}}^{(\mathfrak{p})}| + 1.$$

It remains to prove the equality

$$|N_{\mathfrak{g}}^{(\mathfrak{p})}| = \prod_{\substack{\text{simple factor} \\ \text{of } [\mathfrak{t}_{\mathfrak{p}}, \mathfrak{t}_{\mathfrak{p}}]}} |N'_{\mathfrak{t}}|, \tag{12}$$

for any \mathfrak{p} in $\Upsilon_{\mathfrak{g}}$. Let \mathfrak{p} be in $\Upsilon_{\mathfrak{g}}$. By Lemma 50, the map $Y \mapsto Y + \mathfrak{u}_{\mathfrak{p}}$ is a bijection from $N_{\mathfrak{t}_{\mathfrak{p}}}$ to the subset of irreducible components of $\mathcal{N}_{\mathfrak{g},e}$ contained in \mathfrak{p} . Moreover, $Y + \mathfrak{u}_{\mathfrak{p}}$ belongs to $N_{\mathfrak{g}}^{(\mathfrak{p})}$ if and only if Y belongs to $N'_{\mathfrak{t}_{\mathfrak{p}}}$. So $|N'_{\mathfrak{t}_{\mathfrak{p}}}| = |N_{\mathfrak{g}}^{(\mathfrak{p})}|$.

Let $\mathfrak{l}_1, \dots, \mathfrak{l}_m$ be the simple factors of \mathfrak{l}_p and let Y be in $N_{\mathfrak{l}_p}$. As remarked in Subsection 0.3, $Y = Y_1 \times \dots \times Y_m$ where Y_i is an irreducible component of $\mathcal{N}_{\mathfrak{l}_i, e_i}$, if e_i is the component of $\varpi_{\mathfrak{l}_p}(e)$ on \mathfrak{l}_i for $i = 1, \dots, m$. Then, $Y \in N'_{\mathfrak{l}_p}$ if and only if, for any $i \in \{1, \dots, m\}$, Y_i is not contained in any element of $\Upsilon_{\mathfrak{l}_i}$, that is, $Y_i \in N'_{\mathfrak{l}_p}$. Hence,

$$|N'_{\mathfrak{l}_p}| = \prod_{i=1}^m |N'_{\mathfrak{l}_i}|,$$

and the proposition follows, by relation (12). \square

Remark 9. Considering the formula established in Proposition 52, we see that the number $|N_{\mathfrak{g}}|$ becomes considerably bigger while the dimension of \mathfrak{g} grows up. For example, we have

$$|N_{\mathfrak{sl}_2}| = 1, \quad |N_{\mathfrak{sl}_3}| \geq 2, \quad |N_{\mathfrak{sl}_4}| \geq 4, \quad |N_{\mathfrak{sl}_5}| \geq 7, \quad |N_{\mathfrak{sl}_6}| \geq 12, \quad \dots$$

References

- [Ba] V. V. Batyrev, *Stringy Hodge numbers of varieties with Gorenstein canonical singularities*, in: *Integrable Systems and Algebraic Geometry* (Kobe/Kyoto), World Scientific, River Edge, NJ, 1998, pp. 1–32.
- [Bol] A. V. Bolsinov, *Commutative families of functions related to consistent Poisson brackets*, Acta Appl. Math. **24** (1991), 253–274.
- [Bou] N. Bourbaki, *Lie Groups and Lie Algebras*, Chaps. 4–6, Translated from the 1968 French original by Andrew Pressley, Springer-Verlag, Berlin, 2002.
- [Br] W. Bruns, J. Herzog, *Cohen–Macaulay Rings*, Cambridge Studies in Advanced Mathematics, Vol. 39, Cambridge University Press, Cambridge, 1996.
- [Ch] J.-Y. Charbonnel, *Complexe canonique de deuxième espèce, variété commutante et bicône nilpotent d'une algèbre de Lie réductive*, arXiv:math.RT/0509272.
- [Cr] A. Craw, *An introduction to motivic integration*, in: *Strings and Geometry*, Clay Math. Proc., Vol. 3, Amer. Math. Soc., Providence, RI, 2004, pp. 203–225.
- [DL1] J. Denef, F. Loeser, *Motivic Igusa zeta function*, J. Algebraic Geom. **7** (1998), no. 3, 505–537.
- [DL2] J. Denef, F. Loeser, *Germes of arcs on singular algebraic varieties and motivic integration*, Invent. Math. **135** (1999), 201–232.
- [DGT] D. Deturck, H. Goldschmidt, J. Talvacchia, *Connections with prescribed curvature and Yang–Mills currents: The semi-simple case*, Ann. Sci. 'Ecole Norm. Sup. **24** (1991), 57–112.
- [Di] J. Dixmier, *Champs de vecteurs adjoints sur les groupes et algèbres de Lie semi-simples*, J. Reine Angew. Math. **309** (1979), 183–190.
- [El1] R. Elkik, *Singularités rationnelles et déformations*, Invent. Math. **47** (1978), 139–147.
- [El2] R. Elkik, *Rationalité des singularités canoniques*, Invent. Math. **64** (1981), 1–6.
- [Fl] H. Flenner, *Rationale quasihomogene Singularitäten*, Arch. Math. **36** (1981), 35–44.

NILPOTENT BICONE

- [GG] W. L. Gan, V. Ginzburg, *Almost-commuting variety, \mathcal{D} -modules, and Cherednik algebras*, IMRP, Int. Math. Res. Pap. (2006), no. 2, 1–54.
- [Ha] R. Hartshorne, *Algebraic Geometry*, Graduate Texts in Mathematics, Vol. 52, Springer-Verlag, Berlin, 1977.
- [He] W. H. Hesselink, *Cohomology and the resolution of nilpotent variety*, Math. Ann. **223** (1976), 249–252.
- [Hi] H. Hironaka, *Resolution of singularities of an algebraic variety over a field of characteristic zero: I, II*, Ann. of Math. **79** (1964), 109–326.
- [Kon] M. Kontsevich, *Motivic integration*, Lecture at Orsay, 1995.
- [Kos1] B. Kostant, *The principal three-dimensional subgroup and the Betti numbers of a complex simple Lie Group*, Amer. J. Math. **81** (1959), 973–1032.
- [Kos2] B. Kostant, *Lie group representations on polynomial rings*, Amer. J. Math. **85** (1963), 327–404.
- [Kr] H. Kraft, *Geometrische Methoden in der Invariantentheorie*, Aspekte der Mathematik, Vol. D1, Vieweg Verlag, Braunschweig/Wiesbaden, 1985, (2., durchgesehene Auflage).
- [KrW1] H. Kraft, N. Wallach, *On the nullcone of representation of reductive groups*, Pacific J. Math. **224** (2006), 119–140.
- [KrW2] H. Kraft, N. Wallach, *Polarizations and nullcone of representations of reductive groups*, <http://math.ucsd.edu/~nwallach/KWpolarnullcone.pdf>, 2007.
- [LS] T. Lvasseur, J. T. Stafford, *The kernel of an homomorphism of Harish-Chandra*, Ann. Sci. École Norm. Sup. **29** (1996), 385–397.
- [Lo] E. Looijenga, *Motivic measures*, in: *Séminaire Bourbaki*, Astérisque **276** (1999/2000), 267–297.
- [LMP] M. Losik, P. W. Michor, V. L. Popov, *On polarizations in invariant theory*, J. Algebra **301** (2006), 406–424.
- [Ma] H. Matsumura, *Commutative Ring Theory*, Cambridge Studies in Advanced Mathematics, Vol. 8, Cambridge University Press, Cambridge, 1986).
- [Mu] M. Mustață, *Jet schemes of locally complete intersection canonical singularities*, with an appendix by D. Eisenbud and E. Frenkel, Invent. Math. **145** (2001), 397–424.
- [PYa] D. I. Panyushev, O. Yakimova, *The argument shift method and maximal commutative subalgebras of Poisson algebras*, Math. Res. Lett. **15** (2008), no. 2, 239–249.
- [Po] В. Л. Попов, *Конус нуль-форм Гильберта*, Труды Математического ин-та им. В. А. Стеклова **241** (2003), 192–209. English transl.: V. L. Popov, *The cone of Hilbert nullforms*, Proc. Steklov Inst. Math. **241** (2003), 177–194.
- [Pu] L. Pukanszky, *Leçons sur les Représentations des Groupes*, Monographie de la Société Mathématique de France, Vol. 2, Dunod, Paris, 1967.
- [Ri1] R. W. Richardson, *Commuting varieties of semisimple Lie algebras and algebraic groups*, Compos. Math. **38** (1979), 311–322.
- [Ri2] R. W. Richardson, *Normality of G -stable subvarieties of a semisimple Lie algebra*, in: *Algebraic Groups*, Proc. Symp., Utrecht/Neth. 1986, Lecture Notes in Mathematics, Vol. 1271, Springer-Verlag, Berlin, 1987, pp. 243–264.

- [Ri3] R. W. Richardson, *Derivatives of invariant polynomials on a semisimple Lie algebra*, in: *Microconference on Harmonic Analysis and Operator Algebras*, Proceedings of the Centre of Mathematical Analysis, Australian National University 15, Australian National University, Canberra, 1987, pp. 228–241.
- [Ri4] R. W. Richardson, *Conjugacy classes of n -tuples in Lie algebras and algebraic groups*, *Duke Math. J.* **57** (1988), no. 1, 1–35.
- [Ri5] R. W. Richardson, *Irreducible components of the nullcone*, in: *Invariant Theory* (Denton, TX, 1986), Contemporary Mathematics, Vol. 88, American Mathematical Society, Providence, RI, 1989, pp. 409–434.
- [TYu] P. Tauvel, R. W. T. Yu, *Lie Algebras and Algebraic groups*, Monographs in Mathematics, Springer, Berlin, 2005.
- [Ve] F. D. Veldkamp, *The centre of the universal enveloping algebra of a Lie algebra in characteristic p* , *Ann. Sci. École Norm. Sup.* **5** (1972), 217–240.