

GENERALIZED RANDOM FIELDS AND LÉVY'S CONTINUITY THEOREM ON THE SPACE OF TEMPERED DISTRIBUTIONS

HERMINE BIERMÉ, OLIVIER DURIEU, AND YIZAO WANG

ABSTRACT. In this note, we recall main properties of generalized random fields and present a proof of the continuity theorem of Paul Lévy for generalized random fields in the space of tempered distributions. This theorem was first proved by Fernique [6] in a more general setting. The aim of this note is to provide a self-contained proof that in particular avoids the abstract theory of nuclear spaces.

1. INTRODUCTION

The notion of generalized random fields has been introduced in the 50's by Gel'fand [7]. Such a field is a collection of real random variables that is indexed by a class of functions and that satisfies linearity and certain regularity conditions. It can also be seen as a single random variable taking values in the space of generalized functions (or distributions). The name *random distribution* is also used in the literature. The interest for generalized random fields was first due to their applications in many areas. They were used in constructive quantum field theory (see [3, 18]), in stochastic differential equations in infinite-dimensional spaces (see [14]), and in sparse stochastic modeling (see [4]). In comparison to classical probability theory, two theorems stand out for generalized random fields. The first is a Bochner-type theorem obtained by Minlos [17] that gives sufficient conditions on a functional on the Schwartz space to be the characteristic functional of some generalized random field. The second is a Lévy-type continuity theorem, saying that the convergence in distribution of generalized random fields can be determined by the pointwise convergence of the corresponding characteristic functionals, provided in addition that the limit is continuous at zero. The Lévy's continuity theorem was obtained by Fernique [6] and generalized by Meyer [16] (see also [2]) in the setting of nuclear spaces, and has found various applications in for example particle systems [10, 11] and image processing [5]. Our own motivation came from the investigation of limit theorems for the so-called random-ball models, which will be discussed in a forthcoming paper. In all these applications, Lévy's continuity theorem is used for generalized random fields in the space of tempered distributions, which is the dual of the Schwartz space.

Although Lévy's continuity theorem in very general setups has been known since Fernique [6] in the 60's, a simple and self-contained proof in the specific case of the

Date: June 27, 2017.

2010 Mathematics Subject Classification. 60G20; 60E10; 60B10.

Key words and phrases. Generalized random fields; tempered distributions; Lévy's continuity theorem; Minlos–Bochner's theorem.

space of tempered distributions, without using the general framework of nuclear spaces and their dual spaces, cannot be found in the literature. The space of tempered distributions as the dual of Schwartz space, a countably Hilbert space, enjoys several nice properties. Standard references as Gel'fand and Vilenkin [9] and Kallianpur and Xiong [14] provide excellent introductions to generalized random fields taking values in the space of tempered distributions. However, both treatments stop at Minlos–Bochner's theorem. It is the purpose of this paper to fill in the gap by presenting a proof of Lévy's continuity theorem for generalized random fields, without using abstract theory on nuclear spaces.

Throughout the paper, the integer $d \geq 1$ is fixed; \mathbb{N} stands for the set of non-negative integers $\{0, 1, 2, \dots\}$; for $n \in \mathbb{N}$, Γ_n denotes the net-square $\{0, \dots, n\}^d$; for each integer $s \geq 2$, $\zeta(s)$ denotes the finite constant $\sum_{n \geq 1} n^{-s}$.

2. GENERALIZED RANDOM FIELDS AND MAIN RESULTS

Recall that the *Schwartz space* $\mathcal{S} = \mathcal{S}(\mathbb{R}^d)$ consists of infinitely differentiable functions $f : \mathbb{R}^d \rightarrow \mathbb{R}$ that are rapidly decreasing, that is, for all $m \in \mathbb{N}$ and $j = (j_1, \dots, j_d) \in \mathbb{N}^d$,

$$(1) \quad \|f\|_{m,j} = \sup_{x \in \mathbb{R}^d} (1 + |x|)^m |D^j f(x)| < \infty,$$

where $D^j f(x) = \frac{\partial^{j_1} \dots \partial^{j_d}}{\partial x_1^{j_1} \dots \partial x_d^{j_d}} f(x)$ denotes the partial derivative of order j . It is a linear vector space and it is equipped with the topology generated by the family of semi-norms $\|\cdot\|_{m,j}$, $m \in \mathbb{N}$ and $j \in \mathbb{N}^d$. It is well known that \mathcal{S} is a Fréchet space (i.e. a locally convex metrizable complete space).

The *space of tempered distributions* $\mathcal{S}' = \mathcal{S}'(\mathbb{R}^d)$ is the topological dual of \mathcal{S} , the space of continuous linear functionals on \mathcal{S} . For $F \in \mathcal{S}'$ and $f \in \mathcal{S}$, the real number $F(f)$ is usually denoted by (F, f) . The duality bracket $(\cdot, \cdot) : \mathcal{S}' \times \mathcal{S} \rightarrow \mathbb{R}$ is a bilinear form.

We shall work with two topologies on \mathcal{S}' , the strong topology τ_s and the weak topology τ_w . The *strong topology* τ_s on \mathcal{S}' is generated by the family of semi-norms

$$q_B(F) = \sup_{f \in B} |(F, f)|, \quad B \subset \mathcal{S} \text{ bounded.}$$

Recall that a subset B of a topological vector space is bounded if for all neighborhood V of 0, there exists $\lambda > 0$ such that $B \subset \lambda V$. The *weak topology* on \mathcal{S}' is simply induced by the family of semi-norms $|(\cdot, \varphi)|$, $\varphi \in \mathcal{S}$. In particular F_n converges to F in \mathcal{S}' with respect to the weak topology, when $\lim_{n \rightarrow \infty} (F_n, \varphi) = (F, \varphi)$ for all $\varphi \in \mathcal{S}$. We show in Section 3.3 that the Borel σ -fields corresponding to both topologies coincide. Therefore we shall talk about the Borel σ -fields $\mathcal{B}(\mathcal{S}')$ of \mathcal{S}' without specifying the topology.

Let $(\Omega, \mathcal{A}, \mathbb{P})$ be a probability space. By a *generalized random field* defined on $(\Omega, \mathcal{A}, \mathbb{P})$, we refer to a random variable X with values in $(\mathcal{S}', \mathcal{B}(\mathcal{S}'))$, namely, a measurable map $X : (\Omega, \mathcal{A}) \rightarrow (\mathcal{S}', \mathcal{B}(\mathcal{S}'))$. In this case, $X(f)$ is a real-valued random variable for all $f \in \mathcal{S}$ and $f \in \mathcal{S} \mapsto X(f)(\omega)$ is linear and continuous for all $\omega \in \Omega$. Here, we are mostly interested in the law of the generalized random field

$$\mathbb{P}_X = \mathbb{P} \circ X^{-1},$$

the induced probability measure on $(\mathcal{S}', \mathcal{B}(\mathcal{S}'))$. We shall nevertheless talk about convergence of generalized random fields instead of convergence of laws of generalized random fields, for the sake of simplicity.

A fundamental object in the investigation of generalized random fields is the characteristic functional, playing the role as the characteristic function for a random variable in \mathbb{R} . First, recall that the law of a real random variable Y defined on $(\Omega, \mathcal{A}, \mathbb{P})$ is uniquely determined by its characteristic function \mathcal{L}_Y defined as

$$\mathcal{L}_Y(t) = \mathbb{E}(e^{itY}) = \int_{\Omega} e^{itY} d\mathbb{P}, \quad t \in \mathbb{R}.$$

The function \mathcal{L}_Y is positive-definite, continuous and satisfies $\mathcal{L}_Y(0) = 1$. Here we understand that a complex-valued function \mathcal{L} is *positive-definite* if for all $m \geq 1$, all complex numbers c_1, \dots, c_m , and all $t_1, \dots, t_m \in \mathbb{R}$, $\sum_{i,j=1}^m c_i \bar{c}_j \mathcal{L}(t_i - t_j) \geq 0$. The well known Bochner's theorem states that conversely, if a map $\mathcal{L} : \mathbb{R} \rightarrow \mathbb{C}$ is positive-definite, continuous at 0, and $\mathcal{L}(0) = 1$, then there exists a real random variable Y such that $\mathcal{L}_Y = \mathcal{L}$.

Similarly, for a generalized random field X with law \mathbb{P}_X , we define its *characteristic functional* by

$$\mathcal{L}_X(f) = \mathbb{E}(e^{iX(f)}) = \int_{\mathcal{S}'} e^{iF(f)} d\mathbb{P}_X(F), \quad f \in \mathcal{S}.$$

Note that \mathcal{L}_X is again positive-definite, continuous, and $\mathcal{L}_X(0) = 1$. Here and in the sequel, we say that a functional $\mathcal{L} : \mathcal{S} \rightarrow \mathbb{C}$ is positive-definite, if for all $m \geq 1$, $c_1, \dots, c_m \in \mathbb{C}$ and $f_1, \dots, f_m \in \mathcal{S}$,

$$\sum_{i=1}^m \sum_{j=1}^m c_i \bar{c}_j \mathcal{L}(f_i - f_j) \geq 0.$$

The Minlos–Bochner theorem says that the converse is true.

Theorem 2.1 (Minlos–Bochner). *If a functional $\mathcal{L} : \mathcal{S} \rightarrow \mathbb{C}$ is positive-definite, continuous at 0, and $\mathcal{L}(0) = 1$, there exists a generalized random field X defined on a probability space $(\Omega, \mathcal{A}, \mathbb{P})$ such that $\mathcal{L}_X = \mathcal{L}$.*

In order to be self-contained, we give a proof of this theorem in Section 4 as well as a similar result for random variables taking values in \mathcal{S} (Theorem 4.3), which will be needed in the proof of Lévy's continuity theorem. Minlos–Bochner's theorem has a direct application in proving existence of generalized random fields. For example, consider the functional $\mathcal{L} : \mathcal{S} \rightarrow \mathbb{C}$ defined by $\mathcal{L}(f) = \exp(-\|f\|_2^2/2)$, where $\|f\|_2$ denotes the L^2 -norm of f . Theorem 2.1 yields the existence of a generalized random field X such that $\mathcal{L}_X(\cdot) = \mathcal{L}(\cdot)$. This field X is known as the standard generalized Gaussian field in the literature (see e.g. [12, 15]).

More generally, there are situations that a collection of real random variables $X = (X(f))_{f \in \mathcal{S}}$ are given, such that for all $m \geq 1$, $a_1, \dots, a_m \in \mathbb{R}$, $f_1, \dots, f_m \in \mathcal{S}$,

$$(2) \quad X \left(\sum_{i=1}^m a_i f_i \right) = \sum_{i=1}^m a_i X(f_i), \quad \text{a.s.}$$

Although a priori this does not necessarily imply that $(X(f))_{f \in \mathcal{S}}$ is determined by a generalized random field X , Minlos–Bochner's theorem yields a simple criterion for X to admit a version as a generalized random field. By a version of X , we mean a generalized random field \tilde{X} , not necessarily defined on the same probability space

as X , such that $(X(f))_{f \in \mathcal{S}}$ and $(\tilde{X}(f))_{f \in \mathcal{S}}$ have the same finite-dimensional distributions. Given a collection of random variables $(X(f))_{f \in \mathcal{S}}$, with slightly abused notation we write $\mathcal{L}_X(f) = \mathbb{E}(e^{iX(f)})$. The linearity (2) shall imply that $\mathcal{L}_X(f)$ is positive-definite. We hence have the following.

Corollary 2.2. *Let $X = (X(f))_{f \in \mathcal{S}}$ be a collection of real random variables on $(\Omega, \mathcal{A}, \mathbb{P})$ that are linear in the sense of (2). If $\mathcal{L}_X(f)$ is continuous at zero on \mathcal{S} , then X admits a version that is a generalized random field.*

Our notion of version is weaker than the notion of regularization in [12] where \tilde{X} is required to be on the same probability space, and $\mathbb{P}(\tilde{X}(f) = X(f)) = 1$ for all $f \in \mathcal{S}$. However, our notion of version suffices for many applications on convergence in distribution of generalized random fields.

Another important theorem concerning characteristic functional is the Lévy's continuity theorem that characterizes convergence in distribution. Let $(X_n)_{n \geq 1}$ and X be generalized random fields with laws $(\mathbb{P}_{X_n})_{n \geq 1}$ and \mathbb{P}_X respectively. We say that X_n converges in distribution to X with respect to the strong topology (of \mathcal{S}') if

$$(3) \quad \lim_{n \rightarrow \infty} \int_{\mathcal{S}'} \varphi(F) d\mathbb{P}_{X_n}(F) = \int_{\mathcal{S}'} \varphi(F) d\mathbb{P}_X(F), \text{ for all } \varphi \in \mathcal{C}_b(\mathcal{S}', \tau_s),$$

where $\mathcal{C}_b(\mathcal{S}', \tau_s)$ is the space of bounded continuous functions on \mathcal{S}' given the strong topology.

The main result of the paper is a self-contained proof of the following version of Lévy's continuity theorem for generalized random fields.

Theorem 2.3. *Let $(X_n)_{n \geq 1}$ be a sequence of generalized random fields. If \mathcal{L}_{X_n} converges pointwise to a functional $\mathcal{L} : \mathcal{S} \rightarrow \mathbb{C}$ which is continuous at 0, then there exists a generalized random field X such that $\mathcal{L}_X = \mathcal{L}$ and X_n converges in distribution to X with respect to the strong topology.*

Another natural notion of convergence in the space \mathcal{S}' is the convergence in distribution with respect to the weak topology, which is defined similarly as in (3), with test functions from $\mathcal{C}_b(\mathcal{S}', \tau_w)$ instead of $\mathcal{C}_b(\mathcal{S}', \tau_s)$. It is straightforward to show that the convergence with respect to the strong topology implies the convergence with respect to the weak topology, and that the latter convergence implies the convergence of the characteristic functionals. The main theorem above shows that the three types of convergence are equivalent.

Corollary 2.4. *Let $(X_n)_{n \geq 1}$, X be generalized random fields. The following conditions are equivalent:*

- (i) X_n converges in distribution to X with respect to the strong topology,
- (ii) X_n converges in distribution to X with respect to the weak topology,
- (iii) $\lim_{n \rightarrow \infty} \mathcal{L}_{X_n}(f) = \mathcal{L}_X(f)$ for all $f \in \mathcal{S}$,
- (iv) $X_n(f)$ converges in distribution to $X(f)$ in \mathbb{R} for all $f \in \mathcal{S}$.

As we pointed out before, Theorem 2.3 and the discussion above are a special case of a more general result of [6, 16], where, however, the background on nuclear spaces are required. The aim of the paper is to present a simpler and self-contained proof in this special case of the space \mathcal{S}' that cannot be found in the literature, to the best of our knowledge.

The paper is organized as follows. Some preliminary results on Schwartz space and the space of tempered distributions are provided in Section 3. Section 4 establishes Milnos–Bochner's theorems. Section 5 proves the Lévy's continuity theorem for generalized random fields. Section 6 provides some topological properties of \mathcal{S} and \mathcal{S}' that are needed in the proofs.

3. THE SCHWARTZ SPACE AND THE SPACE OF TEMPERED DISTRIBUTIONS

3.1. The Schwartz space \mathcal{S} as a countably Hilbert space. A countable orthonormal basis (a complete orthonormal system) of $L^2(\mathbb{R})$ is given by the family of Hermite functions $\{h_n\}_{n \in \mathbb{N}}$, defined as

$$h_n(x) = (-1)^n (2^n n! \sqrt{\pi})^{-1/2} e^{\frac{x^2}{2}} \left(\frac{d}{dx} \right)^n e^{-x^2}, \quad n \in \mathbb{N}, x \in \mathbb{R}.$$

On \mathbb{R}^d , we define the Hermite functions $\{h_n\}_{n \in \mathbb{N}^d}$ by

$$h_n(x) = \prod_{i=1}^d h_{n_i}(x_i), \quad n = (n_1, \dots, n_d) \in \mathbb{N}^d, x = (x_1, \dots, x_d) \in \mathbb{R}^d.$$

The family of Hermite functions on \mathbb{R}^d is an orthonormal basis of $L^2(\mathbb{R}^d)$. Any function $f \in L^2(\mathbb{R}^d)$ can be written as

$$f = \sum_{n \in \mathbb{N}^d} \langle f, h_n \rangle h_n,$$

where $\langle \cdot, \cdot \rangle$ denotes the inner product in $L^2(\mathbb{R}^d)$ given by $\langle f, g \rangle = \int_{\mathbb{R}^d} f(x)g(x)dx$. We can then define a family of increasing norms on $L^2(\mathbb{R}^d)$ by

$$(4) \quad \|f\|_p^2 = \sum_{n \in \mathbb{N}^d} (1+n)^{2p} \langle f, h_n \rangle^2, \quad p \in \mathbb{N}, f \in L^2(\mathbb{R}^d),$$

where here and in the sequel, we write

$$(5) \quad (1+n)^q = \prod_{i=1}^d (1+n_i)^q \quad \text{for } q \in \mathbb{Z}, n = (n_1, \dots, n_d) \in \mathbb{N}^d.$$

For each $p \in \mathbb{N}$, we denote by $\mathcal{S}_p = \mathcal{S}_p(\mathbb{R}^d)$ the space of all $L^2(\mathbb{R}^d)$ -functions such that $\|f\|_p < \infty$. Note that each $\|\cdot\|_p$ is the norm associated to the inner product

$$\langle f, g \rangle_p = \sum_{n \in \mathbb{N}^d} (1+n)^{2p} \langle f, h_n \rangle \langle g, h_n \rangle.$$

The spaces $(\mathcal{S}_p)_{p \in \mathbb{N}}$ play a crucial role in our study.

For each $p \in \mathbb{N}$, \mathcal{S}_p is a separable Hilbert space with a countable orthonormal basis given by $\{h_n^{(p)} := (n+1)^{-p} h_n, n \in \mathbb{N}^d\}$. Note that \mathcal{S}_0 is just the space $L^2(\mathbb{R}^d)$, and for each $p \in \mathbb{N}$, $\mathcal{S}_{p+1} \subset \mathcal{S}_p$ with $\|\cdot\|_p \leq \|\cdot\|_{p+1}$. It is well known that $\mathcal{S} = \bigcap_{p \geq 0} \mathcal{S}_p$ and that both families of semi-norms $\{\|\cdot\|_{m,j} \mid m \in \mathbb{N}, j \in \mathbb{N}^d\}$ and $\{\|\cdot\|_p \mid p \in \mathbb{N}\}$ generate the same topology on \mathcal{S} . See for example [14, Theorem 1.3.2]. A basis of neighborhoods of 0 in \mathcal{S} is then given by the sets

$$(6) \quad B_p(r) = \{f \in \mathcal{S} \mid \|f\|_p < r\}, \quad p \in \mathbb{N}, r > 0.$$

Concerning the space of tempered distributions, we have $\mathcal{S}' = \bigcup_{p \geq 0} \mathcal{S}'_p$, where \mathcal{S}'_p is the dual of \mathcal{S}_p . On \mathcal{S}'_p , we consider the dual norm defined by

$$\|F\|'_p = \sup_{\|f\|_p \leq 1} |(F, f)| \text{ for all } F \in \mathcal{S}'_p.$$

3.2. The multi-sequence representation of \mathcal{S} and \mathcal{S}' . The representation of \mathcal{S} as a countably Hilbert space is convenient to establish the main theorems in Sections 4 and 5. As soon as we have this representation, we can think of the spaces \mathcal{S} and \mathcal{S}' as subspaces of $\mathbb{R}^{\mathbb{N}^d}$, the space of real-valued multi-sequences. It gives a more concrete description of these spaces and it simplifies our arguments in the sequel. In this section we recall this multi-sequence representation as introduced by Reed and Simon [19].

Let $\mathbb{R}^{\mathbb{N}^d}$ be the set of multi-sequences with real values. For $a = (a_n)_{n \in \mathbb{N}^d}$ and $b = (b_n)_{n \in \mathbb{N}^d}$ such that $\sum_{n \in \mathbb{N}^d} |a_n b_n| < \infty$, we denote

$$\langle a, b \rangle = \sum_{n \in \mathbb{N}^d} a_n b_n.$$

For all integer $p \in \mathbb{Z}$, and all multi-sequence $a = (a_n)_{n \in \mathbb{N}^d} \in \mathbb{R}^{\mathbb{N}^d}$, we set

$$\|a\|_p^2 = \sum_{n \in \mathbb{N}^d} (1+n)^{2p} a_n^2.$$

We define

$$\mathbb{S}_p = \{a \in \mathbb{R}^{\mathbb{N}^d} \mid \|a\|_p < \infty\}, \quad p \in \mathbb{Z}, \quad \text{and } \mathbb{S} = \bigcap_{p \in \mathbb{N}} \mathbb{S}_p.$$

Note that for each $p \in \mathbb{Z}$, \mathbb{S}_p is a Hilbert space with respect to the inner product $\langle a, b \rangle_p = \sum_{n \in \mathbb{N}^d} (1+n)^{2p} a_n b_n$, and $\|a\|_p^2 = \langle a, a \rangle_p$. The set \mathbb{S} is then given the topology generated by the family of Hilbert norms $\|\cdot\|_p$, $p \in \mathbb{Z}$.

For each $p \in \mathbb{Z}$, we let \mathbb{S}'_p denote the dual space of the space \mathbb{S}_p . The dual norm on \mathbb{S}'_p is given by $\|F\|'_p = \sup_{\|a\|_p \leq 1} |(F, a)|$, and we have $\mathcal{S}' = \bigcup_{p \geq 0} \mathbb{S}'_p$.

Lemma 3.1. *For each $p \in \mathbb{Z}$, the map $b \mapsto \langle b, \cdot \rangle$ is an isometry from \mathbb{S}_{-p} to \mathbb{S}'_p .*

Proof. Let ϕ denote the map of interest $b \mapsto \langle b, \cdot \rangle$. Note that, by Cauchy–Schwarz inequality, for $a \in \mathbb{S}_p$ and $b \in \mathbb{S}_{-p}$,

$$|\langle b, a \rangle| \leq \|a\|_p \|b\|_{-p},$$

and $\|\phi(b)\|'_p = \sup_{\|a\|_p \leq 1} |\langle b, a \rangle| \leq \|b\|_{-p}$. Thus ϕ maps \mathbb{S}_{-p} into \mathbb{S}'_p . Further, for $b \in \mathbb{S}_{-p}$ by setting $a = (a_n)_{n \in \mathbb{N}^d}$ with $a_n = \|b\|_{-p}^{-1} (n+1)^{-2p} b_n$ we have $\|a\|_p = 1$ and thus

$$\|\phi(b)\|'_p \geq |\langle b, a \rangle| = \|b\|_{-p}^{-1} \sum_{n \in \mathbb{N}^d} (n+1)^{-2p} b_n^2 = \|b\|_{-p}.$$

Thus, for all $b \in \mathbb{S}_{-p}$, $\|\phi(b)\|'_p = \|b\|_{-p}$.

Now, to see that ϕ is surjective, suppose $F \in \mathbb{S}'_p$, $F \neq 0$, let c be an element in \mathbb{S}_p such that $\|c\|_p = 1$ and c is orthogonal to $\text{Ker}(F)$. Define $b = (b_n)_{n \in \mathbb{N}^d}$ with $b_n = F(c) c_n (1+n)^{2p}$ for $n \in \mathbb{N}^d$. Then, it follows that $\|b\|_{-p} = F(c) < \infty$, hence $b \in \mathbb{S}_{-p}$. Now for all $a \in \mathbb{S}_p$, write $a = a_1 + a_2$ with $a_1 = (F(a)/F(c))c$ and $a_2 = a - a_1 \in \text{Ker}(F)$. Then $\langle b, a \rangle = \langle b, a_1 \rangle = F(a)$. \square

In the sequel, we identify the spaces \mathbb{S}'_p and \mathbb{S}_{-p} . We thus have $\mathbb{S}' = \bigcup_{p \geq 0} \mathbb{S}_{-p} \subset \mathbb{R}^{\mathbb{N}^d}$ and then the duality bracket (\cdot, \cdot) corresponds to $\langle \cdot, \cdot \rangle$.

Lemma 3.2. *Let $b = (b_n)_{n \in \mathbb{N}^d} \in \mathbb{R}^{\mathbb{N}^d}$. Then $b \in \mathbb{S}'$ if and only if there exist $p \in \mathbb{N}$ and $c > 0$ such that for all $n \in \mathbb{N}^d$, $|b_n| \leq c(1+n)^p$.*

Proof. If $b \in \mathbb{S}'$, there exists $p \in \mathbb{N}$ such that $b \in \mathbb{S}_{-p}$. Then, for all $n \in \mathbb{N}^d$, $(1+n)^{-p}|b_n| \leq \|b\|_{-p}$ and the first implication, with $c = \|b\|_{-p}$, is proved.

Conversely, if there exist $p \in \mathbb{N}$ and $c > 0$ such that for all $n \in \mathbb{N}^d$, $|b_n| \leq c(1+n)^p$, then

$$\|b\|_{-p-1}^2 = \sum_{n \in \mathbb{N}^d} (n+1)^{-2p-2} b_n^2 \leq c^2 \sum_{n \in \mathbb{N}^d} (n+1)^{-2} = c^2 \zeta(2) < \infty.$$

Thus $b \in \mathbb{S}'_{p+1} \subset \mathbb{S}'$. \square

Now, we show that \mathbb{S} is topologically isomorphic to \mathcal{S} . Recall that two topological vector spaces are said to be *topologically isomorphic* if there exists a linear isomorphism from one to the other which is also a homeomorphism (a bi-continuous map). Such a map is called a *topological isomorphism*.

Proposition 3.3. *The map $\Phi : \mathcal{S} \rightarrow \mathbb{S}$, $f \mapsto (\langle f, h_n \rangle)_{n \in \mathbb{N}^d}$ is a topological isomorphism.*

Proof. For any $f \in \mathcal{S}$ and $p \in \mathbb{N}$,

$$\|\Phi(f)\|_p^2 = \sum_{n \in \mathbb{N}^d} (1+n)^{2p} \langle f, h_n \rangle^2 = \|f\|_p^2 < \infty.$$

Thus the linear map Φ is well-defined and continuous. Further, each $\|\cdot\|_p$ being a norm on \mathcal{S} , the map Φ is injective. Let us show that it is also surjective. Let $a = (a_n)_{n \in \mathbb{N}^d} \in \mathbb{S}$ and define $f_m = \sum_{n \in \Gamma_m} a_n h_n$, for all $m \in \mathbb{N}$. For all $p \in \mathbb{N}$ and $k \in \mathbb{N}$,

$$\|f_{m+k} - f_m\|_p^2 \leq \sum_{n \notin \Gamma_m} (n+1)^{2p} a_n^2 \rightarrow 0 \text{ as } m \rightarrow \infty.$$

Thus $(f_m)_{m \in \mathbb{N}}$ is a Cauchy sequence in \mathcal{S} and then it converges to $f \in \mathcal{S}$. By continuity of Φ , we see that $\Phi(f) = a$. The equalities $\|\Phi(f)\|_p = \|f\|_p$, $p \in \mathbb{N}$, prove that Φ is a homeomorphism. \square

We can also introduce a topology on \mathbb{S}' to make it topologically isomorphic to (\mathbb{S}', τ_s) , thanks to the following result.

Proposition 3.4. *The map $\Psi : \mathbb{S}' \rightarrow \mathbb{S}'$, $F \mapsto ((F, h_n))_{n \in \mathbb{N}^d}$ is a linear isomorphism.*

Proof. For all $F \in \mathbb{S}'$ and $n \in \mathbb{N}^d$, there exists $p \in \mathbb{N}$, such that $F \in \mathbb{S}'_p$ and then, for all $n \in \mathbb{N}^d$,

$$|(F, h_n)| \leq \|F\|'_p \|h_n\|_p = \|F\|'_p (n+1)^p.$$

Thus, by Lemma 3.2, the linear map Ψ is well-defined. Further, Ψ is injective, since $\Psi(F) = 0$ implies that $(F, h_n) = 0$ for all $n \in \mathbb{N}^d$, and then by linearity of F , it implies $F(f) = 0$ for all $f \in \mathcal{S}$. It remains to show that Ψ is surjective. Let $b = (b_n)_{n \in \mathbb{N}^d} \in \mathbb{S}'$ and hence, $b \in \mathbb{S}_{-p}$ for some $p \in \mathbb{N}$. We define the map $F : \mathcal{S} \rightarrow \mathbb{R}$, $f \mapsto \langle b, \Phi(f) \rangle$, where Φ is the topological isomorphism in Proposition 3.3. Then, F is a continuous linear map from \mathcal{S} to \mathbb{R} . Indeed, the linearity is clear and the

continuity comes from $|\langle b, \Phi(f) \rangle| \leq \|b\|_{-p} \|\Phi(f)\|_p = \|b\|_{-p} \|f\|_p$. Thus $F \in \mathcal{S}'$. It is clear that $\Psi(F) = b$. \square

As a consequence, (\mathcal{S}', τ_s) and $(\mathbb{S}', \Psi(\tau_s))$ are topologically isomorphic. The topology $\pi_s = \Psi(\tau_s)$ on \mathbb{S}' is called the strong topology on \mathbb{S}' . It is generated by the semi-norms

$$q_B(\cdot) = \sup_{a \in B} |\langle \cdot, a \rangle|, \quad B \text{ bounded in } \mathbb{S}.$$

In the same way, (\mathcal{S}', τ_w) and $(\mathbb{S}', \Psi(\tau_w))$ are topologically isomorphic. The topology $\pi_w = \Psi(\tau_w)$ on \mathbb{S}' is called the weak topology on \mathbb{S}' . It is generated by the semi-norms $|\langle \cdot, a \rangle|$, $a \in \mathbb{S}$. In the sequel, if we do not precise, \mathbb{S}' is assumed given the strong topology.

The following facts will be useful in the next section.

Proposition 3.5. \mathbb{S} , \mathbb{S}' , and hence \mathcal{S} and \mathcal{S}' are separable spaces.

Proof. The set of elements of $\mathbb{R}^{\mathbb{N}^d}$ with rational coordinates and only finitely many non-zero ones is dense in \mathbb{S}_p for all $p \in \mathbb{Z}$. It is then dense in both \mathbb{S} and \mathbb{S}' . \square

Lemma 3.6. For each $p \in \mathbb{N}$, the injection map $i_p : \mathbb{S}'_p \rightarrow \mathbb{S}'$ is continuous with respect to the strong topology on \mathbb{S}' (and hence also with respect to the weak topology).

Proof. Let \mathbb{S}' be given the strong topology. Let $V = \{b \in \mathbb{S}' \mid q_B(b) < \varepsilon\}$ for some bounded set $B \subset \mathbb{S}$ and some $\varepsilon > 0$. For each $b_0 \in i_p^{-1}(V)$, we have $q_B(b_0) < \varepsilon$ and for all $b \in \mathbb{S}'_p$,

$$\begin{aligned} q_B(b) &\leq \sup_{a \in B} |\langle b - b_0, a \rangle| + \sup_{a \in B} |\langle b_0, a \rangle| \\ &\leq \|b - b_0\|'_p \sup_{a \in B} \|a\|_p + q_B(b_0). \end{aligned}$$

Thus, if $\|b - b_0\|'_p < \frac{\varepsilon - q_B(b_0)}{\sup_{a \in B} \|a\|_p + 1}$, then $b \in i_p^{-1}(V)$. \square

3.3. Cylinder σ -fields and Borel σ -fields. On the space $\mathbb{R}^{\mathbb{N}^d}$, the product σ -field generated by the sets $\{a \in \mathbb{R}^{\mathbb{N}^d} \mid (a_i)_{i \in \Gamma} \subset B\}$ for all finite subsets Γ of \mathbb{N}^d and all Borel sets $B \in \mathcal{B}(\mathbb{R}^{\#\Gamma})$ is the same as the Borel σ -field associated to the product topology on $\mathbb{R}^{\mathbb{N}^d}$ (see [13, Lemma 1.2]). It is denoted by $\mathcal{B}(\mathbb{R}^{\mathbb{N}^d})$.

On \mathbb{S} , the cylinder σ -field, denoted by $\mathcal{C}(\mathbb{S})$, is generated by the cylinder sets

$$\{a \in \mathbb{S} \mid (\langle b_1, a \rangle, \dots, \langle b_m, a \rangle) \in B\},$$

for all $m \geq 1$, $b_1, \dots, b_m \in \mathbb{S}'$, $B \in \mathcal{B}(\mathbb{R}^m)$. We easily see that $\mathcal{C}(\mathbb{S})$ is the σ -field induced by $\mathcal{B}(\mathbb{R}^{\mathbb{N}^d})$ on \mathbb{S} .

On \mathbb{S}' , the cylinder σ -field, denoted by $\mathcal{C}(\mathbb{S}')$, is generated by the cylinder sets

$$\{b \in \mathbb{S}' \mid (\langle b, a_1 \rangle, \dots, \langle b, a_m \rangle) \in B\},$$

for all $m \geq 1$, $a_1, \dots, a_m \in \mathbb{S}$, $B \in \mathcal{B}(\mathbb{R}^m)$. Again, $\mathcal{C}(\mathbb{S}')$ is the σ -field induced by $\mathcal{B}(\mathbb{R}^{\mathbb{N}^d})$ on \mathbb{S}' . Further, by definition, we see that $\mathcal{C}(\mathbb{S}')$ is also the Borel σ -field corresponding to the weak topology π_w on \mathbb{S}' .

Recall that, in our setting, \mathbb{S} is given the topology generated by the family of norms $\|\cdot\|_p$, $p \in \mathbb{N}$, and \mathbb{S}' is given the strong topology π_s . The corresponding Borel σ -fields are denoted by $\mathcal{B}(\mathbb{S})$ and $\mathcal{B}(\mathbb{S}')$, respectively. Of course, we have

$$\Phi^{-1}(\mathcal{B}(\mathbb{S})) = \mathcal{B}(\mathcal{S}) \quad \text{and} \quad \Psi^{-1}(\mathcal{B}(\mathbb{S}')) = \mathcal{B}(\mathcal{S}')$$

for the topological isomorphisms Φ and Ψ from Propositions 3.3 and 3.4, and where $\mathcal{B}(\mathcal{S})$ and $\mathcal{B}(\mathcal{S}')$ are respectively the Borel σ -fields of \mathcal{S} and \mathcal{S}' given the strong topology τ_s .

Proposition 3.7. $\mathcal{C}(\mathbb{S}) = \mathcal{B}(\mathbb{S})$.

Proof. Recall that a countable basis of neighborhoods of $0 \in \mathbb{S}$ is given by the sets $B_p(1/k)$, $p \in \mathbb{N}$, $k \geq 1$ (see (6)). Note that, by Lemma 3.1, $B_p(1/k)$ is an open set that can be written as

$$\begin{aligned} B_p(1/k) &= \left\{ a \in \mathbb{S} \mid \sup_{\|b\|_{-p} \leq 1} |\langle b, a \rangle| < 1/k \right\} \\ &= \bigcup_{n \geq 1} \bigcap_{b \in D_{-p}} \{a \in \mathbb{S} \mid |\langle b, a \rangle| < 1/k - 1/n\}, \end{aligned}$$

where D_{-p} is a countable dense subset in the unit ball of \mathbb{S}_{-p} . So, each $B_p(1/k)$ belongs to $\mathcal{C}(\mathbb{S})$. Now, since \mathbb{S} is separable, there exists a countable dense subset $D \subset \mathbb{S}$, and for any open set U in \mathbb{S} , we have

$$U = \bigcup_{\substack{a \in D, p \in \mathbb{N}, k \geq 1 \\ \text{s.t. } a + B_p(1/k) \subset U}} (a + B_p(1/k))$$

which is a countable union of elements of $\mathcal{C}(\mathbb{S})$, and thus $\mathcal{B}(\mathbb{S}) \subset \mathcal{C}(\mathbb{S})$.

The converse inclusion comes from the fact that the maps $f_b : a \in \mathbb{S} \mapsto \langle b, a \rangle$, $b \in \mathbb{S}'$, are continuous on \mathbb{S} and thus $\mathcal{B}(\mathbb{S})$ -measurable. \square

Proposition 3.8. $\mathcal{C}(\mathbb{S}') = \mathcal{B}(\mathbb{S}')$.

Proof. Since for all $a \in \mathbb{S}$, the map $g_a : b \in \mathbb{S}' \mapsto \langle b, a \rangle$ is continuous on \mathbb{S}' , the inclusion $\mathcal{C}(\mathbb{S}') \subset \mathcal{B}(\mathbb{S}')$ is clear.

For the converse, let V be an open set in \mathbb{S}' with respect to π_s . By Lemma 3.6, for each $p \in \mathbb{N}$, $V_p = V \cap \mathbb{S}'_p$ is open in \mathbb{S}'_p , and we can write $V = \bigcup_{p \in \mathbb{N}} V_p$. To conclude we prove that open sets in \mathbb{S}'_p (with respect to the topology generated by $\|\cdot\|'_p$) belong to $\mathcal{C}(\mathbb{S})$. Indeed, for each $p \in \mathbb{N}$ and $r > 0$, consider

$$B_{-p}(r) = \{b \in \mathbb{S}' \mid \|b\|_{-p} < r\} = \bigcup_{n \geq 1} \bigcap_{a \in D_p} \{b \in \mathbb{S}' \mid |\langle b, a \rangle| < r - 1/n\}$$

where D_p a countable dense subset of the unit ball of \mathbb{S}_p , and we see that $B_{-p}(r) \in \mathcal{C}(\mathbb{S}')$. A countable basis of the topology of \mathbb{S}'_p is given by the sets $b + B_{-p}(1/k)$, for all $k \geq 1$ and b in a given countable dense subset of \mathbb{S}'_p . We infer that $V_p \in \mathcal{C}(\mathbb{S})$. \square

Further, since the cylinder σ -field on \mathbb{S}' corresponds to the Borel σ -field of the weak topology, we have the following corollary.

Corollary 3.9. *On \mathbb{S}' (and \mathcal{S}'), the Borel σ -field of the strong topology and the Borel σ -field of the weak topology coincide.*

4. MINLOS–BOCHNER’S THEOREMS

In this section, we study the existence of generalized random fields. Thanks to Proposition 3.4, in order to define a random variable with value in $(\mathcal{S}', \mathcal{B}(\mathcal{S}'))$, we can define a random variable X with values in $(\mathbb{S}', \mathcal{B}(\mathbb{S}'))$ and then consider $\Psi(X)$. Conversely, for any generalized random field Y with values in $(\mathcal{S}', \mathcal{B}(\mathcal{S}'))$, the variable $X = \Psi^{-1}(Y)$ is a random variable with values in $(\mathbb{S}', \mathcal{B}(\mathbb{S}'))$. By extension, we also call X a generalized random field and we have

- for all $\omega \in \Omega$, $X(\omega) \in \mathbb{S}'$;
- for all $a \in \mathbb{S}$, $\langle X, a \rangle : \omega \mapsto \langle X(\omega), a \rangle$ is a real random variable.

We see that a generalized random field X provides a collection $(\langle X, a \rangle)_{a \in \mathbb{S}}$ of real random variables. The finite-dimensional distributions of the generalized random field X on $(\Omega, \mathcal{A}, \mathbb{P})$ are given by the law of the real vectors $(\langle X, a_1 \rangle, \dots, \langle X, a_m \rangle)$ for all $m \geq 1$, $a_1, \dots, a_m \in \mathcal{S}$. By linearity, the finite-dimensional distributions are determined by the characteristic functional of X defined by

$$\mathcal{L}_X(a) = \mathbb{E}(e^{i\langle X, a \rangle}) = \int_{\Omega} e^{i\langle X, a \rangle} d\mathbb{P}, \quad a \in \mathbb{S}.$$

More generally, for a probability measure μ on $(\mathbb{R}^{\mathbb{N}^d}, \mathcal{B}(\mathbb{R}^{\mathbb{N}^d}))$, we call the characteristic functional of μ the functional

$$\mathcal{L}_\mu(a) = \int_{\mathbb{R}^{\mathbb{N}^d}} e^{i\langle b, a \rangle} d\mu(b),$$

defined for multi-sequences a such that $\langle b, a \rangle$ is finite for μ -almost all $b \in \mathbb{R}^{\mathbb{N}^d}$. It is always defined for multi-sequences $a \in \mathbb{R}^{\mathbb{N}^d}$ with finitely many non-zero coordinates. In particular, for a generalized random field X with law $\mu = \mathbb{P} \circ X^{-1}$, we have $\mathcal{L}_X = \mathcal{L}_\mu$.

Theorem 4.1. *If a functional $\mathcal{L} : \mathbb{S} \rightarrow \mathbb{C}$ is positive-definite, continuous at 0, and $\mathcal{L}(0) = 1$, then it is the characteristic functional of a generalized random field on \mathbb{S}' .*

Theorem 2.1 and Corollary 2.2 are direct consequences. To prove the present theorem, we need the following key lemma.

Lemma 4.2. *Let μ be a Borel probability measure on $\mathbb{R}^{\mathbb{N}^d}$, $p \in \mathbb{Z}$, $c > 0$, and $\varepsilon > 0$ such that for all $a \in \mathbb{R}^{\mathbb{N}^d}$ with finitely many non-zero coordinates,*

$$1 - \operatorname{Re} \mathcal{L}_\mu(a) \leq \varepsilon + c \|a\|_p^2.$$

Then for all $q > p$, and $\sigma > 0$,

$$\int_{\mathbb{R}^{\mathbb{N}^d}} 1 - \exp\left(-\frac{\sigma^2}{2} \|b\|_{-q}^2\right) d\mu(b) \leq \varepsilon + c\sigma^2 \zeta(2(q-p))^d.$$

In particular,

$$\mu(\mathbb{S}_{-q}) \geq 1 - \varepsilon.$$

Proof. Recall the expression of $(1+n)^q$ in (5). For all $\sigma > 0$ and $m \in \mathbb{N}$, we consider the Gaussian probability measure $\nu_{m,\sigma}$ on \mathbb{R}^{Γ_m} defined by, for $a = (a_n)_{n \in \Gamma_m} \in \mathbb{R}^{\Gamma_m}$,

$$d\nu_{m,\sigma}(a) = \bigotimes_{n \in \Gamma_m} (2\pi\sigma^2(n+1)^{-2q})^{-1/2} \exp\left(-\frac{a_n^2}{2\sigma^2(n+1)^{-2q}}\right) da_n.$$

Then $\nu_{m,\sigma}$ satisfies:

- (i) $\nu_{m,\sigma}(\mathbb{R}^{\Gamma_m}) = 1$,
- (ii) $\int_{\mathbb{R}^{\Gamma_m}} a_n a_m d\nu_{m,\sigma}(a) = \sigma^2 (n+1)^{-2q} \mathbf{1}_{\{n=m\}}$, and
- (iii) $\int_{\mathbb{R}^{\Gamma_m}} e^{i\langle b,a \rangle} d\nu_{m,\sigma}(a) = \exp\left(-\frac{\sigma^2}{2} \sum_{n \in \Gamma_m} (n+1)^{-2q} b_n^2\right)$.

By Fubini's theorem, we have

$$\begin{aligned}
& \int_{\mathbb{R}^{\mathbb{N}^d}} 1 - \exp\left(-\frac{\sigma^2}{2} \sum_{n \in \Gamma_m} (n+1)^{-2q} b_n^2\right) d\mu(b) \\
&= \int_{\mathbb{R}^{\mathbb{N}^d}} \int_{\mathbb{R}^{\Gamma_m}} 1 - e^{i\langle b,a \rangle} d\nu_{m,\sigma}(a) d\mu(b) \\
&= \int_{\mathbb{R}^{\Gamma_m}} \int_{\mathbb{R}^{\mathbb{N}^d}} 1 - e^{i\langle b,a \rangle} d\mu(b) d\nu_{m,\sigma}(a) \\
&= \int_{\mathbb{R}^{\Gamma_m}} 1 - \mathcal{L}_\mu(a) d\nu_{m,\sigma}(a),
\end{aligned}$$

which is then a real quantity, and by assumption,

$$\begin{aligned}
\int_{\mathbb{R}^{\Gamma_m}} 1 - \mathcal{L}_\mu(a) d\nu_{m,\sigma}(a) &\leq \varepsilon + c \int_{\mathbb{R}^{\Gamma_m}} \|a\|_p^2 d\nu_{m,\sigma}(a) \\
&= \varepsilon + c \sum_{n \in \Gamma_m} (n+1)^{2p} \int_{\mathbb{R}^{\Gamma_m}} a_n^2 d\nu_{m,\sigma}(a) \\
&= \varepsilon + c\sigma^2 \sum_{n \in \Gamma_m} (n+1)^{-2(q-p)}.
\end{aligned}$$

Since $\sum_{n \in \Gamma_m} (n+1)^{-2(q-p)} \uparrow \zeta(2(q-p))^d$ as $m \rightarrow \infty$, by dominated convergence, we get

$$\int_{\mathbb{R}^{\mathbb{N}^d}} 1 - \exp\left(-\frac{\sigma^2}{2} \sum_{n \in \mathbb{N}^d} (n+1)^{-2q} b_n^2\right) d\mu(b) \leq \varepsilon + c\sigma^2 \zeta(2(q-p))^d.$$

Now, since $\exp(-\sigma^2 \|b\|_{-q}^2 / 2) = 0$ for all $b \notin \mathbb{S}_{-q}$, it follows that

$$\mu(\mathbb{S}_{-q}) \geq \int_{\mathbb{R}^{\mathbb{N}^d}} \exp\left(-\frac{\sigma^2}{2} \|b\|_{-q}^2\right) d\mu(b) \geq 1 - \varepsilon - c\sigma^2 \zeta(2(q-p))^d.$$

The last desired statement follows by taking $\sigma \downarrow 0$. \square

Proof of Theorem 4.1. Let $\mathcal{L} : \mathbb{S} \rightarrow \mathbb{C}$ be positive-definite, continuous at 0, and $\mathcal{L}(0) = 1$. For any finite subset Γ of \mathbb{N}^d , we can consider the restriction of \mathcal{L} to $\mathbb{R}^\Gamma \subset \mathbb{S}$ (by completing the multi-sequences of \mathbb{R}^Γ by zeros). Thus, by Bochner's theorem, there exists a unique probability measure μ_Γ on the product σ -field of \mathbb{R}^Γ , such that, for all $a \in \mathbb{R}^\Gamma$,

$$\mathcal{L}(a) = \int_{\mathbb{R}^\Gamma} e^{i\langle b,a \rangle} d\mu_\Gamma(b).$$

Further, if $\Gamma \subset \Gamma'$ are finite subsets of \mathbb{N}^d and $\pi_{\Gamma',\Gamma}$ is the canonical projection from $\mathbb{R}^{\Gamma'}$ to \mathbb{R}^Γ , we have $\mu_{\Gamma'} \circ \pi_{\Gamma',\Gamma}^{-1} = \mu_\Gamma$. By Kolomogorov's extension theorem [13, Theorem 5.16] there exists a unique probability measure μ on the product σ -field of $\mathbb{R}^{\mathbb{N}^d}$ such that for all finite Γ , $\mu \circ \pi_\Gamma^{-1} = \mu_\Gamma$, where π_Γ is the canonical projection

from $\mathbb{R}^{\mathbb{N}^d}$ to \mathbb{R}^Γ . Since any element $a \in \mathbb{S}$ can be approximated by multi-sequences with only finitely many non-zero coefficients, by dominated convergence and the fact that \mathcal{L} is continuous, we get that for all $a \in \mathbb{S}$, $\mathcal{L}(a) = \mathcal{L}_\mu(a)$.

It remains to show that $\mu(\mathbb{S}') = 1$. Fix $\varepsilon > 0$. The continuity of \mathcal{L} implies that there exists $p \in \mathbb{N}$ and $\delta > 0$,

$$1 - \operatorname{Re} \mathcal{L}(a) \leq \varepsilon, \quad \text{for all } a \in \mathbb{S} \text{ such that } \|a\|_p < \delta.$$

Thus,

$$1 - \operatorname{Re} \mathcal{L}_\mu(a) \leq \varepsilon + 2\delta^{-2}\|a\|_p^2, \quad \text{for all } a \in \mathbb{S}.$$

By Lemma 4.2 for $q = p + 1$, we have $\mu(\mathbb{S}_{-q}) \geq 1 - \varepsilon$. We infer $\mu(\mathbb{S}') \geq 1 - \varepsilon$ for all $\varepsilon > 0$, that is $\mu(\mathbb{S}') = 1$. \square

In order to prove Lévy's continuity theorem in the next section, we need another Minlos–Bochner's theorem. Now, we consider random variables with values in $(\mathbb{S}, \mathcal{B}(\mathbb{S}))$. For such a random variable X , one can similarly define its characteristic functional by

$$\mathcal{L}_X(b) = \mathcal{L}_\mu(a) = \int_{\mathbb{S}} e^{i\langle b, a \rangle} d\mu(a), \quad b \in \mathbb{S}',$$

where $\mu = \mathbb{P} \circ X^{-1}$ is the law of X on $(\mathbb{S}, \mathcal{B}(\mathbb{S}))$.

Theorem 4.3. *If a functional $\mathcal{L} : \mathbb{S}' \rightarrow \mathbb{C}$ is positive-definite, continuous at 0 (with respect to the topology π_s), and $\mathcal{L}(0) = 1$, then it is the characteristic functional of a random variable on $(\mathbb{S}, \mathcal{B}(\mathbb{S}))$.*

Proof. Let $\mathcal{L} : \mathbb{S}' \rightarrow \mathbb{C}$ be as in the statement. As we did in the proof of Theorem 4.1, using Bochner's theorem and Kolmogorov's extension theorem, we can show that there exists a probability measures μ on $\mathcal{B}(\mathbb{R}^{\mathbb{N}^d})$ such that for all $b \in \mathbb{S}'$, $\mathcal{L}_\mu(b) = \mathcal{L}(b)$.

Again, it remains to show that $\mu(\mathbb{S}) = 1$. Fix $\varepsilon > 0$. Since \mathcal{L} is continuous on \mathbb{S}' , by Lemma 3.6, its restriction to \mathbb{S}_{-p} is also continuous for each $p \in \mathbb{N}$. Thus, for all $p \in \mathbb{N}$, there exists $\delta_p > 0$,

$$1 - \operatorname{Re} \mathcal{L}(b) \leq \varepsilon, \quad \text{for all } b \in \mathbb{S}_{-p} \text{ such that } \|b\|_{-p} < \delta_p.$$

Thus,

$$1 - \operatorname{Re} \mathcal{L}_\mu(b) \leq \varepsilon + 2\delta_p^{-2}\|b\|_{-p}^2, \quad \text{for all } b \in \mathbb{S}_{-p}.$$

By Lemma 4.2 for $q = -p + 1$, $\mu(\mathbb{S}_{p-1}) \geq 1 - \varepsilon$. Since this is true for all $p \geq 1$, and since the sets $(\mathbb{S}_p)_{p \in \mathbb{N}}$ form a non-increasing sequence, we get

$$\mu(\mathbb{S}) = \mu \left(\bigcap_{p \geq 0} \mathbb{S}_p \right) \geq 1 - \varepsilon.$$

Finally, $\varepsilon > 0$ being arbitrary, $\mu(\mathbb{S}) = 1$. \square

5. LÉVY'S CONTINUITY THEOREM

For Borel probability measures $(\mu_n)_{n \geq 1}$ and μ on $(\mathbb{S}', \mathcal{B}(\mathbb{S}'))$, we say that μ_n converges weakly to μ with respect to the strong topology (of \mathbb{S}'), if

$$\lim_{n \rightarrow \infty} \int_{\mathbb{S}'} \varphi(b) d\mu_n(b) = \int_{\mathbb{S}'} \varphi(b) d\mu(b), \quad \text{for all } \varphi \in \mathcal{C}_b(\mathbb{S}', \pi_s).$$

Accordingly, for generalized random fields $(X_n)_{n \geq 1}$ and X in \mathbb{S}' , we say X_n converges in distribution to X with respect to the strong topology (of \mathbb{S}'), if the corresponding probability law converges weakly. Thanks to Proposition 3.4, to prove Theorem 2.3 it is equivalent to prove the following.

Theorem 5.1. *If $(\mu_n)_{n \geq 1}$ is a sequence of Borel probability measures on \mathbb{S}' such that \mathcal{L}_{μ_n} converges pointwise to a functional $\mathcal{L} : \mathbb{S} \rightarrow \mathbb{C}$ which is continuous at 0, then there exists a Borel probability measure μ on \mathbb{S}' such that $\mathcal{L}_\mu = \mathcal{L}$ and μ_n converges weakly to μ with respect to the strong topology.*

Proof. Let $(\mu_n)_{n \geq 1}$ be as in the statement. By pointwise convergence, \mathcal{L} is also positive-definite and $\mathcal{L}(0) = 1$. By Theorem 4.1, there exists a Borel probability measure μ on \mathbb{S}' such that $\mathcal{L}_\mu = \mathcal{L}$.

Let $\mathcal{P}(\mathbb{S}')$ denote the set of all Borel probability measures on \mathbb{S}' . The main step of the proof is to show that, under the assumptions, the sequence $(\mu_n)_{n \geq 1}$ is tight in $\mathcal{P}(\mathbb{S}')$, i.e. for all $\varepsilon > 0$, there exists a strongly compact set K in \mathbb{S}' such that $\mu_n(K) \geq 1 - \varepsilon$ for all $n \geq 1$.

Assume we have proved it and let us conclude the proof. From tightness, we infer that the sequence $(\mu_n)_{n \geq 1}$ is relatively sequentially compact with respect to the topology of the weak convergence in $\mathcal{P}(\mathbb{S}')$, see [14, Theorem 2.2.1]. Thus, from any subsequence of $(\mu_n)_{n \geq 1}$, we can extract a sub-subsequence that converges weakly to a probability measure $\nu \in \mathcal{P}(\mathbb{S}')$. But, by weak convergence, the characteristic functional of ν is necessarily the pointwise limit of $(\mathcal{L}_{\mu_n})_{n \geq 1}$, which is \mathcal{L} . Hence $\nu = \mu$ and we conclude that μ_n converges weakly to μ .

It remains to prove that $(\mu_n)_{n \geq 1}$ is tight in $\mathcal{P}(\mathbb{S}')$. This follows from the following two lemmas. \square

Lemma 5.2. *Let $(\mu_n)_{n \geq 1}$ be a sequence of Borel probability measures on \mathbb{S}' . If $(\text{Re } \mathcal{L}_{\mu_n})_{n \geq 1}$ is equicontinuous on \mathbb{S} , then $(\mu_n)_{n \geq 1}$ is tight in $\mathcal{P}(\mathbb{S}')$.*

Proof. Let $\varepsilon > 0$. The equicontinuity of $(\text{Re } \mathcal{L}_{\mu_n})_{n \geq 1}$ implies that there exist $p \in \mathbb{N}$ and $\delta > 0$ such that for all $n \geq 1$,

$$\|a\|_p < \delta \text{ implies } 1 - \text{Re } \mathcal{L}_{\mu_n}(a) < \varepsilon.$$

By Lemma 4.2 for $q = p + 1$, for all $n \geq 1$ and all $\sigma > 0$,

$$\int_{\mathbb{S}'} 1 - \exp\left(-\frac{\sigma^2}{2} \|b\|_{-p-1}^2\right) d\mu_n(b) \leq \varepsilon + 2\delta^{-2} \sigma^2 \zeta(2)^d.$$

Then, fix $\sigma > 0$ such that $2\delta^{-2} \sigma^2 \zeta(2)^d = \varepsilon$ and choose $\kappa > 0$ such that $\exp(-\sigma^2 \kappa^2 / 2) \leq \varepsilon$. We need the fact that the set

$$K = \{b \in \mathbb{S}' \mid \|b\|_{-p-1} \leq \kappa\}$$

is strongly compact in \mathbb{S}' . This follows from the fact that in \mathbb{S}' , the strongly closed and bounded subsets are strongly compact. We provide a proof of this fact in

Proposition 6.1 in the appendix. The set $\{b \in \mathbb{S}' \mid \|b\|_{-p} \leq C\}$ is strongly (or weakly) closed as the intersection of the closed sets $\{b \in \mathbb{S}' \mid |\langle b, a \rangle| \leq C\}$, $a \in \mathbb{S}$ with $\|a\|_p \leq 1$. It is strongly bounded since for all bounded set $B \in \mathbb{S}$, with $M = \sup_{a \in B} \|a\|_p < \infty$, we have $q_B(b) \leq M\|b\|_{-p}$.

We have, for all $n \geq 1$,

$$\begin{aligned} \mu_n(K) &\geq \int_{\mathbb{S}'} \exp\left(-\frac{\sigma^2}{2}\|b\|_{-p-1}^2\right) d\mu_n(b) - \int_{K^c} \exp\left(-\frac{\sigma^2}{2}\|b\|_{-p-1}^2\right) d\mu_n(b) \\ &\geq 1 - 2\varepsilon - \varepsilon. \end{aligned}$$

Thus $(\mu_n)_{n \geq 1}$ is tight. \square

Lemma 5.3. *(Re \mathcal{L}_{μ_n}) $_{n \geq 1}$ is equicontinuous.*

Proof. Assume that $(\text{Re } \mathcal{L}_{\mu_n})_{n \geq 1}$ is not equicontinuous. Then, there exist $\varepsilon > 0$ and a sequence of elements $a_k = (a_{k,n})_{n \in \mathbb{N}^d} \in \mathbb{S}$, $k \in \mathbb{N}$, such that $a_k \rightarrow 0$ in \mathbb{S} as $k \rightarrow \infty$, and an increasing sequence of integers $(n_k)_{k \geq 1}$ such that

$$(7) \quad 1 - \text{Re } \mathcal{L}_{\mu_{n_k}}(a_k) \geq \varepsilon, \quad \text{for all } k \geq 1.$$

Since $a_k \rightarrow 0$ in \mathbb{S} , considering a subsequence if necessary, we can assume that $2^k a_k \rightarrow 0$ in \mathbb{S} . That is, for all $p \in \mathbb{N}$, $\lim_{k \rightarrow \infty} \|2^k a_k\|_p = 0$. For a positive integer k_0 to be fixed later, we define the functional $Q : \mathbb{S}' \rightarrow \mathbb{R}_+$ by

$$Q(b) = \sum_{k \geq k_0} \langle b, a_k \rangle^2, \quad b \in \mathbb{S}'.$$

Note that, if $b \in \mathbb{S}'$, there exists $p \geq k_0$ such that $b \in \mathbb{S}_{-p}$ and thus

$$Q(b) \leq \sum_{k=k_0}^p \langle b, a_k \rangle^2 + \|b\|_{-p} \sum_{k \geq p+1} \|a_k\|_p^2 < \infty$$

because $\|a_k\|_p^2 = o(2^{-2k})$. This proves that Q is well-defined. Further, Q is continuous at 0. Indeed, since $2^k a_k \rightarrow 0$, the set $B = \{2^k a_k \mid k \in \mathbb{N}\}$ is bounded in \mathbb{S} , and for all $\eta > 0$, the neighborhood $V_\eta = \{b \in \mathbb{S}' \mid q_B(b) < \eta\}$ of 0 in \mathbb{S}' satisfies that for all $b \in V_\eta$, $|\langle b, a_k \rangle| \leq \eta 2^{-k}$, and thus

$$|Q(b) - Q(0)| = Q(b) \leq \eta^2 \sum_{k \geq k_0} 2^{-k} \leq \eta^2.$$

Next, Q is negative-definite, i.e. for all $m \in \mathbb{N}$, all $c_1, \dots, c_m \in \mathbb{C}$ such that $\sum_{i=1}^m c_i = 0$, and all $b^1, \dots, b^m \in \mathbb{S}'$,

$$\sum_{i=1}^m \sum_{j=1}^m c_i c_j Q(b^i - b^j) \leq 0.$$

Indeed, it is sufficient to notice that

$$\sum_{i=1}^m \sum_{j=1}^m c_i c_j Q(b^i - b^j) = -2 \left(\sum_{i=1}^m c_i \sum_{k \geq k_0} \langle b^i, a_k \rangle \right)^2.$$

The functional $U : \mathbb{S}' \rightarrow [0, 1]$, defined by

$$U(b) = \exp(-Q(b)), \quad b \in \mathbb{S}'.$$

is then positive-definite (see [1, Chap. 3, Theorem 2.2]), continuous at 0 and $U(0) = 1$. Hence, by Theorem 4.3 there exists a Borel probability measure ν on \mathbb{S} such that $\mathcal{L}_\nu = U$.

Note that, since it can be written as the limit of the measurable maps $(b, a) \mapsto \sum_{n \in \Gamma_N} a_n b_n$ as $N \rightarrow \infty$, the bilinear map $\langle \cdot, \cdot \rangle$ is $\mathcal{B}(\mathbb{S}') \times \mathcal{B}(\mathbb{S})$ -measurable. Using Fubini's theorem, for all $n \in \mathbb{N}$, we can write

$$\int_{\mathbb{S}'} U(b) d\mu_n(b) = \int_{\mathbb{S}'} \int_{\mathbb{S}} e^{i\langle b, a \rangle} d\nu(a) d\mu_n(b) = \int_{\mathbb{S}} \mathcal{L}_{\mu_n}(a) d\nu(a).$$

But, by dominated convergence,

$$\lim_{n \rightarrow \infty} \int_{\mathbb{S}} \mathcal{L}_{\mu_n}(a) d\nu(a) = \int_{\mathbb{S}} \mathcal{L}_\mu(a) d\nu(a).$$

Again, by Fubini's theorem, $\int_{\mathbb{S}} \mathcal{L}_\mu(a) d\nu(a) = \int_{\mathbb{S}'} U(b) d\mu(b)$, and thus,

$$\lim_{n \rightarrow \infty} \int_{\mathbb{S}'} U(b) d\mu_n(b) = \int_{\mathbb{S}'} U(b) d\mu(b).$$

Now, recalling the definition of U , we can fix k_0 such that

$$\int_{\mathbb{S}'} 1 - U(b) d\mu(b) < \varepsilon/2,$$

and find $N \in \mathbb{N}$ such that for all $n \geq N$,

$$\int_{\mathbb{S}'} 1 - U(b) d\mu_n(b) < \varepsilon.$$

As a consequence, for all $n \geq N$, and $k \geq k_0$,

$$\int_{\mathbb{S}'} 1 - \exp(-\langle b, a_k \rangle^2) d\mu_n(b) \leq \int_{\mathbb{S}'} 1 - U(b) d\mu_n(b) < \varepsilon.$$

We infer that for all $n \geq N$, and $k \geq k_0$,

$$1 - \operatorname{Re} \mathcal{L}_{\mu_n}(a_k) = \int_{\mathbb{S}'} 1 - \cos(-\langle b, a_k \rangle) d\mu_n(b) < M\varepsilon,$$

where $M = \sup_{x \geq 0} \frac{1 - \cos(x)}{1 - \exp(-x^2)} < \infty$. This contradicts (7). From Lemma 5.2, we infer that $(\mu_n)_{n \geq 1}$ is tight in $\mathcal{P}(\mathbb{S}')$. \square

6. APPENDIX

The goal of the appendix is to prove the following proposition.

Proposition 6.1. *In \mathbb{S}' , the strongly closed and strongly bounded subsets are strongly compact.*

This proposition tells that \mathbb{S}' is a Montel space. See Remark 6.6 at the end. We provide a self-contained proof here, which consists of several steps.

Proposition 6.2. *In \mathcal{S} (and hence in \mathbb{S}), the bounded subsets are relatively compact.*

Proof. As \mathcal{S} is metrizable, compactness and sequential compactness are equivalent. Let $(f_n)_{n \in \mathbb{N}}$ be a bounded sequence in \mathcal{S} . Let $(K_\ell)_{\ell \geq 1}$ be an increasing sequence of compact subsets of \mathbb{R}^d whose union is \mathbb{R}^d . Recall that $(\|\cdot\|_{m,j})_{m \in \mathbb{N}, j \in \mathbb{N}^d}$ in (1) and $(\|\cdot\|_p)_{p \in \mathbb{N}}$ in (4) generate the same topology in \mathcal{S} . For each $m \in \mathbb{N}$ and $j \in \mathbb{N}^d$, the sequence of functions $x \rightarrow (1 + |x|)^m D^j f_n(x)$, $n \in \mathbb{N}$, is uniformly bounded

and equicontinuous on K_1 and hence, by Arzèla–Ascoli’s theorem, there exists a subsequence that converges uniformly on K_1 . By diagonal extraction, we can find a subsequence $(f_{\varphi(n)})_{n \in \mathbb{N}}$ for which $(1 + |\cdot|)^m D^j f_{\varphi(n)}(\cdot)$ converges uniformly on K_1 for all $m \in \mathbb{N}$ and $j \in \mathbb{N}^d$. We can successively repeat the same procedure on each K_ℓ , $\ell \geq 1$, and finally, another diagonal extraction leads to a subsequence that converges with respect to all the semi-norms $\|\cdot\|_{m,j}$, and hence in \mathcal{S} . \square

Lemma 6.3. *If $H \subset \mathcal{S}'$ is weakly closed and weakly bounded, then it is weakly compact.*

Proof. Let $H \subset \mathcal{S}'$ be weakly closed and weakly bounded. Then, for all $n \in \mathbb{N}^d$, the set $H_n = \{b_n \mid (b_k)_{k \in \mathbb{N}^d} \in H\}$ is closed and bounded in \mathbb{R} . By Tychonoff’s theorem, the set $\prod_{n \in \mathbb{N}^d} H_n$ is compact in $\mathbb{R}^{\mathbb{N}^d}$. Therefore H , being a closed subset of $\prod_{n \in \mathbb{N}^d} H_n$, is compact. \square

We say that a set $H \subset \mathcal{S}'$ is *equicontinuous* if for all $\varepsilon > 0$, there exists a neighborhood V of 0 in \mathcal{S} such that, for all $b \in H$ and $a \in V$, $|\langle b, a \rangle| < \varepsilon$.

Lemma 6.4. *If $H \subset \mathcal{S}'$ is equicontinuous, the weak topology and the strong topology are identical on H .*

Proof. It suffices to prove that the induced strong topology on H is weaker than the induced weak topology. For this purpose, we show that for all $\varepsilon > 0$, B a bounded set in \mathcal{S} and $b_0 \in H$, the strongly open set $O := \{b \in H \mid q_B(b - b_0) < \varepsilon\}$ contains a weak neighborhood of b_0 . By equicontinuity, there exists a neighborhood V of 0 in \mathcal{S} such that $|\langle b, a \rangle| < \varepsilon/3$ for all $b \in H$ and $a \in V$. Since B is bounded, by Proposition 6.2, it is relatively compact in \mathcal{S} . Thus, we can find $a_1, \dots, a_n \in B$, such that

$$B \subset \bigcup_{i=1}^n (a_i + V).$$

The set $O' := \{b \in H \mid |\langle b - b_0, a_i \rangle| < \varepsilon/3 \text{ for all } i = 1, \dots, n\}$ is a weak neighborhood of b_0 and we have $O' \subset O$. Indeed, for all $b \in O'$ we can find, for all $a \in B$, $i \in \{1, \dots, n\}$, such that $a - a_i \in V$. Then,

$$|\langle b - b_0, a \rangle| \leq |\langle b, a - a_i \rangle| + |\langle b - b_0, a_i \rangle| + |\langle b_0, a - a_i \rangle| < \varepsilon/3 + \varepsilon/3 + \varepsilon/3.$$

Thus, $q_B(b - b_0) < \varepsilon$ and hence $b \in O$. \square

Lemma 6.5. *If $H \subset \mathcal{S}'$ is strongly bounded, then it is equicontinuous.*

Proof. Let $H \subset \mathcal{S}'$ be strongly bounded. In the sequel, we denote $B_p(\varepsilon) = \{a \in \mathcal{S} \mid \|a\|_p < \varepsilon\}$ for $p \in \mathbb{N}$ and $\varepsilon > 0$. We first show that the set $\{q_{B_{p_0}(1)}(b) \mid b \in H\}$ is bounded for some $p_0 \in \mathbb{N}$. Assume it is not the case. Then, for each $p \in \mathbb{N}$, there exist $a_p \in B_p(1/p)$ and $b_p \in H$ such that $|\langle b_p, a_p \rangle| > p$. Since each a_p belongs to $B_p(1/p)$, the sequence $(a_p)_{p \in \mathbb{N}}$ converges to 0 in \mathcal{S} and thus it is bounded. But then, for the bounded set $B = \{a_p \mid p \in \mathbb{N}\}$, we have $q_B(b_p) > p$ for all $p \in \mathbb{N}$ which is a contradiction with the fact that H is strongly bounded.

Now for such a p_0 , letting $M = \sup_{b \in H} q_{B_{p_0}(1)}(b) < \infty$, we have that for all $\varepsilon > 0$ and for all $a \in B_{p_0}(\varepsilon/M)$,

$$|\langle b, a \rangle| \leq \|b\|'_{p_0} \|a\|_{p_0} \leq M\varepsilon/M = \varepsilon, \text{ for all } b \in H.$$

\square

Proof of Proposition 6.1. Let H be a strongly closed and strongly bounded subset of \mathcal{S}' . In particular, H is weakly bounded and, by Lemma 6.3, its weak closure \bar{H} is weakly compact. Now, by Lemma 6.5, H is equicontinuous and then \bar{H} is also equicontinuous. Hence, strong and weak topology are identical on \bar{H} by Lemma 6.4. As H is strongly closed, it is also weakly closed and we infer that $\bar{H} = H$. Hence H is weakly compact and then strongly compact. \square

Remark 6.6. Proposition 6.2 tells that \mathcal{S} and \mathbb{S} are actually Montel spaces (i.e. a locally convex Hausdorff barreled space in which every closed bounded set is compact). See also [20, Example 10.IV and Proposition 34.4] and [8, Section I.6.5]. Proposition 6.1 tells that \mathcal{S}' and \mathbb{S}' are also Montel spaces. Furthermore, as a consequence of Banach–Steinhaus theorem, any weakly convergent sequence of \mathcal{S}' (or \mathbb{S}') is also strongly convergent (see [20, Corollary 1 of Proposition 34.6] and [8, Section I.6.4]).

Acknowledgements. YW's research was partially supported by the NSA grants H98230-14-1-0318 and H98230-16-1-0322, the ARO grant W911NF-17-1-0006, and Charles Phelps Taft Research Center at University of Cincinnati.

REFERENCES

- [1] Berg, C., Christensen, J. P. R., and Ressel, P. (1984). *Harmonic analysis on semigroups*, volume 100 of *Graduate Texts in Mathematics*. Springer-Verlag, New York. Theory of positive definite and related functions.
- [2] Boulicaut, P. (1973/74). Convergence cylindrique et convergence étroite d'une suite de probabilités de Radon. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 28:43–52.
- [3] Dobrušin, R. L. and Minlos, R. A. (1976). A study of the properties of generalized Gaussian random fields. In *Problems in mechanics and mathematical physics (Russian)*, pages 117–165, 298. Izdat. “Nauka”, Moscow.
- [4] Fageot, J., Amini, A., and Unser, M. (2014). On the continuity of characteristic functionals and sparse stochastic modeling. *J. Fourier Anal. Appl.*, 20(6):1179–1211.
- [5] Fageot, J., Bostan, E., and Unser, M. (2015). Wavelet statistics of sparse and self-similar images. *SIAM J. Imaging Sci.*, 8(4):2951–2975.
- [6] Fernique, X. (1968). Généralisations du théorème de continuité de P. Lévy. *C. R. Acad. Sci. Paris Sér. A-B*, 266:A25–A28.
- [7] Gel'fand, I. M. (1955). Generalized random processes. *Dokl. Akad. Nauk SSSR (N.S.)*, 100:853–856.
- [8] Gel'fand, I. M. and Shilov, G. E. (1968). *Generalized functions. Vol. 2. Spaces of fundamental and generalized functions*. Translated from the Russian by Morris D. Friedman, Amiel Feinstein and Christian P. Peltzer. Academic Press, New York-London.
- [9] Gel'fand, I. M. and Vilenkin, N. Y. (1964). *Generalized functions. Vol. 4*. Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London. Applications of harmonic analysis, Translated from the Russian by Amiel Feinstein.
- [10] Gorostiza, L. G. and Nualart, D. (1994). Nuclear Gel'fand triples on Wiener space and applications to trajectorial fluctuations of particle systems. *J. Funct. Anal.*, 125(1):37–66.

- [11] Holley, R. A. and Stroock, D. W. (1978). Generalized Ornstein-Uhlenbeck processes and infinite particle branching Brownian motions. *Publ. Res. Inst. Math. Sci.*, 14(3):741–788.
- [12] Itô, K. (1983). Distribution-valued processes arising from independent Brownian motions. *Math. Z.*, 182(1):17–33.
- [13] Kallenberg, O. (1997). *Foundations of modern probability*. Probability and its Applications (New York). Springer-Verlag, New York.
- [14] Kallianpur, G. and Xiong, J. (1995). *Stochastic differential equations in infinite-dimensional spaces*. Institute of Mathematical Statistics Lecture Notes—Monograph Series, 26. Institute of Mathematical Statistics, Hayward, CA.
- [15] Kuo, H.-H. (1996). *White noise distribution theory*. Probability and Stochastics Series. CRC Press, Boca Raton, FL.
- [16] Meyer, P.-A. (1995). Le théorème de continuité de P. Lévy sur les espaces nucléaires (d’après X. Fernique). In *Séminaire Bourbaki, Vol. 9*, pages Exp. No. 311, 509–522. Soc. Math. France, Paris.
- [17] Minlos, R. A. (1959). Generalized random processes and their extension in measure. *Trudy Moskov. Mat. Obšč.*, 8:497–518.
- [18] Nelson, E. (1973). Quantum fields and Markoff fields. In *Partial differential equations (Proc. Sympos. Pure Math., Vol. XXIII, Univ. California, Berkeley, Calif., 1971)*, pages 413–420. Amer. Math. Soc., Providence, R.I.
- [19] Reed, M. and Simon, B. (1980). *Methods of modern mathematical physics. I*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, second edition. Functional analysis.
- [20] Trèves, F. (1967). *Topological vector spaces, distributions and kernels*. Academic Press, New York-London.

HERMINE BIERMÉ, LABORATOIRE DE MATHÉMATIQUES ET APPLICATIONS UMR CNRS 7348, UNIVERSITÉ DE POITIERS, BOULEVARD MARIE ET PIERRE CURIE 86962 FUTUROSCOPE CHASSENEUIL CEDEX, FRANCE.

E-mail address: `hermine.bierme@math.univ-poitiers.fr`

OLIVIER DURIEU, LABORATOIRE DE MATHÉMATIQUES ET PHYSIQUE THÉORIQUE, UMR-CNRS 7350, FÉDÉRATION DENIS POISSON, FR-CNRS 2964, UNIVERSITÉ FRANÇOIS-RABELAIS DE TOURS, PARC DE GRANDMONT, 37200 TOURS, FRANCE.

E-mail address: `olivier.durieu@lmpt.univ-tours.fr`

YIZAO WANG, DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF CINCINNATI, 2815 COMMONS WAY, ML-0025, CINCINNATI, OH, 45221-0025.

E-mail address: `yizao.wang@uc.edu`