

# ISOMETRIES FOR THE CARATHÉODORY METRIC

MARCO ABATE AND JEAN-PIERRE VIGUÉ

## 1. INTRODUCTION

The following problem has been studied by many authors. Let  $D_1$  and  $D_2$  be two bounded domains in complex Banach spaces and let  $f: D_1 \rightarrow D_2$  be a holomorphic map such that  $f'(a)$  is a surjective isometry for the Carathéodory infinitesimal metric at a point  $a$  of  $D_1$ . The problem is to know whether  $f$  is an analytic isomorphism of  $D_1$  onto  $D_2$ . For example, J.-P. Vigué [8] proved this is the case when  $D_1$  and  $D_2$  are two bounded domains in  $\mathbb{C}^n$  and  $D_1$  is convex. Similar results have been obtained when  $D_2$  is convex using the Kobayashi infinitesimal metric (I. Graham [3] and L. Belkhchicha [1]). We have to remark that all these results are based on the theorem of L. Lempert ([5] et [6]; one can also consult M. Jarnicki and P. Pflug [4]) on the equality of Kobayashi and Carathéodory metrics on a bounded convex domain in  $\mathbb{C}^n$ . J.-P. Vigué [9] proved the first results on this subject in the case of bounded domains in complex Banach spaces.

Now, we can study the same problem dropping the hypothesis that  $f'(a)$  is surjective. So, we only suppose that  $f'(a)$  is an isometry for the Carathéodory infinitesimal metric. Does this imply that  $f(D_1)$  is a complex analytic closed submanifold of  $D_2$  and that  $f$  is an analytic isomorphism of  $D_1$  onto  $f(D_1)$ ?

Some results have been obtained by J.-P. Vigué [10] and P. Mazet [7] assuming that  $D_1$  and  $D_2$  are open unit balls in complex Banach spaces, that  $a = 0$ , and that the image of  $f'(0)$  contains enough complex extremal points of the boundary of  $D_2$ . Under these hypotheses they proved that  $f$  is linear equal to  $f'(0)$ . This result shows that  $f(D_1)$  is an analytic submanifold of  $D_2$  and that  $f$  is an analytic isomorphism of  $D_1$  onto  $f(D_1)$ .

Of course, if we do not suppose the existence of complex extremal points in the image of  $f'(0)$ , the map  $f$  has no reason to be linear. However, one can hope that  $f(D_1)$  still is a complex analytic submanifold of  $D_2$ . In this paper we shall be able to prove such a result for maps of unit balls of complex Banach spaces, under some additional hypotheses on the Banach spaces involved.

The authors would like to thank the referee for his/her useful remarks.

## 2. THE MAIN RESULTS

We shall prove the following theorem:

**Theorem 1.** *Let  $(E_j, \|\cdot\|_j)$  be complex Banach spaces and let  $B_j = \{x \in E_j \mid \|x\|_j < 1\}$ , for  $j = 1, 2$ . Let  $f: B_1 \rightarrow B_2$  be a holomorphic mapping with  $f(0) = 0$  and  $\|f'(0)(X)\|_2 = \|X\|_1$  for all  $X \in E_1$ . Then the following statements are equivalent :*

- (1) *there exists a direct decomposition  $E_2 = f'(0)(E_1) \oplus F$  such that the corresponding projection  $\pi: E_2 \rightarrow f'(0)(E_1)$  has norm 1;*
- (2)  *$f(B_1)$  is a closed complex direct submanifold of  $B_2$ , the map  $f$  is a biholomorphism of  $B_1$  onto  $f(B_1)$ , and there exists a holomorphic retraction of  $B_2$  onto  $f(B_1)$ .*

To apply this theorem, we give the following definition:

**Definition 1.** *We say that a pair  $(E_1, E_2)$  of complex Banach spaces has the property (V) if for every linear isometry  $L: E_1 \rightarrow E_2$  there exists a direct decomposition  $E_2 = L(E_1) \oplus F$  such that the corresponding projection  $\pi: E_2 \rightarrow L(E_1)$  has norm 1.*

From theorem 1 and definition 1, we deduce the following

**Theorem 2.** *Assume that the pair  $(E_1, E_2)$  of complex Banach spaces has the property (V), and let  $B_1$  and  $B_2$  be their open unit balls. Let  $f: B_1 \rightarrow B_2$  be a holomorphic map such that*

- (1)  *$f(0) = 0$ , and  $f'(0)$  is an isometry for the Carathéodory infinitesimal metric,*  
or
- (2)  *$B_1$  and  $B_2$  are homogeneous, and there exists  $a \in B_1$  such that  $f'(a)$  is an isometry for the Carathéodory infinitesimal metric.*

*Then  $f(B_1)$  is a closed complex direct submanifold of  $B_2$ , the map  $f$  is a biholomorphism of  $B_1$  onto  $f(B_1)$ , and there exists a holomorphic retraction of  $B_2$  onto  $f(B_1)$ .*

Now, we clearly need examples of pairs of complex Banach spaces satisfying property (V). The first (easy) example is given by Hilbert spaces.

**Proposition 1.** *Let  $E_2$  a complex Hilbert space. Then, for every complex Banach space  $E_1$  the pair  $(E_1, E_2)$  has property (V).*

More interesting is the following theorem:

**Theorem 3.** *Let  $I$  be a set and let  $l^\infty(I)$  be the complex Banach space of bounded sequences indexed by  $I$ , with the usual norm. Let  $E_2$  be any Banach space. Then, the pair  $(l^\infty(I), E_2)$  has property (V).*

Other pairs enjoying property (V) can be constructed using suitable subspaces of  $l^\infty(I)$ . For instance, let  $c_0(I) \subset l^\infty(I)$  be the subspace given by the elements  $(a_i)_{i \in I} \in l^\infty(I)$  such that for every  $\varepsilon > 0$  there exists a finite subset  $K \subseteq I$  so that  $|a_i| < \varepsilon$  when  $i \notin K$ . Then:

**Theorem 4.** *For any sets  $I, J$  the pair  $(c_0(I), c_0(J))$  has property (V).*

Applying Theorem 2 and 3 with  $I$  finite, we get in particular a new result in the finite-dimensional case:

**Corollary 1.** *Let  $f: \Delta^n \rightarrow D$  be a holomorphic map between a polydisk  $\Delta^n \subset \mathbb{C}^n$  and an open convex circular bounded domain  $D \subset \mathbb{C}^N$  (i.e.,  $D$  is the unit ball for a suitable norm in  $\mathbb{C}^N$ ). We also assume  $n \leq N$ , and that  $D$  is homogeneous (for instance,  $D = \Delta^N, B^N$  or a bounded symmetric domain). Assume that there exists  $a \in \Delta^n$  such that  $f'(a)$  is an isometry for the Carathéodory infinitesimal metrics. Then  $f(\Delta^n)$  is a closed complex submanifold of  $D$ , the map  $f$  is a biholomorphism onto its image, and  $f(\Delta^n)$  is a holomorphic retract of  $D$ .*

Before proving these results, we need to recall some facts.

### 3. SOME CLASSICAL RESULTS

The definition and the main properties of Carathéodory and Kobayashi infinitesimal metrics  $E_D$  and  $F_D$  on a bounded domain  $D$  are given in the book of T. Franzoni et E. Vesentini [2] (see also the book of M. Jarnicki and P. Pflug [4]).

Let  $B$  be the open unit ball of a complex Banach space  $E$ . It is well known that

$$E_B(0, x) = F_B(0, x) = \|x\|.$$

Furthermore, every biholomorphism  $f: D_1 \rightarrow D_2$  between domains in complex Banach spaces is an isometry for the Carathéodory and Kobayashi infinitesimal metrics.

Finally, let us recall that, the open unit balls  $B$  of the complex Banach spaces  $c_0(I)$  and  $l^\infty(I)$  are homogeneous. Indeed, it is easy to check that, for every  $a \in B$ , the map  $\varphi_a: B \rightarrow B$  given by

$$\forall i \in I \quad \varphi_a(f)_i = \frac{f_i + a_i}{1 + \overline{a_i} f_i},$$

is an analytic automorphism of  $B$ .

Another example of homogeneous unit ball is given by the open unit ball  $B$  of the space  $C(S, \mathbb{C})$  of continuous complex functions on a

compact space  $S$ , because for every  $a \in B$  the map  $\varphi_a: B \rightarrow B$  given by

$$\varphi_a(f) = \frac{f + a}{1 + \bar{a}f}$$

is a biholomorphism of  $B$ .

#### 4. PROOF OF THEOREMS 1 AND 2

To begin, let us prove theorem 1.

*Proof of Theorem 1.* First, if  $r: B_2 \rightarrow f(B_1)$  is a holomorphic retraction,  $r'(0)$  is a projection of norm  $\leq 1$  for the Carathéodory infinitesimal metrics, and,

as the Carathéodory infinitesimal metric at the origin is equal to the given norm, we get  $\|r'(0)\| = 1$ . This proves that (2) implies (1).

To prove that (1) implies (2), let us consider

$$\varphi = \pi \circ f: B_1 \rightarrow f'(0)(E_1).$$

We have  $\varphi(0) = 0$ ,  $\varphi(B_1) \subseteq f'(0)(E_1) \cap B_2$  (because  $\pi$  has norm 1), and  $\varphi'(0) = \pi \circ f'(0) = f'(0)$ . So  $\varphi'(0)$  is a linear isometry from  $E_1$  onto  $f'(0)(E_1)$ . Using Cartan's uniqueness theorem (see [2]), one easily proves that  $\varphi$  is a linear isometry from  $B_1$  onto  $B'_2 = f'(0)(E_1) \cap B_2$ .

Finally, let  $\psi: B'_2 \rightarrow F$  be defined by

$$\psi(y) = (\text{id} - \pi)(f(\varphi^{-1}(y))).$$

Then the set  $f(B_1)$  is the graph of  $\psi$ , the map  $(\pi, \psi \circ \pi): B_2 \rightarrow f(B_1)$  is a holomorphic retraction of  $B_2$  onto  $f(B_1)$ , and  $\varphi^{-1} \circ \pi|_{f(B_1)}: f(B_1) \rightarrow B_1$  is a holomorphic inverse of  $f$ , and the theorem is proved.  $\square$

Now, we can prove Theorem 2.

*Proof of Theorem 2.* First, let us remark that, in case (2), by pre-composing  $f$  with an analytic automorphism of  $B_1$  and post-composing it with an analytic automorphism of  $B_2$ , we can assume that  $f(0) = 0$  and that 0 is precisely the point  $a$  such that  $f'(0)$  is an isometry for the Carathéodory infinitesimal metrics. Thus without loss of generality in both cases we can assume that  $f'(0)$  is an isometry for the norms of  $E_1$  and  $E_2$ .

Since  $(E_1, E_2)$  satisfies the property (V), there exists a direct decomposition  $E_2 = f'(0)(E_1) \oplus F$  such that the corresponding projection  $\pi: E_2 \rightarrow f'(0)(E_1)$  has norm 1 and we can apply Theorem 1.  $\square$

## 5. EXAMPLES OF PAIR OF BANACH SPACES WITH PROPERTY (V)

Now we have to give examples of pair of complex Banach spaces satisfying property (V). Proposition 1 (the case of Hilbert spaces) is easy and left as an exercise. Let us now give the

*Proof of Theorem 3.* We suppose that  $E_1 = l^\infty(I)$  and we consider an isometry  $L: l^\infty(I) \rightarrow E_2$ . Let  $G: L(E_1) \rightarrow l^\infty(I)$  be the inverse of  $L$ . So,  $G$  is a linear map of norm 1; for every  $i \in I$ , let  $G_i$  be the  $i$ -component of  $G$ . Then  $G_i$  is a linear form from  $L(E_1)$  to  $\mathbb{C}$  of norm 1. By the Hahn-Banach Theorem, we can extend  $G_i$  to a linear form  $H_i: E_2 \rightarrow \mathbb{C}$  still of norm 1. Setting  $H = (H_i)_{i \in I}$  we obtain a linear map  $H: E_2 \rightarrow l^\infty(I)$  of norm 1 extending  $G$ . Then it is clear that  $L \circ H$  is a projection of  $E_2$  onto  $L(l^\infty(I))$  of norm 1, and taking  $F = \text{Ker}(L \circ H)$  the theorem is proved.  $\square$

*Proof of Theorem 4.* Let  $L: c_0(I) \rightarrow c_0(J)$  be an isometry, and let  $(e^k)_{k \in I}$  be the canonical basis of  $c_0(I)$ . Since  $L$  is an isometry, for every  $k \in I$  there exists  $j(k) \in J$  such that  $|L(e^k)_{j(k)}| = 1$ . Now, if we consider an element  $v = (v_i)_{i \in I}$  of  $c_0(I)$  such that  $v_k = 0$ , then  $L(v)_{j(k)} = 0$ . In fact, suppose that  $L(v)_{j(k)} \neq 0$ . For  $\lambda \in \mathbb{C}$  small enough, we have  $\|e^k + \lambda v\| = 1$ . But

$$L(e^k + \lambda v)_{j(k)} = L(e^k)_{j(k)} + \lambda L(v)_{j(k)} = e^{i\theta} + \lambda L(v)_{j(k)}.$$

Therefore if  $L(v)_{j(k)} \neq 0$ , there exists  $\lambda \in \mathbb{C}$  small enough such that the modulus of  $L(e^k + \lambda v)_{j(k)}$  is greater than 1, and thus  $\|L(e^k + \lambda v)\| > 1$ , contradiction. It follows that the map  $k \mapsto j(k)$  is injective.

Let  $M = \{j(k) \mid k \in I\} \subseteq J$ , and let  $\pi: c_0(J) \rightarrow c_0(M)$  be the canonical projection. The previous argument shows

that  $\pi \circ L(e^k) = \lambda_k e^{j(k)}$  with  $|\lambda_k| = 1$  for all  $k \in I$ ; it is then easy

to check that  $\varphi = \pi \circ L: c_0(I) \rightarrow c_0(M)$  is a linear surjective isometry, and that  $L \circ \varphi^{-1} \circ \pi: c_0(J) \rightarrow L(c_0(I))$  is a linear projection of norm 1 of  $c_0(J)$  onto  $L(c_0(I))$ , as required.  $\square$

It might be interesting to remark that the same proof yields that a pair  $(E_1, E_2)$  of complex Banach spaces satisfies property (V) if each  $E_j$  has a Schauder basis  $(e_j^k)$  such that

$$\left\| \sum_k \lambda_k e_j^k \right\|_{E_j} = \sup_k |\lambda_k|.$$

## 6. FINAL REMARKS

Not all pairs of complex Banach spaces have property (V); so we do not know whether Theorem 2 holds in general.

For example, the Banach spaces  $c_0(\mathbb{N})$  is not complemented in  $l^\infty(\mathbb{N})$ , and so the pair  $(c_0(\mathbb{N}), l^\infty(\mathbb{N}))$  does not have property (V).

It is also possible to build finite dimensional examples. Take  $E_2 = (\mathbb{C}^3, \|\cdot\|_\infty)$ , so that the unit ball of  $E_2$  is the open polydisk  $\Delta^3$ . If  $L: \mathbb{C}^2 \rightarrow \mathbb{C}^3$  is given by  $L(x, y) = (x, y, x + y)$ , then  $B = L^{-1}(\Delta^3)$  is the open unit ball in  $\mathbb{C}^2$  for a norm  $\|\cdot\|$ ; set  $E_1 = (\mathbb{C}^2, \|\cdot\|)$ . We claim that the pair  $(E_1, E_2)$  does not satisfy property (V). By construction,  $L: E_1 \rightarrow E_2$  is a linear isometry. The set  $(1, 0, 1) + (\{0\} \times \Delta \times \{0\})$  is contained in the boundary of  $\Delta^3$ . Since  $(1, 0, 1) \in L(\partial B)$ , it is easy to check that if there exists a projection  $\pi$  of norm 1 from  $\mathbb{C}^3$  onto  $L(\mathbb{C}^2)$ , then  $\pi$  must vanish on  $\{0\} \times \mathbb{C} \times \{0\}$ . Considering the point  $(0, 1, 1)$ , we analogously see that  $\pi$  must vanish on  $\Delta \times \{0\} \times \{0\}$ ; and thus such a projection cannot exist.

## REFERENCES

- [1] L. Belkhchicha. Caractérisation des isomorphismes analytiques sur la boule-unité de  $\mathbb{C}^n$  pour une norme. *Math. Z.* **215** (1994), 129–141.
- [2] T. Franzoni and E. Vesentini. Holomorphic maps and invariant distances. *Notas de Matematica [Mathematical Notes]*, 69. North-Holland Publishing Co., Amsterdam-New York, 1980.
- [3] I. Graham. Holomorphic mappings into strictly convex domains which are Kobayashi isometries at one point. *Proc. Amer. Math. Soc.* **105** (1989), 917–921.
- [4] M. Jarnicki and P. Pflug. Invariant distances and metrics in complex analysis. *de Gruyter Expositions in Mathematics*, 9. Walter de Gruyter and Co., Berlin, 1993.
- [5] L. Lempert. La métrique de Kobayashi et la représentation des domaines sur la boule. *Bull. Soc. Math. France* **109** (1981), 427–474.
- [6] L. Lempert. Holomorphic retracts and intrinsic metrics in convex domains. *Anal. Math.* **8** (1982), 257–261.
- [7] P. Mazet. Principe du maximum et lemme de Schwarz à valeurs vectorielles. *Canad. Math. Bull.* **40** (1997), 356–363.
- [8] J.-P. Vigué. Sur les points fixes d'applications holomorphes. *C. R. Acad. Sci. Paris Sér. I Math.* **303** (1986), 927–930.
- [9] J.-P. Vigué. Sur la caractérisation des isomorphismes analytiques entre domaines bornés d'un espace de Banach complexe. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **21** (1994), 145–155.
- [10] J.-P. Vigué. Un lemme de Schwarz pour les boules-unités ouvertes. *Canad. Math. Bull.* **40** (1997), no. 1, 117–128.

Marco Abate, Dipartimento di Matematica, Università di Pisa, Largo Pontecorvo 5, 56127 Pisa, Italy.

e-mail : [abate@dm.unipi.it](mailto:abate@dm.unipi.it)

J.-P. V., LMA, Université de Poitiers, CNRS, Mathématiques, SP2MI,  
BP 30179, 86962 FUTUROSCEPE.

e-mail : [vigue@math.univ-poitiers.fr](mailto:vigue@math.univ-poitiers.fr)